

Advice following treatment with Botox

We have already discussed the possible side effects of using botox, before you were given your first injection. These are as follows:

- bruising, discomfort at the site of injection - this should be mild and should pass quickly
- MILD headache (within the first 24 hours).
- drooping of the brows (if used in the forehead area)
- drooping of the eyelid
- allergy: you should be aware of the possibility of allergy with any new medication. In the case of botox allergy is very rare, but worth bearing in mind.

If you have *any* concerns, please contact me via email in the first instance - on sageaesthetic@me.com, and I will contact you ASAP (usually within a couple of hours) to arrange a time for you to be reviewed in the clinic.

To maximise the effectiveness of your Botox:

Botox injected into muscles: for the first 24 hours (but not when you are sleeping!), try to make repeated facial expressions - this will allow the Botox to enter the “active” nerve ending. For example, frown / raise eyebrows / smile every waking hour.

For Botox injected into the skin: avoid applying deodorant and skin preparations containing alcohol to the affected area, as the alcohol can destroy the botox protein.

EMERGENCY ACTIONS:

If you think you are experiencing a severe allergic reaction, please **dial 999 immediately**. Symptoms of this might include: Hives/nettle rash, swelling (including swelling of the face and throat), wheezing, feeling faint, shortness of breath.

If you are having difficulty in breathing, speaking or swallowing, or have noticed that food is accidentally going ‘down the wrong way’ (into your lungs, causing coughing/choking when eating), or if you have muscle weakness (not including the target muscles of course!), then please either contact me as above by email, or if more urgent then call an ambulance on 999.

Any questions... just ask!

