

Literature: Weyl 1946 Classical groups, pages 177–229;

Muirhead 1982, Aspects of multivariate statistical theory, chapter 2, pages 50–72;

Katz, Sarnak, Chapter 5, pages 107–121;

1.1 $U(n)$.

For a matrix A , denote

$$\delta A = A^{-1} dA,$$

whatever this means.

For $A \in U(n)$, represent

$$A = U \Lambda U^*, \quad (1.1)$$

where

$$\Lambda = \text{diag}(e^{i\varphi_1}, \dots, e^{i\varphi_n}), \quad 0 < \varphi_1 < \dots < \varphi_n < 2\pi, \quad (1.2)$$

and $H^* \in U(n)$ is fixed by the condition that the in every column, the first (from up down over rows) non-zero column is positive. (Otherwise representation (1.1) is not unique, $U \rightarrow UD$ with diagonal D). Furthermore, U is parametrized by $n^2 - n = l$ real parameters, Λ is parametrized by n parameters. The representation (1.1) is unique for $A \in U(n)$ with all different eigenvalues, and hence we can consider it as a change of variables. Now, since $UU^* = \mathbf{1}$, and hence $\delta U = -\delta U^*$ is skew-symmetric, we obtain by formal manipulations

$$\delta B := U^{-1} \delta A U = (\Lambda^{-1} \delta U \Lambda - \delta U) + \delta \Lambda. \quad (1.3)$$

From (1.3), considering separately diagonal and off-diagonal terms,

$$\begin{aligned} \delta b_{jj} &= i d\varphi_j, \\ \delta b_{jk} &= (e^{i(\varphi_k - \varphi_j)} - 1) \delta u_{jk}, \quad j \neq k. \end{aligned}$$

Now, after all matrix multiplications, we write δB in a row of n^2 real elements:

$$\delta b_{11}, \dots, \delta b_{nn}, \delta b_{12}, \delta b_{21}, \dots, \delta b_{n-1,n}, \delta b_{n,n-1},$$

and take a (wedge) product of them (as of differentials; we obtain an n^2 dimensional differential). Writing δB as a (scalar rather than matrix) n^2 dimensional differential, and ignoring sign,

$$\delta B = \prod_{j=1}^n d\varphi_j \prod_{j \neq k} (e^{i\varphi_j} - e^{i\varphi_k}) \delta b_{j,k} = \prod_{j \neq k} (e^{i\varphi_j} - e^{i\varphi_k}) \prod_{j=1}^n d\varphi_j d\omega_U.$$

Now, when we integrate over $U(n)$ (or average over group) a function that depends only on eigenvalues, first of all we integrate only over matrices A with all distinct eigenvalues (other A form a sub-manifold of smaller dimension, and hence do not contribute in the integral)

$$\begin{aligned} \int_{U(n)} f(A) d\text{Haar} &= \int_{U(n)} f(A) A^{-1} dA = \int_{U(n)} f(A) \delta A = \int_{U(n)} f(A) \delta B = \\ &= \int_{U(n)} f(\varphi_1, \dots, \varphi_n) \prod_{j \neq k} (e^{i\varphi_j} - e^{i\varphi_k}) \prod_{j=1}^n d\varphi_j d\omega_U = \\ &= \left(\frac{1}{n!} \int_{[0, 2\pi]^n} f(\varphi_1, \dots, \varphi_n) \prod_{j \neq k} |e^{i\varphi_j} - e^{i\varphi_k}| \prod_{j=1}^n d\varphi_j \right) \cdot \text{Vol}(U) \end{aligned}$$

where we separated contribution in the integral coming from φ_j , and from U . The factor $n!$ appeared since we changed integral from over $0 < \varphi_1 < \dots < \varphi_n < 2\pi$ to over $\varphi_j \in [0, 2\pi]$. To have it normalized to 1, probably when we integrate over $d\varphi$ over $[0, 2\pi]$, it is better to divide by 2π . However, Katz Sarnak, chapter 5, divide by 2π not only in the case $U(n)$, when integrate over $[0, 2\pi]$, but also in the cases $O(2n)$, when they integrate over $[0, \pi]$, but still divide by 2π . Furthermore, normalizing, we need to divide by analogous expression with $f = 1$. In that way we get rid of $Vol(U)$.

$$\int_{U(n)} f(A) dHaar = \left(\frac{1}{n!} \int_{[0, 2\pi]^n} f(\varphi_1, \dots, \varphi_n) \prod_{j \neq k} |e^{i\varphi_j} - e^{i\varphi_k}| \prod_{j=1}^n \frac{d\varphi_j}{2\pi} \right). \quad (1.4)$$

1.2 $O^+(2n)$.

We will take $n = 2$, so we consider $O^+(4)$. Here (1.1) and (1.3) still hold,

$$A = U\Lambda U^*, \quad \delta B := U^{-1} \delta A U = (\Lambda^{-1} \delta U \Lambda - \delta U) + \delta \Lambda,$$

but Λ changes: instead of (1.2), we have

$$\Lambda = \begin{bmatrix} \begin{bmatrix} \cos \varphi_1 & -\sin \varphi_1 \\ \sin \varphi_1 & \cos \varphi_1 \end{bmatrix} & 0 \\ 0 & \begin{bmatrix} \cos \varphi_2 & -\sin \varphi_2 \\ \sin \varphi_2 & \cos \varphi_2 \end{bmatrix} \end{bmatrix}, \quad 0 < \varphi_1 < \varphi_2 < \pi. \quad (1.5)$$

Again, U should be fixed somehow to ensure uniqueness. It is parametrized by

$$\frac{2n(2n-1)}{2} - n = 2n(n-1)$$

parameters. Λ is parametrized by n real parameters. What is δU ? It is a skew-symmetric matrix, parametrized by low-diagonal elements, whose are $2n(2n-1)/2$, where we need to subtract n , since we fixed somehow U . We have

$$\delta U = \begin{bmatrix} \begin{bmatrix} 0 & -\delta u_{21} \\ \delta u_{21} & 0 \end{bmatrix} & \begin{bmatrix} -\delta u_{31} & -\delta u_{41} \\ -\delta u_{32} & -\delta u_{42} \end{bmatrix} \\ \begin{bmatrix} \delta u_{31} & \delta u_{32} \\ \delta u_{41} & \delta u_{42} \end{bmatrix} & \begin{bmatrix} 0 & -\delta u_{43} \\ \delta u_{43} & 0 \end{bmatrix} \end{bmatrix}, \quad \text{where now the scalar meaning is } \delta U = \prod_{j>k} \delta u_{jk}.$$

What is $\Lambda^{-1} \delta U \Lambda$? But first, what is

$$\delta \Lambda = \Lambda^{-1} d\Lambda = \begin{bmatrix} \begin{bmatrix} 0 & -d\varphi_1 \\ d\varphi_1 & 0 \end{bmatrix} & 0 \\ 0 & \begin{bmatrix} 0 & -d\varphi_2 \\ d\varphi_2 & 0 \end{bmatrix} \end{bmatrix}, \quad \text{with the scalar meaning } \delta \Lambda = d\varphi_1 d\varphi_2.$$

Second, diagonal block terms in $\Lambda^{-1} \delta U \Lambda$ are conjugated by the same φ :

$$\begin{bmatrix} \cos \varphi_1 & \sin \varphi_1 \\ -\sin \varphi_1 & \cos \varphi_1 \end{bmatrix} \begin{bmatrix} 0 & -\delta u_{21} \\ \delta u_{21} & 0 \end{bmatrix} \begin{bmatrix} \cos \varphi_1 & -\sin \varphi_1 \\ \sin \varphi_1 & \cos \varphi_1 \end{bmatrix} = \begin{bmatrix} 0 & -\delta u_{21} \\ \delta u_{21} & 0 \end{bmatrix}, \quad \text{with meaning } \delta_{21}.$$

Third, off-diagonal block terms in $\Delta^{-1} \delta U \Delta$ are conjugated by different φ :

$$\begin{bmatrix} \cos \varphi_2 & \sin \varphi_2 \\ -\sin \varphi_2 & \cos \varphi_2 \end{bmatrix} \begin{bmatrix} \delta u_{31} & \delta u_{32} \\ \delta u_{41} & \delta u_{42} \end{bmatrix} \begin{bmatrix} \cos \varphi_1 & -\sin \varphi_1 \\ \sin \varphi_1 & \cos \varphi_1 \end{bmatrix} - \begin{bmatrix} \delta u_{31} & \delta u_{32} \\ \delta u_{41} & \delta u_{42} \end{bmatrix} =: \begin{bmatrix} \delta v_1 & \delta v_2 \\ \delta v_3 & \delta v_4 \end{bmatrix}. \quad (1.6)$$

The scalar meaning of $\begin{bmatrix} \delta v_1 & \delta v_2 \\ \delta v_3 & \delta v_4 \end{bmatrix}$ itself is just the (wedge) product of all the elements: $\delta v_1 \delta v_2 \delta v_3 \delta v_4$.

Formula (1.6) can be rewritten, elementwise, as (we denote for shortness $c = \cos, s = \sin$)

$$\begin{bmatrix} \begin{bmatrix} c_1 c_2 - 1 & s_1 c_2 \\ -s_1 c_2 & c_1 c_2 - 1 \end{bmatrix} & \begin{bmatrix} c_1 s_2 & s_1 s_2 \\ -s_1 s_2 & c_1 s_2 \end{bmatrix} \\ \begin{bmatrix} -c_1 s_2 & -s_1 s_2 \\ s_1 s_2 & -c_1 s_2 \end{bmatrix} & \begin{bmatrix} c_1 c_2 - 1 & s_1 c_2 \\ -s_1 c_2 & c_1 c_2 - 1 \end{bmatrix} \end{bmatrix} \cdot \begin{bmatrix} \delta u_{31} \\ \delta u_{32} \\ \delta u_{41} \\ \delta u_{42} \end{bmatrix}.$$

We need to take a covariant product of those differential forms, or, equivalently, take the determinant of the matrix, which equals

$$\det = 4(\cos \varphi_1 - \cos \varphi_2)^2.$$

Hence, the contribution from off-diagonal blocks are

$$4(\cos \varphi_j - \cos \varphi_k)^2 \prod \delta u_{...},$$

and the contribution from each diagonal block is

$$\delta u_{...}.$$

Finally,

$$\begin{aligned} \int_{O^+(2n)} f(A) dH_{aar} &= \int_{O^+(2n)} f(A) \delta A = \int_{O^+(2n)} f(A) \delta B = \\ &= \left(\frac{1}{n!} \int_{[0, \pi]^n} f(\varphi_1, \dots, \varphi_n) \prod_{j < k} 4(\cos \varphi_j - \cos \varphi_k)^2 \prod_j d\varphi_j \right) \cdot Vol(U). \end{aligned}$$

or, cheating with constant, using mathematical packages to check the correct constant for the first several n ,

$$\int_{O^+(2n)} f(A) dH_{aar} = \frac{2}{n!} \int_{[0, \pi]^n} f(\varphi_1, \dots, \varphi_n) \prod_{j < k} 4(\cos \varphi_j - \cos \varphi_k)^2 \prod_j \frac{d\varphi_j}{2\pi}. \quad (1.7)$$

1.3 $O^-(2n+2)$.

In this case there are 1, -1 among eigenvalues. Let again $n = 2$. Denote by

$$T = \begin{bmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{bmatrix}$$

the block matrix. Then Λ can be parametrized in the form

$$\Lambda = \begin{pmatrix} T_1 & 0 & 0 \\ 0 & T_2 & 0 \\ 0 & 0 & \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \end{pmatrix}, \quad 0 < \varphi_1 < \varphi_2 < \pi.$$

What changes in our fundamental relation (1.3)?

$$\delta B = U^{-1} \delta A U = (\Lambda^{-1} \delta U \Lambda - \delta U) + \delta \Lambda.$$

We have

$$\delta U = \begin{bmatrix} \begin{bmatrix} 0 & -\delta u_{21} \\ \delta u_{21} & 0 \end{bmatrix} & \begin{bmatrix} -\delta u_{31} & -\delta u_{41} \\ -\delta u_{32} & -\delta u_{42} \end{bmatrix} & \begin{bmatrix} -\delta u_{51} & -\delta u_{61} \\ -\delta u_{52} & -\delta u_{62} \end{bmatrix} \\ \begin{bmatrix} \delta u_{31} & \delta u_{32} \\ \delta u_{41} & \delta u_{42} \end{bmatrix} & \begin{bmatrix} 0 & -\delta u_{43} \\ \delta u_{43} & 0 \end{bmatrix} & \begin{bmatrix} -\delta u_{53} & -\delta u_{63} \\ -\delta u_{54} & -\delta u_{64} \end{bmatrix} \\ \begin{bmatrix} \delta u_{51} & \delta u_{52} \\ \delta u_{61} & \delta u_{62} \end{bmatrix} & \begin{bmatrix} \delta u_{53} & \delta u_{54} \\ \delta u_{63} & \delta u_{64} \end{bmatrix} & \begin{bmatrix} 0 & -\delta u_{65} \\ \delta u_{65} & 0 \end{bmatrix} \end{bmatrix},$$

where now the scalar meaning is $\delta U = \prod_{j>k} \delta u_{jk}$. New pieces comparing to the previous case are now lying on the last block row (i.e., in last 2 rows).

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} \delta u_{51} & \delta u_{52} \\ \delta u_{61} & \delta u_{62} \end{bmatrix} \begin{bmatrix} c_1 & -s_1 \\ s_1 & c_1 \end{bmatrix} - \begin{bmatrix} \delta u_{51} & \delta u_{52} \\ \delta u_{61} & \delta u_{62} \end{bmatrix},$$

and the latter can be written line-by-line in the form of two decomposed relations, for δ_{51} , δ_{52} , and distinctively for δ_{61} , δ_{62} . We have

$$\begin{pmatrix} c_1 - 1 & s_1 \\ -s_1 & c_1 - 1 \end{pmatrix} \begin{pmatrix} \delta u_{51} \\ \delta u_{52} \end{pmatrix}, \quad \begin{pmatrix} c_1 + 1 & s_1 \\ -s_1 & c_1 + 1 \end{pmatrix} \begin{pmatrix} \delta u_{61} \\ \delta u_{62} \end{pmatrix}$$

and the determinant equals

$$(2 - 2 \cos \varphi_1)(2 + 2 \cos \varphi_1) = 4 \sin^2 \varphi_1.$$

Finally,

$$\begin{aligned} \int_{O^-(2n+2)} f(A) dH_{aar} &= \int_{O^-(2n+2)} f(A) \delta A = \int_{O^-(2n+2)} f(A) \delta B = \\ &= \left(\frac{1}{n!} \int_{[0, \pi]^n} f(1, -1, \varphi_1, \dots, \varphi_n) \prod_{j < k} 4(\cos \varphi_j - \cos \varphi_k)^2 \prod_j 4 \sin^2 \varphi_j \prod_j d\varphi_j \right) \cdot Vol(U). \end{aligned}$$

and, cheating with constant,

$$\begin{aligned} \int_{O^-(2n+2)} f(A) dH_{aar} &= \int_{O^-(2n+2)} f(A) \delta A = \int_{O^-(2n+2)} f(A) \delta B = \\ &= \left(\frac{1}{n!} \int_{[0, \pi]^n} f(1, -1, \varphi_1, \dots, \varphi_n) \prod_{j < k} 4(\cos \varphi_j - \cos \varphi_k)^2 \prod_j 4 \sin^2 \varphi_j \prod_j \frac{d\varphi_j}{2\pi} \right). \end{aligned} \tag{1.8}$$

1.4 $O^+(2n+1)$.

In this case there is an eigenvalue 1. We have (again $n = 2$)

$$\Lambda = \begin{pmatrix} T_1 & 0 & 0 \\ 0 & T_2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad 0 < \varphi_1 < \varphi_2 < \pi.$$

Furthermore, δU is

$$\delta U = \begin{bmatrix} \begin{bmatrix} 0 & -\delta u_{21} \\ \delta u_{21} & 0 \end{bmatrix} & \begin{bmatrix} -\delta u_{31} & -\delta u_{41} \\ -\delta u_{32} & -\delta u_{42} \end{bmatrix} & \begin{bmatrix} -\delta u_{51} \\ -\delta u_{52} \\ -\delta u_{53} \end{bmatrix} \\ \begin{bmatrix} \delta u_{31} & \delta u_{32} \\ \delta u_{41} & \delta u_{42} \end{bmatrix} & \begin{bmatrix} 0 & -\delta u_{43} \\ \delta u_{43} & 0 \end{bmatrix} & \begin{bmatrix} -\delta u_{54} \\ -\delta u_{55} \\ -\delta u_{56} \end{bmatrix} \\ \begin{bmatrix} \delta u_{51} & \delta u_{52} \end{bmatrix} & \begin{bmatrix} \delta u_{53} & \delta u_{54} \end{bmatrix} & 0 \end{bmatrix}.$$

In $\Lambda^{-1} \delta U \Lambda - \delta U$, comparing to the case case $O^+(2n)$, the new pieces are on the last row. We have

$$(\delta u_{51}, \delta u_{52}) \begin{pmatrix} c_1 - 1 & s_1 \\ -s_1 & c_1 - 1 \end{pmatrix}, \quad \text{or} \quad \begin{pmatrix} c - 1 & s \\ -s & c - 1 \end{pmatrix} \begin{pmatrix} \delta u_{51} \\ \delta u_{52} \end{pmatrix},$$

from where the denominator is $2(1 - \cos \varphi_1)$. The result now is

$$\begin{aligned} \int_{O^+(2n+1)} f(A) d\text{Haar} &= \int_{O^+(2n+1)} f(A) \delta A = \int_{O^+(2n+1)} f(A) \delta B = \\ &= \left(\frac{1}{n!} \int_{[0, \pi]^n} f(1, \varphi_1, \dots, \varphi_n) \prod_{j < k} 4(\cos \varphi_j - \cos \varphi_k)^2 \prod_j 2(1 - \cos \varphi_j) \prod_j d\varphi_j \right) \cdot \text{Vol}(U), \end{aligned}$$

and cheating with a constant,

$$\begin{aligned} \int_{O^+(2n+1)} f(A) d\text{Haar} &= \int_{O^+(2n+1)} f(A) \delta A = \int_{O^+(2n+1)} f(A) \delta B = \\ &= \left(\frac{1}{n!} \int_{[0, \pi]^n} f(1, \varphi_1, \dots, \varphi_n) \prod_{j < k} 4(\cos \varphi_j - \cos \varphi_k)^2 \prod_j 2(1 - \cos \varphi_j) \prod_j \frac{d\varphi_j}{2\pi} \right). \end{aligned} \quad (1.9)$$

1.5 $O^-(2n+1)$.

In this case the representation Λ will have -1 instead of 1 in the last row. The rest is the same as for $O^+(2n+1)$.

$$\begin{aligned} \int_{O^-(2n+1)} f(A) d\text{Haar} &= \\ &= \left(\frac{1}{n!} \int_{[0, \pi]^n} f(-1, \varphi_1, \dots, \varphi_n) \prod_{j < k} 4(\cos \varphi_j - \cos \varphi_k)^2 \prod_j 2(1 - \cos \varphi_j) \prod_j \frac{d\varphi_j}{2\pi} \right). \end{aligned} \quad (1.10)$$

All the wrapped formulas coincide with Katz, Sarnak, Chapter 5, pages 107-108, up to a constant, with the difference that he divides all $d\varphi$ by 2π , and discard $Vol(U)$. Indeed, the latter will cancel if we want our integral be normalized. In the case $O^+(2n)$, Katz Sarnak also multiply by 2.

2 Meet Andreeiv.

Now a general statement,

$$\frac{1}{N!} \int_{X^N} \det(g_k(\theta_j))_{k,j=1}^N \cdot \det(h_k(\theta_j))_{k,j=1}^N \cdot \prod_{j=1}^N d\sigma(\theta_j) = \det \left[\int_X g_k(\theta) h_j(\theta) d\sigma(\theta) \right]_{j,k=1}^N.$$

Set

$$f()$$

2.1 $U(N)$.

In formula (1.4), set

$$f(A) = \prod_j f(e^{i\varphi_j}), \quad g_k(\varphi_j) = e^{i\varphi_j(k-1)}, \quad h_k(\varphi_j) = e^{-i\varphi_j(k-1)}, \quad d\sigma(\varphi) = f(e^{i\varphi}) \frac{d\varphi}{2\pi},$$

and since

$$g_k(\varphi) h_j(\varphi) d\sigma(\varphi) = e^{i\varphi(k-j)} f(e^{i\varphi}) \frac{d\varphi}{2\pi}, \quad \int_0^{2\pi} g_k(\varphi) h_j(\varphi) d\sigma(\varphi) = \int_0^{2\pi} e^{i\varphi(k-j)} f(e^{i\varphi}) \frac{d\varphi}{2\pi} =: m_{k-j},$$

and (1.4) equals

$$\int_{U(n)} f(A) d\text{Haar} = \left(\frac{1}{n!} \int_{[0,2\pi]^n} f(\varphi_1, \dots, \varphi_n) \prod_{j \neq k} |e^{i\varphi_j} - e^{i\varphi_k}| \prod_{j=1}^n \frac{d\varphi_j}{2\pi} \right) = \det(m_{k-j})_{j,k=1}^n.$$

2.2 $O^+(2n)$.

In (1.7) set

$$f(A) = \prod_j f(e^{i\varphi}) f(e^{-i\varphi}), \quad g_k(\varphi_j) = h_k(\varphi_j) = (2 \cos \varphi_j)^{k-1}, \quad d\sigma(\varphi) = f(e^{i\varphi}) f(e^{-i\varphi}) \frac{d\varphi}{2\pi},$$

and

$$\int_0^\pi g_k(\varphi) h_j(\varphi) d\sigma(\varphi) = \int_0^\pi (\cos \varphi)^{k-j} f(e^{i\varphi}) f(e^{-i\varphi}) \frac{d\varphi}{2\pi},$$

but this is not the form we want.

$$\int_{O^+(2n)} f(A) dA = \det \left(\int_0^\pi f(e^{i\varphi}) f(e^{-i\varphi}) (\cos \varphi)^{k-j} \frac{d\varphi}{2\pi} \right)_{k,j=1}^n.$$

2.3 $O^-(2n+2)$

In (1.8) put

$$f(A) = f(1)f(-1) \prod_j f(e^{i\varphi})f(e^{-i\varphi}), \quad g_k(\varphi) = h_k(\varphi) = (2 \cos \varphi)^{k-1}, \quad d\sigma(\varphi) = f(e^{i\varphi})f(e^{-i\varphi})4 \sin^2 \varphi \frac{d\varphi}{2\pi}.$$

We get

$$\int_{O^-(2n+2)} f(A) dH_{aar} = f(1)f(-1) \det \left(\int_0^\pi (2 \cos \varphi)^{k-j} 4 \sin^2 \varphi f(e^{i\varphi})f(e^{-i\varphi}) \frac{d\varphi}{2\pi} \right)_{j,k=1}^n.$$

2.4 $O^\pm(2n+1)$.

We set in (1.9), (1.10)

$$f(A) = f(\pm 1) \prod_j f(e^{i\varphi})f(e^{-i\varphi}), \quad g_k(\varphi) = h_k(\varphi) = (2 \cos \varphi)^{k-1}, \quad d\sigma(\varphi) = 2(1 - \cos \varphi) \frac{d\varphi}{2\pi},$$

hence

$$\int_{O^\pm(2n+1)} f(A) dH_{aar} = \det \left(\int_0^\pi (2 \cos \varphi)^{k-j} f(e^{i\varphi})f(e^{-i\varphi})2(1 - \cos \varphi) \frac{d\varphi}{2\pi} \right)_{j,k=1}^n.$$

Ha-ha, I wrote divergent integrals. Great.