

Math Camp

Justin Grimmer

Associate Professor
Department of Political Science
University of Chicago

August 28th, 2017

< Course >

The Systematic Analysis of Politics

Social Science: systematic analysis of society

The Systematic Analysis of Politics

Social Science: systematic analysis of society (Political Science: who gets what, when, and how).

The Systematic Analysis of Politics

Social Science: systematic analysis of society (Political Science: who gets what, when, and how).

Methodology: Develop and disseminate tools to make inferences about society

The Systematic Analysis of Politics

Social Science: systematic analysis of society (Political Science: who gets what, when, and how).

Methodology: Develop and disseminate tools to make inferences about society

- Mathematical models of social world

The Systematic Analysis of Politics

Social Science: systematic analysis of society (Political Science: who gets what, when, and how).

Methodology: Develop and disseminate tools to make inferences about society

- Mathematical models of social world
- Probability and Statistics used across sciences

The Systematic Analysis of Politics

Social Science: systematic analysis of society (Political Science: who gets what, when, and how).

Methodology: Develop and disseminate tools to make inferences about society

- Mathematical models of social world
- Probability and Statistics used across sciences

This class (introduction):

The Systematic Analysis of Politics

Social Science: systematic analysis of society (Political Science: who gets what, when, and how).

Methodology: Develop and disseminate tools to make inferences about society

- Mathematical models of social world
- Probability and Statistics used across sciences

This class (introduction):

- Math Camp: Develop Tools for Analysis

The Systematic Analysis of Politics

Social Science: systematic analysis of society (Political Science: who gets what, when, and how).

Methodology: Develop and disseminate tools to make inferences about society

- Mathematical models of social world
- Probability and Statistics used across sciences

This class (introduction):

- Math Camp: Develop Tools for Analysis
- Probability theory: systematic model of randomness

Course Goals

First stop in methodology sequence

Course Goals

First stop in methodology sequence

Big Goal: prepare you to make **discoveries** about social world

Course Goals

First stop in methodology sequence

Big Goal: prepare you to make **discoveries** about social world

Proximate Goals

Course Goals

First stop in methodology sequence

Big Goal: prepare you to make **discoveries** about social world

Proximate Goals

- 1) Mathematical tools to comprehend and use statistical methods
- 2) Foundation in probability theory/analytic reasoning

Course Goals

First stop in methodology sequence

Big Goal: prepare you to make **discoveries** about social world

Proximate Goals

- 1) Mathematical tools to comprehend and use statistical methods
- 2) Foundation in probability theory/analytic reasoning
- 3) Practical Computing Tools: R

Course Staff

Me: Justin Grimmer

Course Staff

Me: Justin Grimmer

- Office: Pick Hall 423

Course Staff

Me: Justin Grimmer

- Office: Pick Hall 423
- Email: grimmer@uchicago.edu

Course Staff

Me: Justin Grimmer

- Office: Pick Hall 423
- Email: grimmer@uchicago.edu
- Cell: 617-710-6803

Course Staff

Me: Justin Grimmer

- Office: Pick Hall 423
- Email: grimmer@uchicago.edu
- Cell: 617-710-6803
- Office Hours: I'm generally here all the time (9am to 5pm), just stop by [but if you need to see me with 100% probability, schedule a visit]

Course Staff

Me: Justin Grimmer

- Office: Pick Hall 423
- Email: grimmer@uchicago.edu
- Cell: 617-710-6803
- Office Hours: I'm generally here all the time (9am to 5pm), just stop by [but if you need to see me with 100% probability, schedule a visit]

TA Info

- Joshua Mausolf
Ryan Hughes
- We will hold twice weekly labs, that will occur in this room from 130-300pm (or so)
- Github for class: `github/justingrimmer/MathCamp17`

Prerequisites

No Formal Prerequisites

Prerequisites

No Formal Prerequisites
BUT

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)
 - 2) Derivatives (tangent lines, differentiation rules)

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)
 - 2) Derivatives (tangent lines, differentiation rules)
 - 3) Integrals (fundamental theorem of calculus/antidifferentiation rules)

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)
 - 2) Derivatives (tangent lines, differentiation rules)
 - 3) Integrals (fundamental theorem of calculus/antidifferentiation rules)
- We are here to help

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)
 - 2) Derivatives (tangent lines, differentiation rules)
 - 3) Integrals (fundamental theorem of calculus/antidifferentiation rules)
- We are here to help
 - No mystery to learning math: just hard work

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)
 - 2) Derivatives (tangent lines, differentiation rules)
 - 3) Integrals (fundamental theorem of calculus/antidifferentiation rules)
- We are here to help
 - No mystery to learning math: just hard work
 - Political science increasingly requires math

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)
 - 2) Derivatives (tangent lines, differentiation rules)
 - 3) Integrals (fundamental theorem of calculus/antidifferentiation rules)
- We are here to help
 - No mystery to learning math: just hard work
 - Political science increasingly requires math
 - Empirical: calculus and linear algebra

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)
 - 2) Derivatives (tangent lines, differentiation rules)
 - 3) Integrals (fundamental theorem of calculus/antidifferentiation rules)
- We are here to help
 - No mystery to learning math: just hard work
 - Political science increasingly requires math
 - Empirical: calculus and linear algebra
 - Quantitative Methodologist: Real Analysis and Grad level statistics

Prerequisites

No Formal Prerequisites

BUT

- Successful students will know differential and integral calculus
 - 1) Limits (intuitive)
 - 2) Derivatives (tangent lines, differentiation rules)
 - 3) Integrals (fundamental theorem of calculus/antidifferentiation rules)
- We are here to help
 - No mystery to learning math: just hard work
 - Political science increasingly requires math
 - Empirical: calculus and linear algebra
 - Quantitative Methodologist: Real Analysis and Grad level statistics
 - Formal Theory: Real Analysis (through measure theory), Topology

Evaluation

You're not taking this class for a grade

Evaluation

You're not taking this class for a grade \rightsquigarrow that shouldn't matter:

Evaluation

You're not taking this class for a grade \rightsquigarrow that shouldn't matter:

- Math Camp Exam

Evaluation

You're not taking this class for a grade \rightsquigarrow that shouldn't matter:

- Math Camp Exam

Grad School Irony

Evaluation

You're not taking this class for a grade \rightsquigarrow that shouldn't matter:

- Math Camp Exam

Grad School Irony Or: How I Learned to Stop Worrying and Love C's

Evaluation

You're not taking this class for a grade \rightsquigarrow that shouldn't matter:

- Math Camp Exam

Grad School Irony Or: How I Learned to Stop Worrying and Love C's

- Grades no longer matter

Evaluation

You're not taking this class for a grade \rightsquigarrow that shouldn't matter:

- Math Camp Exam

Grad School Irony Or: How I Learned to Stop Worrying and Love C's

- Grades no longer matter
- Learn as much material as possible

Evaluation

You're not taking this class for a grade \rightsquigarrow that shouldn't matter:

- Math Camp Exam

Grad School Irony Or: How I Learned to Stop Worrying and Love C's

- Grades no longer matter
- Learn as much material as possible
- If you truly only care about learning material, you'll get amazing grades

Homework

Math camp: assigned daily \rightsquigarrow Mechanics of solving problems

Lab Assignment: Twice weekly assignments, help you develop computational and mathematical skills.

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: R

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools and makes learning other programs easier

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools and makes learning other programs easier
- More start up costs than STATA, but more payoff

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools and makes learning other programs easier
- More start up costs than STATA, but more payoff

Paper writeup: \LaTeX

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools and makes learning other programs easier
- More start up costs than STATA, but more payoff

Paper writeup: \LaTeX

- Hard to write equations in Word:

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools and makes learning other programs easier
- More start up costs than STATA, but more payoff

Paper writeup: \LaTeX

- Hard to write equations in Word:
- Relatively easy in \LaTeX

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools and makes learning other programs easier
- More start up costs than STATA, but more payoff

Paper writeup: \LaTeX

- Hard to write equations in Word:
- Relatively easy in \LaTeX

$$f(x) = \frac{\exp(-\frac{(x-\mu)^2}{2\sigma^2})}{\sqrt{2\pi\sigma^2}}$$

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools and makes learning other programs easier
- More start up costs than STATA, but more payoff

Paper writeup: \LaTeX

- Hard to write equations in Word:
- Relatively easy in \LaTeX

$$f(x) = \frac{\exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)}{\sqrt{2\pi\sigma^2}}$$

- Tables/Figures/General type/Nice Presentations setting: easier in \LaTeX

Computing/Homeworks

Greatest scientific discovery of 20th Century:

Powerful personal computer (standardize science)

1956: \$10,000 megabyte

2015: <<< \$ 0.0001 per megabyte

Statistical Computing: **R**

- R: Scripting language
- Flexible, Cutting Edge Software, great visualization tools and makes learning other programs easier
- More start up costs than STATA, but more payoff

Paper writeup: \LaTeX

- Hard to write equations in Word:
- Relatively easy in \LaTeX

$$f(x) = \frac{\exp(-\frac{(x-\mu)^2}{2\sigma^2})}{\sqrt{2\pi\sigma^2}}$$

- Tables/Figures/General type/Nice Presentations setting: easier in \LaTeX
- **If you start using \LaTeX , you'll soon love it**

Course Books

- 1) Simon, Carl and Blume, Lawrence (SB). Mathematics for Economists.
- 2) Bertsekas, Dimitri P. and Tsitsiklis, John (BT) Introduction to Probability Theory (second edition)

Life in Graduate School/Academy

Three part mixture:

Life in Graduate School/Academy


Three part mixture:



George Strait

Life in Graduate School/Academy

Three part mixture:



George Strait

Kanye West

Life in Graduate School/Academy


Three part mixture:



George Strait



Kanye West



Steve Prefontaine

$\frac{1}{3}$ George Strait



$\frac{1}{3}$ George Strait



- Amarillo By Morning [Terry Stafford 1973, George Strait 1982]

$\frac{1}{3}$ George Strait



- Amarillo By Morning [Terry Stafford 1973, George Strait 1982]
- Ostensibly: song about rodeo cowboys

$\frac{1}{3}$ George Strait



- Amarillo By Morning [Terry Stafford 1973, George Strait 1982]
- Ostensibly: song about rodeo cowboys
- Really: song about being academic

$\frac{1}{3}$ George Strait



- Amarillo By Morning [Terry Stafford 1973, George Strait 1982]
- Ostensibly: song about rodeo cowboys
- Really: song about being academic
- “I ain’t got a dime/but what I got is mine/I ain’t rich/ but lord I’m free”

$\frac{1}{3}$ George Strait



- Amarillo By Morning [Terry Stafford 1973, George Strait 1982]
- Ostensibly: song about rodeo cowboys
- Really: song about being academic
- “I ain’t got a dime/but what I got is mine/I ain’t rich/ but lord I’m free”
- Academics ain’t rich (counterfactually)

$\frac{1}{3}$ George Strait




- Amarillo By Morning [Terry Stafford 1973, George Strait 1982]
- Ostensibly: song about rodeo cowboys
- Really: song about being academic
- “I ain’t got a dime/but what I got is mine/I ain’t rich/ but lord I’m free”
- Academics ain’t rich (counterfactually)
- But (lord) we’re free

$\frac{1}{3}$ George Strait




- Amarillo By Morning [Terry Stafford 1973, George Strait 1982]
- Ostensibly: song about rodeo cowboys
- Really: song about being academic
- “I ain’t got a dime/but what I got is mine/I ain’t rich/ but lord I’m free”
- Academics ain’t rich (counterfactually)
- But (lord) we’re free
- If you’re good at methods, you’ll be more rich [in expectation] and equally free

$\frac{1}{3}$ Kanye West



$\frac{1}{3}$ Kanye West

- Deal with explicit criticism (part of Hip/Hop culture)




$\frac{1}{3}$ Kanye West




- Deal with explicit criticism (part of Hip/Hop culture)
- On masterpiece album **My Beautiful Dark Twisted Fantasy**

$\frac{1}{3}$ Kanye West




- Deal with explicit criticism (part of Hip/Hop culture)
- On masterpiece album **My Beautiful Dark Twisted Fantasy**
- “Screams from the haters, got a nice ring to it/I guess every superhero needs his theme music”

$\frac{1}{3}$ Kanye West




- Deal with explicit criticism (part of Hip/Hop culture)
- On masterpiece album **My Beautiful Dark Twisted Fantasy**
- “Screams from the haters, got a nice ring to it/I guess every superhero needs his theme music”
- Kid Cudi: “These motherf**kers can’t fathom the wizardry”

$\frac{1}{3}$ Kayne West




- Deal with explicit criticism (part of Hip/Hop culture)
- On masterpiece album **My Beautiful Dark Twisted Fantasy**
- “Screams from the haters, got a nice ring to it/I guess every superhero needs his theme music”
- Kid Cudi: “These motherf**kers can’t fathom the wizadry”
- Academics: intense criticism of ideas

$\frac{1}{3}$ Kanye West



- Deal with explicit criticism (part of Hip/Hop culture)
- On masterpiece album **My Beautiful Dark Twisted Fantasy**
- “Screams from the haters, got a nice ring to it/I guess every superhero needs his theme music”
- Kid Cudi: “These motherf**kers can’t fathom the wizadry”
- Academics: intense criticism of ideas
- **Very rarely will you be told you’re doing a great job**

$\frac{1}{3}$ Kanye West



- Deal with explicit criticism (part of Hip/Hop culture)
- On masterpiece album **My Beautiful Dark Twisted Fantasy**
- “Screams from the haters, got a nice ring to it/I guess every superhero needs his theme music”
- Kid Cudi: “These motherf**kers can’t fathom the wizardry”
- Academics: intense criticism of ideas
- **Very rarely will you be told you’re doing a great job**
- Self confidence: believe in work

$\frac{1}{3}$ Steve Prefontaine

"It's not a sprint, it's a marathon".



$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons
- Marathon: 4:40 minute mile, for 26.2 miles.

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons
- Marathon: 4:40 minute mile, for 26.2 miles.
- How to train?

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons
- Marathon: 4:40 minute mile, for 26.2 miles.
- How to train?
 - Old way: get in shape (run far) rely on adrenaline in race

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons
- Marathon: 4:40 minute mile, for 26.2 miles.
- How to train?
 - Old way: get in shape (run far) rely on adrenaline in race
 - Now: races more tactical **and agonizing**

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons
- Marathon: 4:40 minute mile, for 26.2 miles.
- How to train?
 - Old way: get in shape (run far) rely on adrenaline in race
 - Now: races more tactical **and agonizing**
 - Need to prepare for agony

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons
- Marathon: 4:40 minute mile, for 26.2 miles.
- How to train?
 - Old way: get in shape (run far) rely on adrenaline in race
 - Now: races more tactical **and agonizing**
 - Need to prepare for agony
- Mantra: **sustained agony**

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons
- Marathon: 4:40 minute mile, for 26.2 miles.
- How to train?
 - Old way: get in shape (run far) rely on adrenaline in race
 - Now: races more tactical **and agonizing**
 - Need to prepare for agony
- Mantra: **sustained agony**
- Graduate School/Academics: **Sustained Agony**

$\frac{1}{3}$ Steve Prefontaine

“It’s not a sprint, it’s a marathon”.

- World class distance running: it is **hard**
- But not for the obvious reasons
- Marathon: 4:40 minute mile, for 26.2 miles.
- How to train?
 - Old way: get in shape (run far) rely on adrenaline in race
 - Now: races more tactical **and agonizing**
 - Need to prepare for agony
- Mantra: **sustained agony**
- Graduate School/Academics: **Sustained Agony**

$\frac{1}{3}$ Steve Prefontaine

Not crazy to work 40 hours on methods
alone

Not crazy to work 40 hours on methods
alone

- Methods \rightsquigarrow skills use for rest of career

Not crazy to work 40 hours on methods alone

- Methods \rightsquigarrow skills use for rest of career
- Methods \rightsquigarrow often takes deep thinking, practice

Not crazy to work 40 hours on methods
alone

- Methods \rightsquigarrow skills use for rest of career
- Methods \rightsquigarrow often takes deep thinking, practice

TAKE BREAKS!

Not crazy to work 40 hours on methods alone

- Methods \rightsquigarrow skills use for rest of career
- Methods \rightsquigarrow often takes deep thinking, practice

TAKE BREAKS!

- Regular physical activity \rightsquigarrow improve focus

Not crazy to work 40 hours on methods alone

- Methods \rightsquigarrow skills use for rest of career
- Methods \rightsquigarrow often takes deep thinking, practice

TAKE BREAKS!

- Regular physical activity \rightsquigarrow improve focus
- Time away from lab \rightsquigarrow more productive when back

$\frac{1}{3}$ Steve Prefontaine

Why work so hard?

$\frac{1}{3}$ Steve Prefontaine

Why work so hard?

- You are all smart

$\frac{1}{3}$ Steve Prefontaine

Why work so hard?

- You are all smart Really Smart

$\frac{1}{3}$ Steve Prefontaine

Why work so hard?

- You are all smart Really Smart Mother-in-law brags about you smart

Why work so hard?

- **You are all smart** Really Smart Mother-in-law brags about you smart
- Everyone entering graduate school at top programs this fall

Why work so hard?

- **You are all smart** Really Smart Mother-in-law brags about you smart
- Everyone entering graduate school at top programs this fall
- Success: **work**

Why work so hard?

- **You are all smart** Really Smart Mother-in-law brags about you smart
- Everyone entering graduate school at top programs this fall
- Success: **work**
- Treat grad school like a job

Why work so hard?

- **You are all smart** Really Smart Mother-in-law brags about you smart
- Everyone entering graduate school at top programs this fall
- Success: **work**
- Treat grad school like a job
- Who gets ahead? who gets the most work done on the smartest ideas

Preliminaries

What can you learn in a math camp?

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar
- 4) This will not substitute for a richer math background and we won't expect it to

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar
- 4) This will not substitute for a richer math background and we won't expect it to

Do not let yourself get lost.

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar
- 4) This will not substitute for a richer math background and we won't expect it to

Do not let yourself get lost.

If at.

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar
- 4) This will not substitute for a richer math background and we won't expect it to

Do not let yourself get lost.

If at. any.

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar
- 4) This will not substitute for a richer math background and we won't expect it to

Do not let yourself get lost.

If at. any. point.

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar
- 4) This will not substitute for a richer math background and we won't expect it to

Do not let yourself get lost.

If at. any. point. you have a question

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar
- 4) This will not substitute for a richer math background and we won't expect it to

Do not let yourself get lost.

If at. any. point. you have a question please ask !

Preliminaries

What can you learn in a math camp?

- 1) Introduction to more sophisticated mathematics (notation)
- 2) Getting acquainted with proof techniques and proofs
- 3) I'm going to introduce ideas/example problems common in research that will help with your seminar
- 4) This will not substitute for a richer math background and we won't expect it to

Do not let yourself get lost.

If at. any. point. you have a question please ask !

Smartest people ask the most questions!

Let's get to work

Sets

A **set** is a collection of objects.

$$A = \{1, 2, 3\}$$

$$B = \{4, 5, 6\}$$

$$C = \{\text{First year cohort}\}$$

$$D = \{\text{Stanford Faculty}\}$$

Definition

If A is a set, we say that x is an element of A by writing, $x \in A$. If x is not an element of A then, we write $x \notin A$.

Definition

If A is a set, we say that x is an element of A by writing, $x \in A$. If x is not an element of A then, we write $x \notin A$.

$$- 1 \in \{1, 2, 3\}$$

Definition

If A is a set, we say that x is an element of A by writing, $x \in A$. If x is not an element of A then, we write $x \notin A$.

- $1 \in \{1, 2, 3\}$
- $4 \in \{4, 5, 6\}$

Definition

If A is a set, we say that x is an element of A by writing, $x \in A$. If x is not an element of A then, we write $x \notin A$.

- $1 \in \{1, 2, 3\}$
- $4 \in \{4, 5, 6\}$
- Will $\notin \{\text{First year cohort}\}$

Definition

If A is a set, we say that x is an element of A by writing, $x \in A$. If x is not an element of A then, we write $x \notin A$.

- $1 \in \{1, 2, 3\}$
- $4 \in \{4, 5, 6\}$
- Will $\notin \{\text{First year cohort}\}$
- Justin $\in \{\text{Stanford Faculty}\}$

Definition

If A is a set, we say that x is an element of A by writing, $x \in A$. If x is not an element of A then, we write $x \notin A$.

- $1 \in \{1, 2, 3\}$
- $4 \in \{4, 5, 6\}$
- Will $\notin \{\text{First year cohort}\}$
- Justin $\in \{\text{Stanford Faculty}\}$

Why Care?

Definition

If A is a set, we say that x is an element of A by writing, $x \in A$. If x is not an element of A then, we write $x \notin A$.

- $1 \in \{1, 2, 3\}$
- $4 \in \{4, 5, 6\}$
- Will $\notin \{\text{First year cohort}\}$
- Justin $\in \{\text{Stanford Faculty}\}$

Why Care?

- Sets are necessary for probability theory

Definition

If A is a set, we say that x is an element of A by writing, $x \in A$. If x is not an element of A then, we write $x \notin A$.

- $1 \in \{1, 2, 3\}$
- $4 \in \{4, 5, 6\}$
- Will $\notin \{\text{First year cohort}\}$
- Justin $\in \{\text{Stanford Faculty}\}$

Why Care?

- Sets are necessary for probability theory
- Defining **set** is equivalent to choosing population of interest (usually)

Definition

If A and B are sets, then we say that $A = B$ if, for all $x \in A$ then $x \in B$ and for all $y \in B$ then $y \in A$.

Definition

If A and B are sets, then we say that $A = B$ if, for all $x \in A$ then $x \in B$ and for all $y \in B$ then $y \in A$.

- Test to determine equality:

Definition

If A and B are sets, then we say that $A = B$ if, for all $x \in A$ then $x \in B$ and for all $y \in B$ then $y \in A$.

- Test to determine equality:
 - Take all elements of A , see if in B

Definition

If A and B are sets, then we say that $A = B$ if, for all $x \in A$ then $x \in B$ and for all $y \in B$ then $y \in A$.

- Test to determine equality:
 - Take all elements of A , see if in B
 - Take all elements of B , see if in A

Definition

If A and B are sets, then we say that $A = B$ if, for all $x \in A$ then $x \in B$ and for all $y \in B$ then $y \in A$.

- Test to determine equality:
 - Take all elements of A , see if in B
 - Take all elements of B , see if in A

Definition

If A and B are sets, then we say that $A \subset B$ is, for all $x \in A$, then $x \in B$.

Definition

If A and B are sets, then we say that $A = B$ if, for all $x \in A$ then $x \in B$ and for all $y \in B$ then $y \in A$.

- Test to determine equality:
 - Take all elements of A , see if in B
 - Take all elements of B , see if in A

Definition

If A and B are sets, then we say that $A \subset B$ is, for all $x \in A$, then $x \in B$.

Difference between definitions?

Theorem

Let A and B be sets. If $A = B$ then $A \subset B$ and $B \subset A$

Theorem

Let A and B be sets. If $A = B$ then $A \subset B$ and $B \subset A$

Proof.

Suppose $A = B$. By definition, if $x \in A$ then $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$. □

Theorem

Let A and B be sets. If $A = B$ then $A \subset B$ and $B \subset A$

Proof.

Suppose $A = B$. By definition, if $x \in A$ then $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$. \square

Theorem

Let A and B be sets. If $A \subset B$ and $B \subset A$ then $A = B$

Theorem

Let A and B be sets. If $A = B$ then $A \subset B$ and $B \subset A$

Proof.

Suppose $A = B$. By definition, if $x \in A$ then $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$. \square

Theorem

Let A and B be sets. If $A \subset B$ and $B \subset A$ then $A = B$

Proof.

Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. \square

Theorem

Let A and B be sets. Then $A = B$ *if and only if* $A \subset B$ and $B \subset A$.



Theorem

Let A and B be sets. Then $A = B$ *if and only if* $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.



Theorem

Let A and B be sets. Then $A = B$ *if and only if* $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

Theorem

Let A and B be sets. Then $A = B$ *if and only if* $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

When a proof says if and only if it is showing two things.

Theorem

Let A and B be sets. Then $A = B$ *if and only if* $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

When a proof says if and only if it is showing two things.

- **If** or that a condition is **sufficient**

Theorem

Let A and B be sets. Then $A = B$ **if and only if** $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

When a proof says if and only if it is showing two things.

- **If** or that a condition is **sufficient**
- **Only If** or that a condition is necessary

Theorem

Let A and B be sets. Then $A = B$ **if and only if** $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

When a proof says if and only if it is showing two things.

- **If** or that a condition is **sufficient**
- **Only If** or that a condition is necessary

Example of sufficient, but not necessary

Theorem

Let A and B be sets. Then $A = B$ **if and only if** $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

When a proof says if and only if it is showing two things.

- **If** or that a condition is **sufficient**
- **Only If** or that a condition is necessary

Example of sufficient, but not necessary

- If candidate wins the electoral college, then president (can be president through vote of House too)

Theorem

Let A and B be sets. Then $A = B$ *if and only if* $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

When a proof says if and only if it is showing two things.

- **If** or that a condition is **sufficient**
- **Only If** or that a condition is necessary

Example of sufficient, but not necessary

- If candidate wins the electoral college, then president (can be president through vote of House too)

Example of necessary, but not sufficient

Theorem

Let A and B be sets. Then $A = B$ **if and only if** $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

When a proof says if and only if it is showing two things.

- **If** or that a condition is **sufficient**
- **Only If** or that a condition is necessary

Example of sufficient, but not necessary

- If candidate wins the electoral college, then president (can be president through vote of House too)

Example of necessary, but not sufficient

- Only if a candidate is older than 35 can s/he be president (but clearly not sufficient)

Theorem

Let A and B be sets. Then $A = B$ **if and only if** $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. For all $x \in A$, then $x \in B$. And for all $y \in B$, $y \in A$. Or, every element in A is in B and each element of B is in A . $A = B$. □

When a proof says if and only if it is showing two things.

- **If** or that a condition is **sufficient**
- **Only If** or that a condition is necessary

Example of sufficient, but not necessary

- If candidate wins the electoral college, then president (can be president through vote of House too)

Example of necessary, but not sufficient

- Only if a candidate is older than 35 can s/he be president (but clearly not sufficient)

Contradiction



Contradiction

- Many ways to prove the same theorem.



Contradiction

- Many ways to prove the same theorem.
- **Contradiction**: assume theorem is false, show that this leads to logical contradiction



Contradiction

- Many ways to prove the same theorem.
- **Contradiction**: assume theorem is false, show that this leads to logical contradiction
- **Indirect proof**: setting up proof hardest part



Contradiction

- Many ways to prove the same theorem.
- **Contradiction**: assume theorem is false, show that this leads to logical contradiction
- **Indirect proof**: setting up proof hardest part

Theorem

Let A and B be sets. Then $A = B$ if and only if $A \subset B$ and $B \subset A$.



Contradiction

- Many ways to prove the same theorem.
- **Contradiction**: assume theorem is false, show that this leads to logical contradiction
- **Indirect proof**: setting up proof hardest part

Theorem

*Let A and B be sets. Then $A = B$ **if and only if** $A \subset B$ and $B \subset A$.*

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.



Contradiction

- Many ways to prove the same theorem.
- **Contradiction**: assume theorem is false, show that this leads to logical contradiction
- **Indirect proof**: setting up proof hardest part

Theorem

Let A and B be sets. Then $A = B$ **if and only if** $A \subset B$ and $B \subset A$.

Proof.

\Rightarrow Suppose $A = B$. By definition, if $x \in A$, $x \in B$. So $A \subset B$. Again, by definition, if $y \in B$ then $y \in A$. So $B \subset A$.

\Leftarrow Suppose $A \subset B$ and that $B \subset A$. Now, by way of contradiction, suppose that $A \neq B$. $A \neq B$ only if there is $x \in A$ and $x \notin B$ or if $y \in B$ and $y \notin A$. But then, either $A \not\subset B$ or $B \not\subset A$, contradicting our initial assumption. □

Set Builder Notation

- Some famous sets
 - $J = \{1, 2, 3, \dots\}$
 - $Z = \{\dots, -2, -1, 0, 1, 2, \dots\}$
 - $\mathbb{R} = \text{real numbers}$ (more to come about this)
- Use **set builder notation** to identify subsets
 - $[a, b] = \{x : x \in \mathbb{R} \text{ and } a \leq x \leq b\}$
 - $(a, b] = \{x : x \in \mathbb{R} \text{ and } a < x \leq b\}$
 - $[a, b) = \{x : x \in \mathbb{R} \text{ and } a \leq x < b\}$
 - $(a, b) = \{x : x \in \mathbb{R} \text{ and } a < x < b\}$
 - \emptyset

Set Operations

We can build new sets with **set operations**.

Set Operations

We can build new sets with **set operations**.

Definition

*Suppose A and B are sets. Define the **Union** of sets A and B as the new set that contains all elements in set A **or** in set B . In notation,*

$$\begin{aligned} C &= A \cup B \\ &= \{x : x \in A \text{ or } x \in B\} \end{aligned}$$

Set Operations

We can build new sets with **set operations**.

Definition

*Suppose A and B are sets. Define the **Union** of sets A and B as the new set that contains all elements in set A **or** in set B . In notation,*

$$\begin{aligned} C &= A \cup B \\ &= \{x : x \in A \text{ or } x \in B\} \end{aligned}$$

- $A = \{1, 2, 3\}, B = \{3, 4, 5\}$, then $C = A \cup B = \{1, 2, 3, 4, 5\}$

Set Operations

We can build new sets with **set operations**.

Definition

*Suppose A and B are sets. Define the **Union** of sets A and B as the new set that contains all elements in set A **or** in set B . In notation,*

$$\begin{aligned} C &= A \cup B \\ &= \{x : x \in A \text{ or } x \in B\} \end{aligned}$$

- $A = \{1, 2, 3\}$, $B = \{3, 4, 5\}$, then $C = A \cup B = \{1, 2, 3, 4, 5\}$
- $D = \{\text{First Year Cohort}\}$, $E = \{\text{Me}\}$, then
 $F = D \cup E = \{\text{First Year Cohort}, \text{ME}\}$

Set Operations

Definition

Suppose A and B are sets. Define the *Intersection* of sets A and B as the new set that contains all elements in set A *and* set B . In notation,

$$\begin{aligned} C &= A \cap B \\ &= \{x : x \in A \text{ and } x \in B\} \end{aligned}$$

Set Operations

Definition

Suppose A and B are sets. Define the *Intersection* of sets A and B as the new set that contains all elements in set A *and* set B . In notation,

$$\begin{aligned} C &= A \cap B \\ &= \{x : x \in A \text{ and } x \in B\} \end{aligned}$$

- $A = \{1, 2, 3\}, B = \{3, 4, 5\}$, then, $C = A \cap B = \{3\}$

Set Operations

Definition

Suppose A and B are sets. Define the *Intersection* of sets A and B as the new that contains all elements in set A *and* set B . In notation,

$$\begin{aligned} C &= A \cap B \\ &= \{x : x \in A \text{ and } x \in B\} \end{aligned}$$

- $A = \{1, 2, 3\}, B = \{3, 4, 5\}$, then, $C = A \cap B = \{3\}$
- $D = \{\text{First Year Cohort}\}, E = \{\text{Me}\}$, then $F = D \cap E = \emptyset$

Some Facts about Sets (No Venn Diagrams!!!)

$$1) A \cap B = B \cap A$$

Some Facts about Sets (No Venn Diagrams!!!)

1) $A \cap B = B \cap A$

Proof.

This fact (theorem) says that the **set** $A \cap B$ is equal to the set $B \cap A$. We can use the definition of equal sets to test this. Suppose $x \in A \cap B$. Then $x \in A$ and $x \in B$. By definition, then, $x \in B \cap A$. Now, suppose $y \in B \cap A$. Then $y \in B$ and $y \in A$. So, by definition of intersection $y \in A \cap B$. This implies $A \cap B = B \cap A$ □

Some Facts about Sets (No Venn Diagrams!!!)

$$1) A \cap B = B \cap A$$

$$5) A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

Some Facts about Sets (No Venn Diagrams!!!)

$$1) A \cap B = B \cap A$$

$$5) A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

Proof.

Suppose $x \in A \cap (B \cup C)$. Then $x \in B$ or $x \in C$ and $x \in A$. This implies that $x \in (A \cap B)$ or $x \in (A \cap C)$. Or, $x \in (A \cap B) \cup (A \cap C)$. Now, suppose $y \in (A \cap B) \cup (A \cap C)$. Then, $y \in A$ and $y \in B$ or $y \in C$. Well, this implies $y \in A \cap (B \cup C)$. And we have established equality \square

Some Facts about Sets (No Venn Diagrams!!!)

1) $A \cap B = B \cap A$

2) $A \cup B = B \cup A$

3) $(A \cap B) \cap C = A \cap (B \cap C)$

4) $(A \cup B) \cup C = A \cup (B \cup C)$

5) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

6) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

Break into groups, derive for the remaining facts

Ordered Pair

You've seen an **ordered pair** before,

$$(a, b)$$

Definition

*Suppose we have two sets, A and B . Define the **Cartesian product** of A and B , $A \times B$ as the set of all ordered pairs (a, b) , where $a \in A$ and $b \in B$. In other words,*

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$$

Example:

$A = \{1, 2\}$ and $B = \{3, 4\}$, then,

$$A \times B = \{(1, 3); (1, 4); (2, 3); (2, 4)\}$$

Function

Start with general and move to specific— (abstract just takes time to get acquainted)

Function

Start with general and move to specific— (abstract just takes time to get acquainted)

Definition

A *relation* is a set of ordered pairs. A *function* F is a relation such that,

$$(x, y) \in F \quad ; \quad (x, z) \in F \Rightarrow y = z$$

We will commonly write a function as $F(x)$, where $x \in \text{Domain } F$ and $F(x) \in \text{Codomain } F$. It is common to see people write,

$$F : A \rightarrow B$$

where A is domain and B is codomain

Function

Start with general and move to specific— (abstract just takes time to get acquainted)

Definition

A *relation* is a set of ordered pairs. A *function* F is a relation such that,

$$(x, y) \in F \quad ; \quad (x, z) \in F \Rightarrow y = z$$

We will commonly write a function as $F(x)$, where $x \in \text{Domain } F$ and $F(x) \in \text{Codomain } F$. It is common to see people write,

$$F : A \rightarrow B$$

where A is domain and B is codomain

Examples

Function

Start with general and move to specific— (abstract just takes time to get acquainted)

Definition

A *relation* is a set of ordered pairs. A *function* F is a relation such that,

$$(x, y) \in F \quad ; \quad (x, z) \in F \Rightarrow y = z$$

We will commonly write a function as $F(x)$, where $x \in \text{Domain } F$ and $F(x) \in \text{Codomain } F$. It is common to see people write,

$$F : A \rightarrow B$$

where A is domain and B is codomain

Examples

- $F(x) = x$

Function

Start with general and move to specific— (abstract just takes time to get acquainted)

Definition

A *relation* is a set of ordered pairs. A *function* F is a relation such that,

$$(x, y) \in F \quad ; \quad (x, z) \in F \Rightarrow y = z$$

We will commonly write a function as $F(x)$, where $x \in \text{Domain } F$ and $F(x) \in \text{Codomain } F$. It is common to see people write,

$$F : A \rightarrow B$$

where A is domain and B is codomain

Examples

- $F(x) = x$
- $F(x) = x^2$

Function

Start with general and move to specific— (abstract just takes time to get acquainted)

Definition

A *relation* is a set of ordered pairs. A *function* F is a relation such that,

$$(x, y) \in F \quad ; \quad (x, z) \in F \Rightarrow y = z$$

We will commonly write a function as $F(x)$, where $x \in \text{Domain } F$ and $F(x) \in \text{Codomain } F$. It is common to see people write,

$$F : A \rightarrow B$$

where A is domain and B is codomain

Examples

- $F(x) = x$
- $F(x) = x^2$
- $F(x) = \sqrt{x}$

R Computing Language

- We're going to use R throughout the course
- R as calculator :


```
> 1 + 1  
[1] 2  
> 'Hello World'  
[1] "Hello World"
```

- `object<- 2` ## assign numbers to objects
- R has functions defined, we can define them to objects as well

```
first.func<- function(x) {  
  out<- 2*x  
  return(out) }
```


```
first.func(2)  
[1] 4
```

Plotting Functions



```
x<- seq(-2, 2,  
len=1000)  
plot(x~x) ##  
Results may vary
```


Plotting Functions



```
x<- seq(-2, 2,  
len=1000)  
x.2<- x*x  
plot(x.2~x)
```


Plotting Functions

$$f(x) = \sin(2 \cdot x)$$



```
x<- seq(-2, 2,  
len=1000)  
sin.2x<- sin(2*x)  
plot(sin.2x~x)
```

Plotting Functions



Exponents, Logarithms, and All That

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

$$a^x \times a^y = a^{x+y}$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

$$a^x \times a^y = a^{x+y}$$

$$(a^x)^y = a^{x \times y}$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

$$a^x \times a^y = a^{x+y}$$

$$(a^x)^y = a^{x \times y}$$

$$\frac{a^x}{a^y} = a^{x-y}$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

$$a^x \times a^y = a^{x+y}$$

$$(a^x)^y = a^{x \times y}$$

$$\frac{a^x}{a^y} = a^{x-y}$$

$$\frac{1}{a^x} = a^{-x}$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

$$a^x \times a^y = a^{x+y}$$

$$(a^x)^y = a^{x \times y}$$

$$\frac{a^x}{a^y} = a^{x-y}$$

$$\frac{1}{a^x} = a^{-x}$$

$$a^x \times b^x = (a \times b)^x$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

$$a^x \times a^y = a^{x+y}$$

$$(a^x)^y = a^{x \times y}$$

$$\frac{a^x}{a^y} = a^{x-y}$$

$$\frac{1}{a^x} = a^{-x}$$

$$a^x \times b^x = (a \times b)^x$$

$$a^0 = 1$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

$$a^x \times a^y = a^{x+y}$$

$$(a^x)^y = a^{x \times y}$$

$$\frac{a^x}{a^y} = a^{x-y}$$

$$\frac{1}{a^x} = a^{-x}$$

$$a^x \times b^x = (a \times b)^x$$

$$a^0 = 1$$

$$a^1 = a$$

Exponents, Logarithms, and All That

$$f(x) = 2^x$$

$$g(x) = e^x$$

Some rules of exponents remember a could equal e

$$a^x \times a^y = a^{x+y}$$

$$(a^x)^y = a^{x \times y}$$

$$\frac{a^x}{a^y} = a^{x-y}$$

$$\frac{1}{a^x} = a^{-x}$$

$$a^x \times b^x = (a \times b)^x$$

$$a^0 = 1$$

$$a^1 = a$$

$$1^x = 1$$

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$
- $\log e = 1$ (because $e^1 = e$)

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)
- $\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)
- $\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$
- $\log(a^b) = b \log(a)$

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)
- $\log(\frac{a}{b}) = \log(a) - \log(b)$
- $\log(a^b) = b \log(a)$
- $\log(1) = 0$

Exponents, Logarithms, and All That

Logarithm \log is a **class** of functions.

- $\log_e z =$ what number x solves $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log_e = \log$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)
- $\log(\frac{a}{b}) = \log(a) - \log(b)$
- $\log(a^b) = b \log(a)$
- $\log(1) = 0$
- $\log(e) = 1$

Properties of Functions

Two important properties of functions

Properties of Functions

Two important properties of functions

Definition

A function $f : A \rightarrow B$ is 1-1 (one-to-one, or injective) if for all $y \in A$ and $z \in A$ in Domain, $f(y) = f(z)$ implies $y = z$. In other words, preserves distinctiveness.

Properties of Functions

Two important properties of functions

Definition

A function $f : A \rightarrow B$ is 1-1 (one-to-one, or injective) if for all $y \in A$ and $z \in A$ in Domain, $f(y) = f(z)$ implies $y = z$. In other words, preserves distinctiveness.

- $f(x) = x$

Properties of Functions

Two important properties of functions

Definition

A function $f : A \rightarrow B$ is 1-1 (one-to-one, or injective) if for all $y \in A$ and $z \in A$ in Domain, $f(y) = f(z)$ implies $y = z$. In other words, preserves distinctiveness.

- $f(x) = x$
- $f(x) = x^2$

Properties of Functions

Two important properties of functions

Definition

A function $f : A \rightarrow B$ is 1-1 (one-to-one, or injective) if for all $y \in A$ and $z \in A$ in Domain, $f(y) = f(z)$ implies $y = z$. In other words, preserves distinctiveness.

- $f(x) = x$
- $f(x) = x^2$

Definition

A function $f : A \rightarrow B$ is onto (surjective) if for all $b \in B$ there exists (\exists) $a \in A$ such that $f(a) = b$.

Properties of Functions

Two important properties of functions

Definition

A function $f : A \rightarrow B$ is 1-1 (one-to-one, or injective) if for all $y \in A$ and $z \in A$ in Domain, $f(y) = f(z)$ implies $y = z$. In other words, preserves distinctiveness.

- $f(x) = x$
- $f(x) = x^2$

Definition

A function $f : A \rightarrow B$ is onto (surjective) if for all $b \in B$ there exists $(\exists) a \in A$ such that $f(a) = b$.

- $f : \{\dots, -2, -1, 0, 1, 2, \dots\} \rightarrow \{0, 1, 2, \dots\}$ and $f(x) = |x|$. onto, but not 1-1.

Properties of Functions

Two important properties of functions

Definition

A function $f : A \rightarrow B$ is 1-1 (one-to-one, or injective) if for all $y \in A$ and $z \in A$ in Domain, $f(y) = f(z)$ implies $y = z$. In other words, preserves distinctiveness.

- $f(x) = x$
- $f(x) = x^2$

Definition

A function $f : A \rightarrow B$ is onto (surjective) if for all $b \in B$ there exists $(\exists) a \in A$ such that $f(a) = b$.

- $f : \{\dots, -2, -1, 0, 1, 2, \dots\} \rightarrow \{0, 1, 2, \dots\}$ and $f(x) = |x|$. onto, but not 1-1.
- $f : \mathbb{R} \rightarrow \mathbb{R} f(x) = x$. Onto and 1-1, **bijjective**

Composite Functions

Definition

Suppose $f : A \rightarrow B$ and $g : B \rightarrow C$. Then, define,

$$g \circ f = g(f(x))$$

- $f(x) = x$, $g(x) = x^2$. Then $g \circ f = x^2$.
- $f(x) = \sqrt{x}$, $g(x) = e^x$. Then $g \circ f = e^{\sqrt{x}}$.
- $f(x) = \sin(x)$, $g(x) = |x|$. Then $g \circ f = |\sin(x)|$.

Inverse Function

Definition

*Suppose a function f is 1-1. Then we'll define f^{-1} as its **inverse** if,*

$$f^{-1}(f(x)) = x$$

Why do we need 1-1?

Induction

Well Ordering Principle Every non-empty set J has a smallest number

Induction

Well Ordering Principle Every non-empty set J has a smallest number

Theorem

If $P(n)$ is a statement containing the variable n such that

- i. $P(1)$ is a true statement, and*
 - ii. for each $k \in 1, 2, 3, 4, \dots, n, \dots$ if $P(k)$ is true then $P(k + 1)$ is true*
- then $P(n)$ is true for all $n \in 1, 2, 3, 4, \dots, n, \dots$*

Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that

- i. $P(1)$ is true



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that

- i. $P(1)$ is true
- ii. If $P(k)$ is true then $P(k + 1)$ is true.



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that

- i. $P(1)$ is true
- ii. If $P(k)$ is true then $P(k + 1)$ is true.

Now suppose, **by way of contradiction** that there exists N such that $P(N)$ is false. This implies that



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that

- i. $P(1)$ is true
- ii. If $P(k)$ is true then $P(k + 1)$ is true.

Now suppose, **by way of contradiction** that there exists N such that $P(N)$ is false. This implies that

$$S = \{x : P(x) \text{ is not true} \}$$



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that

- i. $P(1)$ is true
- ii. If $P(k)$ is true then $P(k + 1)$ is true.

Now suppose, **by way of contradiction** that there exists N such that $P(N)$ is false. This implies that

$$S = \{x : P(x) \text{ is not true} \}$$

By well ordering principle, there is smallest member of S , call it n_0 .



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that

- i. $P(1)$ is true
- ii. If $P(k)$ is true then $P(k + 1)$ is true.

Now suppose, **by way of contradiction** that there exists N such that $P(N)$ is false. This implies that

$$S = \{x : P(x) \text{ is not true} \}$$

By well ordering principle, there is smallest member of S , call it n_0 . By *i*. we know that $n_0 > 1$. Further, because n_0 is smallest member of S , then $P(n_0)$ is false, but $P(n_0 - 1)$ is true.



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that

- i. $P(1)$ is true
- ii. If $P(k)$ is true then $P(k + 1)$ is true.

Now suppose, **by way of contradiction** that there exists N such that $P(N)$ is false. This implies that

$$S = \{x : P(x) \text{ is not true} \}$$

By well ordering principle, there is smallest member of S , call it n_0 . By *i*. we know that $n_0 > 1$. Further, because n_0 is smallest member of S , then $P(n_0)$ is false, but $P(n_0 - 1)$ is true. But now we have a problem, because if $P(n_0 - 1)$ is true, then $P(n_0)$ is also true.



Induction and Contradiction

We'll use **contradiction** and well ordering to prove that induction works.

Proof.

Suppose $P(n)$ is some statement about the variable n and that

- i. $P(1)$ is true
- ii. If $P(k)$ is true then $P(k + 1)$ is true.

Now suppose, **by way of contradiction** that there exists N such that $P(N)$ is false. This implies that

$$S = \{x : P(x) \text{ is not true} \}$$

By well ordering principle, there is smallest member of S , call it n_0 . By *i*. we know that $n_0 > 1$. Further, because n_0 is smallest member of S , then $P(n_0)$ is false, but $P(n_0 - 1)$ is true. But now we have a problem, because if $P(n_0 - 1)$ is true, then $P(n_0)$ is also true. This implies that there is no smallest element of S . **CONTRADICTION** □

Summing N numbers

Induction is a useful proof technique.

Theorem

$$\sum_{i=1}^N i = 1 + 2 + 3 + 4 + \dots + N = \frac{N(N+1)}{2}$$

Two conditions to show:

i. $\sum_{i=1}^1 i = 1$ and $\frac{1(1+1)}{2} = 1$

Summing N numbers

ii. Suppose true N . Then, for $N + 1$ we have,

$$\begin{aligned}\sum_{i=1}^{N+1} i &= \sum_{i=1}^N i + (N + 1) \\ &= \frac{N(N + 1)}{2} + \frac{2(N + 1)}{2} \\ &= \frac{(N + 1)(N + 2)}{2} \\ &= \frac{(N + 1)((N + 1) + 1)}{2}\end{aligned}$$

Conditions of induction met. Therefore, proof complete

Very Simple R Code

Finite, Countable, and Uncountable

Three sizes of sets

- 1) A set, X is finite if there is a bijective function from $\{1, 2, 3, \dots, n\}$ to X .
 - 2) A set X is **countably infinite** if there is a bijective function from $\{1, 2, 3, 4, \dots, \}$ to X .
 - 3) A set X is **uncountably infinite** if it is not countable
- The **Real numbers** are **uncountably infinite**

Recap

We've covered a lot.

Recap

We've covered a lot.

PLEASE don't worry—we're here to help!

Recap

We've covered a lot.

PLEASE don't worry—we're here to help!

1) Sets + Operations

Recap

We've covered a lot.

PLEASE don't worry—we're here to help!

- 1) Sets + Operations
- 2) Functions

Recap

We've covered a lot.

PLEASE don't worry—we're here to help!

- 1) Sets + Operations
- 2) Functions
- 3) Contradiction, Induction, and direct proofs

Tomorrow:

- Convergence of sequences
- Limits
- Continuity
- Derivatives