

Seminar 1

Operation: $*$: $A \times A \rightarrow A$ with $x, y \in A \Rightarrow x * y \in A$.

Grupoid: $(A, *)$

Semigroup: $(A, *)$ grupoid + associativity

Monoid: $(A, *)$ semigroup + identity element

Group: $(A, *)$ monoid + all elements have a symmetric

Abelian group: $(A, *)$ group + commutativity

Subgroupoid = stable part: $\forall a, b \in A \Rightarrow a * b \in A$

Subgroup: $H \leq (G, *)$ if H is a stable part in G ($H \subseteq G$) and $(H, *)$ is a group.

1. Addition: $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$

Subtraction: $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$

Multiplication: $\mathbb{N}, \mathbb{Z}, \mathbb{Q}^*, \mathbb{R}^*, \mathbb{C}^*$

Division: $\mathbb{Q}^*, \mathbb{R}^*, \mathbb{C}^*$

2. i 3 elements in 3 spaces $\Rightarrow 3^9$

	a	b	c
a			
b			
c			

ii 3^3 (3 elements in 3 free spaces) and 3^3 (3 commutative elements in 3 spaces) $\Rightarrow 3^6$.

	a	b	c
a		c	b
b	c		a
c	b	a	

iii 3^4 (3 elements in 4 free spaces) and 3 elements, which can be e $\Rightarrow 3^5$

	e	b	c
e	e	b	c
b	b		
c	c		

Generalization:

i n^{n^2}

ii $n^n \cdot n^{\frac{n(n-1)}{2}}$

iii $n^{(n-1)^2+1}$

3. $(\mathbb{Z}, +), (\mathbb{Q}, +), (\mathbb{R}, +), (\mathbb{C}, +)$ and $(\mathbb{Q}^*, \cdot), (\mathbb{R}^*, \cdot), (\mathbb{C}^*, \cdot)$.

4. i Stable part: $\forall x, y \in \mathbb{R} \Rightarrow x * y = x + y + xy = (x+1)(y+1) - 1 \in \mathbb{R}$

Associativity: $\forall x, y \in \mathbb{R} \Rightarrow (x * y) * z = x * (y * z)$

Identity element: $\exists e \in \mathbb{R}$ such that $\forall x \in \mathbb{R} \Rightarrow x * e = e * x = x$

Commutativity: $\forall x, y \in \mathbb{R} \Rightarrow x * y = y * x$.

ii Let A be our interval. Then A is a stable subset of $(\mathbb{R}, *) \iff \forall x, y \in A \Rightarrow x * y \in A$.

$x, y \in A \Rightarrow -1 \leq x, -1 \leq y \Rightarrow 0 \leq x+1, 0 \leq y+1 \Rightarrow 0 \leq (x+1)(y+1) \Rightarrow -1 \leq (x+1)(y+1) - 1 \Rightarrow x * y \in A$

5. i Here is interesting to see the associativity: $\forall x, y, z \in \mathbb{N} \Rightarrow (x * y) * z = \gcd(x, y) * z = \gcd(\gcd(x, y), z) = \alpha \Rightarrow \alpha \mid \gcd(x, y)$ and $\alpha \mid z$. From $\gcd(x, y) = d \Rightarrow x = dx_1$ and $y = dy_1$, but $\alpha \mid d \Rightarrow \alpha \mid x$ and $\alpha \mid y \Rightarrow \alpha \mid x, y, z \Rightarrow \alpha \mid \gcd(y, z) \Rightarrow \alpha \mid \gcd(x, \gcd(y, z))$. Analogous for $\gcd(x, \gcd(y, z)) \mid \alpha$.

ii $\forall x, y \in D_n \Rightarrow x \mid n$ and $y \mid n \Rightarrow n = xd_1$ and $n = yd_2$. We compute $x * y = \gcd(x, y) = \alpha \Rightarrow x = \alpha x_1$ and $y = \alpha y_1 \Rightarrow n = \alpha x_1 d_1$ and $n = \alpha y_1 d_2 \Rightarrow \alpha \mid n \Rightarrow \gcd(x, y) \mid n \Rightarrow x * y \in D_n$. Associativity, commutativity and identity element are easy to prove.

iii $D_6 = \{1, 2, 3, 6\}$

	1	2	3	6
1	1	1	1	1
2	1	2	1	2
3	1	1	3	3
6	1	2	3	6

6. $H \subseteq \mathbb{Z}$ and H stable part of $\mathbb{Z} \Rightarrow \exists x \in H \Rightarrow \forall n \in \mathbb{N}^*$ we have $x^n \in H$,
but H is finite $\Rightarrow \exists n \in \mathbb{N}^*$ such that $x^i = x^j, i, j \in \mathbb{N}^*$ and $0 < i < j \Rightarrow$
 $x \in \{-1, 0, 1\} \Rightarrow H$ can be $\emptyset, \{0\}, \{1\}, \{0, 1\}, \{-1, 1\}, \{-1, 0, 1\}$.