

CIS 607 Project 1

Alexandre Chen 951680846

October 21, 2019

Exercise 3

(a)

Did it.

(b)

For $a > 0$, the spatial discretization is given by

$$U_j^{n+1} = U_j^n - \frac{ak}{2h} (3U_j^n - 4U_{j-1}^n + U_{j-2}^n) + \frac{a^2k^2}{2h^2} (U_j^n - 2U_{j-1}^n + U_{j-2}^n)$$

let $v = ak/h$, the stability condition is given by $0 \leq v \leq 2$

(c)

For advection function, there is asymmetry in the equations because the equation models translation at speed a . If $a > 0$, the solution moves rightward. If $a < 0$, the solution moves leftward. For this reason, we would like to use the upwind method which can capture this asymmetry from the equation.

(d)

I modified the “FE_MOL_BS.jl” file for this exercise. The core code that I modified is given by

```
A = zeros(N-1,N-1)
A1 = zeros(N-1,N-1)
for j in 1:N-1
    A1[j , j] = 1
end

A2 = zeros(N-1,N-1)
for j in 1:N-1
    A2[j , j] = 3
end
for j in 1:N-2
    A2[j , j+1] = 1
end

for j in 2:N-1
```

```

    A2[j , j -1] = -4
end

A3 = zeros (N-1,N-1)

for j in 1:N-1
    A3[j , j] = 1
end
for j in 1:N-2
    A3[j , j] = 1
end
for j in 2:N-1
    A3[j , j -1] = -2
end


```

```


A = A1 + (-a*k/(2h))*A2 + (a^2*k^2/(2*h^2))*A3

```

Given our parameter, $a = 0.5, h = 0.05, \lambda = k/h$, the theoretical stable result for λ given by (b) should be $0 \leq \lambda \leq 4$. My test shows that for $\lambda = 0.5, 2$, The numerical results are good. At time $t = 1, t = 2$, the graph plotted is a just a sine wave traslated rightward. But for $\lambda = 4$, the numerical result is not good. Starting from $t = 1$, the graph has been twisted.



(a) $t=1$



(b) $t=2$

Figure 1: $\lambda = 4$

The stability result is obtained given infinitesimal grid width. We are using a rather coarse grid size, this could be the reason why we don't have stable numerical result at $\lambda = 4$. I tested $h = 0.01$, and with this grid size, we can obtain stable numerical result at $\lambda = 4$