
Held–Suarez climate: hs-T42L20-prod-002

T42 retained (stored l_max 43) · L20 uniform σ · 64 × 128 Gaussian state grid · dt 720 s · centred damping
Sections: equal-weight mean of 100 saved states, 200 < day ≤ 1200 (default averaging window); days 210–1200, every 10 days; coverage complete (all 100 nominal save days of the window, every 10 d).


Mean zonal wind [u]

data -23.9 to 32.6 m s⁻¹




Mean temperature [T]

data 189 to 306 K




Eddy momentum flux [u*v*]

data -73.1 to 71.8 m² s⁻²




Eddy heat flux [v*T*]

data -21.8 to 22.2 K m s⁻¹




Eddy kinetic energy ½[u*² + v*²]

data 1.96 to 394 m² s⁻²




Eddy temperature variance [T*²]

data 4.0e-02 to 43.1 K²




Mass: global-mean p_s

max |Δ| 2.02e-06 (day 1200)




Kinetic energy

area- and σ-weighted (not mass-weighted)
window mean 130 (105–148)



Temperature

area- and σ-weighted (not mass-weighted)
window mean 246.26 K (246.14–246.50)



Daily global means of every simulated day (series.json), not the 100 saved states; grey = the sections' window. Mass is conserved by the dry forcing; kinetic energy and temperature are not, and show whether the climate has settled (Gaussian-area means of Δσ-weighted level sums, no p_s factor). Descriptive: the run's acceptance verdicts are in its report.

[ ] = zonal mean on a native σ level, * = departure from that state's [ ]; fixed scales HS_SECTION_SCALES; beyond-scale values hatched.
Definitions and sampling rules: docs/held_suarez/CLIMATE_FIGURE.md.
config sha256 d1fa645db120… · run commit a52ea20550c1 · states sha256 5547924a5df6… · series sha256 d02c6d53534d… · plotted with palintropos 29340e5a