
Held–Suarez climate: hs-T42L20-prod-002

T42 retained (stored l_max 43) · L20 uniform σ · 64 × 128 Gaussian state grid · dt 720 s · centred damping
Time mean: equal-weight mean of 100 saved states with 200 < day ≤ 1200 (default: spinup_days < day ≤ spinup_days + n_blocks × block_days).
Samples: day 210 to day 1200, every 10 days. Coverage: complete (all 100 nominal save days of the window, every 10 d).


Mean zonal wind [u]

data -23.9 to 32.6 m s⁻¹




Mean temperature [T]

data 189 to 306 K




Eddy momentum flux [u*v*]

data -73.1 to 71.8 m² s⁻²




Eddy heat flux [v*T*]

data -21.8 to 22.2 K m s⁻¹




Eddy kinetic energy ½[u*² + v*²]

data 1.96 to 394 m² s⁻²



Eddy temperature variance [T*²]

data 4.0e-02 to 43.1 K²



[ ] = unweighted mean over the 128 equally spaced longitudes at one latitude on one native σ level; * = departure from that state's instantaneous [ ]. Eddy products are formed for each state, then averaged.
All six panels come from the same states. No trapezoidal weighting, interpolation or gap filling. σ-level means are not pressure-level means. The σ axis spans the full levels 0.025 to 0.975 (minor ticks).
u, v, T synthesized exactly from the saved coefficients on the run's Gaussian latitudes; measured zonal bandlimit m_max = 42 < nlon/2, so longitude means of products are exact. Fixed scales: HS_SECTION_SCALES; values beyond the scale are hatched.
config sha256 d1fa645db120… · run commit a52ea20550c1 · protocol sha256 26a8ac31f6da… · states sha256 5547924a5df6… · plotted with palintropos e1f283ed + local changes