

Remote Sensing Analysis of Mangrove Resilience and Cover Change in the Grand-Pierre Bay, Artibonite, Haiti



Alexandre E. S. Georges¹, Mark T. Stacey¹, Deanesh Ramsewak²


¹University of California, Berkeley, ²University of Trinidad & Tobago

¹alexandre_georges@berkeley.edu



Background and Motivation

- Mangroves as Natural Coastal Defense in the Caribbean
- Island Nations in the Caribbean, such as Haiti, are severely threatened by sea-level rise and coastal flooding due to hurricanes' increased frequency and strength.
- Mangroves have the potential to be part of a natural infrastructure strategy for these islands to protect their coastal communities.
- Mangrove forests are, unfortunately, under both climate and anthropogenic pressures.
- One such forest is the Grand-Pierre Bay Mangrove Forest in Haiti, the most extensive mangrove cover in the country.



Data Sources

Satellite Imagery from PlanetLabs is used to conduct this study. We looked at imagery between January 2010 and January 2020 taken by PlanetLabs' RapidEye satellite.

- These images feature high spatial resolution (5 meters) and near-daily temporal resolution, letting us find frequent cloud-free observations to classify and analyze.
- RapidEye (5m) features only five bands (**Red, Green, Blue, Red Edge, and Near Infrared**); however, this is adequate for the indices used and to accurately classify land covers.

The indices used are **NDVI**, **NDWI**, and **UVVR**, as defined below. A Histogram-Based Gradient Boosting Classifier by scikit-learn is used for land classification.

¹Planet Team (2017). Planet Application Program Interface: In Space for Life on Earth. San Francisco, CA. <https://api.planet.com>.

Indices and Models

• Normalized Difference Vegetation Index (NDVI)

$$NDVI = \frac{NIR - Red}{NIR + Red}$$

Quantifies the density and health of vegetation, providing insights in vegetation health changes.

• Normalized Difference Water Index (NDWI)

$$NDWI = \frac{Green - NIR}{Green + NIR}$$

Assesses the presence and extent of surface water bodies.

• Unvegetated to Vegetated Ratio (UVVR)

Helps establish vegetation cover status and tracking changes in wetlands.


• Histogram-based Gradient Boosting (HGB) Classifier²

Ensemble algorithm combining the boosting technique with histogram-based binning. It achieves high-performance and efficient classification. We use it for land classification.

²Scikit-learn: Machine Learning in Python, Pedregosa et al., JMLR 12, pp. 2825-2830, 2011.

How has the GP Mangrove Forest responded to external pressure?

Fig. 1 Mangrove dNDVI between 01-23-2010 and 01-20-2020



• Seaward Retreat, Landward Advance

While the sea-facing region (*seaward*) denotes a consistent drop in NDVI between 2010 and 2020, mangroves in the lagoon-facing region (*landward*) exhibit disperse pockets of NDVI gain. More interior regions show no change to small negative change.

This may indicate mangrove diebacks towards the sea, while new establishment or vegetation reinforcement is occurring towards the lagoon.





Fig. 3 Average NDVI timeseries



• What is driving this decline?

Looking at the timeseries of NDVI and UVVR for both seaward and landward regions shows a period of decreased vegetation health and an increase in the unvegetated ratio between 2013 and 2016.

This matches with an extended drought period experienced in Haiti during the same period. However, the landward region is recovering better from this drought than the seaward one.

Mangrove Cover Change between 01-23-2010 and 01-20-2020





Fig. 5

Fig. 4 Unvegetated to Vegetated Ratio Timeseries



• The forest is migrating landward, but not fast enough


As seen in Fig. 2, the landward section shows increased mangrove cover and distribution, indicating forest expansion, while the seaward section experiences reduced cover but higher distribution, suggesting decreased forest density.

With the landward section being smaller than the seaward, the landward migration is insufficient to offset the seaward

Methods and Workflow

Data Acquisition


Jan. 2010 - Jan. 2020
(Red, Blue, Green, RedEdge, NIR, NDVI, NDWI)



Model Training


Hist. Gradient Boosting Classifier
(Red, Blue, Green, RedEdge, NIR, NDVI, NDWI)

Water Masked NDVI



NDVI Timeseries

Land Classification



Mangrove Cover Mask
Mudflat Cover Mask

Cover Change Map

2MOA

UVVR Timeseries

Trajectories

Mangrove Cover Timeseries*

Future Work

- Future work on remote sensing analysis will involve validating our classifying model using ground truth data from Trinidad, expanding the analysis to other important mangrove sites in the Caribbean, and conducting a comparative study of dynamics between different Caribbean sites.

- In addition to remote sensing, using mangrove cover maps and changes in coupled hydrodynamic and dynamic vegetation models will enable the study of mangrove response to sea-level rise and provide insights into the processes driving seaward retreat.

Acknowledgements

- UC Berkeley EFMH
- PlanetLabs
- Hearts To Humanity (H2H8)

