THE FIBRES OF THE COMPLETION MAP OF A P-RING HAVE PROPERTY P

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied from the pinned Stacks Project chapter file more-algebra.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. The chapter file pinned-original-source.tex carries only \input{preamble} on its first line, so no preamble text lives inside the chapter. The retained file sources/stacks-more-algebra/retained-preamble.tex is a mechanical, line-for-line copy of the project's single shared preamble as staged in the snapshot (Xy-pic setup plus theorem declarations and mathematical macros). Nothing was added, removed or redefined, and the excerpt serves only as preamble, never as a span coordinate.

Editorial note. Both counted spans start at the \begin line of their environment, so the statement and the author's proof are delivered whole; surrounding prose of the section is deliberately outside the counted spans and is carried as standard prerequisites instead.

Editorial note. The citation fields (source_id, source_version, authors, source_url, license.name, source_title) are identical to those of every other card this lane has produced from this chapter, so the batch selector treats them as one source.

Allowed standard prerequisites (editorial).

- Flatness and Noetherianity of completions. Algebra, Lemma \ref{algebra-lemma-completion-flat} and Algebra, Lemma \ref{algebra-lemma-completion-Noetherian-Noetherian}, cited at proof lines 13055-13058. Let the author verify the local criterion for property P for the completion map.
- The local criterion for property P: it is enough to check the third condition on a flat map whose source is Noetherian. Lemma \ref{lemma-P-local}, cited at proof lines 13059-13060. Reduces the claim to the behaviour at a maximal ideal of the completion.
- The ideal of definition is contained in every maximal ideal of the completion, and the quotient of the completion by it is the quotient of the ring. Algebra, Lemma \ref{algebra-lemma-radical-completion}, cited at proof line 13063. Identifies the residue fields of the two local rings and shows that the prime below is maximal.
- A flat local map of Noetherian local rings identifying residue fields and with equal maximal ideals induces an isomorphism on completions. Lemma \ref{lemma-flat-unramified}, cited at proof lines 13073-13074. Identifies the completion of the localised completion with the completion of the localisation.
- The completion map of a local ring is faithfully flat, and property P for fibres is preserved by the composition witnessed by a faithfully flat map. Algebra, Lemma \ref{algebra-lemma-completion-faithfully-flat}, cited at proof lines 13077-13079, and condition (D), used at proof lines 13080-13083.

Date: stacks-project:more-algebra.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

Allow the permanence condition (D) to be applied to the two-step chain of ring maps.

Source reference. Original lemma 0BK9: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 14552–14552. The lemma's own canonical tag, quoted so that readers can resolve the label inside the delivered text; it points at the counted statement itself and carries no proof content.

Source reference. Original lemma 00MB: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 817–817. Navigation pointer to a statement the counted proof refers to but does not contain; nothing load-bearing is delegated to it.

Source reference. Original lemma 0316: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 3973–3973. Navigation pointer to a statement the counted proof refers to but does not contain; nothing load-bearing is delegated to it.

Source reference. Original lemma 0BK8: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 14551–14551. Navigation pointer to a statement the counted proof refers to but does not contain; nothing load-bearing is delegated to it.

Source reference. Original lemma 05GI: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 7062–7062. Navigation pointer to a statement the counted proof refers to but does not contain; nothing load-bearing is delegated to it.

Source reference. Original lemma 0AGX: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 13210–13210. Navigation pointer to a statement the counted proof refers to but does not contain; nothing load-bearing is delegated to it.

Source reference. Original lemma 00MC: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 818–818. Navigation pointer to a statement the counted proof refers to but does not contain; nothing load-bearing is delegated to it.

Lemma 0.2. *Let A be a P-ring where P satisfies (B) and (D). Let $I \subset A$ be an ideal and let $A^\wedge$ be the completion of A with respect to I. Then the fibres of $A \to A^\wedge$ have P.*

Proof. The ring map $A \to A^\wedge$ is flat by Algebra, Lemma 00MB. The ring $A^\wedge$ is Noetherian by Algebra, Lemma 0316. Thus it suffices to check the third condition of Lemma 0BK8. Let $\mathfrak{m}' \subset A^\wedge$ be a maximal ideal lying over $\mathfrak{m} \subset A$. By Algebra, Lemma 05GI we have $IA^\wedge \subset \mathfrak{m}'$. Since $A^\wedge/IA^\wedge = A/I$ we see that $I \subset \mathfrak{m}$, $\mathfrak{m}/I = \mathfrak{m}'/IA^\wedge$, and $A/\mathfrak{m} = A^\wedge/\mathfrak{m}'$. Since $A^\wedge/\mathfrak{m}'$ is a field, we conclude that $\mathfrak{m}$ is a maximal ideal as well. Then $A_\mathfrak{m} \to A^\wedge_{\mathfrak{m}'}$ is a flat local ring homomorphism of Noetherian local rings which identifies residue fields and such that $\mathfrak{m}A^\wedge_{\mathfrak{m}'} = \mathfrak{m}'A^\wedge_{\mathfrak{m}'}$. Thus it induces an isomorphism on complete local rings, see Lemma 0AGX. Let $(A_\mathfrak{m})^\wedge$ be the completion of $A_\mathfrak{m}$ with respect to its maximal ideal. The ring map

$$(A^\wedge)_{\mathfrak{m}'} \to ((A^\wedge)_{\mathfrak{m}'})^\wedge = (A_\mathfrak{m})^\wedge$$

is faithfully flat (Algebra, Lemma 00MC). Thus we can apply (D) to the ring maps

$$A_\mathfrak{m} \to (A^\wedge)_{\mathfrak{m}'} \to (A_\mathfrak{m})^\wedge$$

to conclude because the fibres of $A_\mathfrak{m} \to (A_\mathfrak{m})^\wedge$ have P as A is a P-ring. □