STACKS, DEFORMATION THEORY: LEMMA-INF-MAP-RINGED-TOPOI

THE STACKS PROJECT AUTHORS

Editorial note. Statement span is lines 2787-2820 of sources/stacks-defos/pinned-original-source.tex; proof span is lines 2822-2836, the complete author proof (13 body lines).

Editorial note. Navigation stubs: 2 cited-but-not-included labels are supplied as non-load-bearing pointers with their pinned tags where the snapshot tag index has them.

Editorial note. extra_packages is empty on purpose: the retained preamble already loads xy with options and a second declaration clashes in the assembler wrapper.

Editorial note. Text is retained verbatim, including the author's own typos and placeholders.

Editorial note. Queue index 43 of queue_defos.json (pb=632, stmt 2787-2820, proof 2822-2836).

Allowed standard prerequisites (editorial).

- Explicit prerequisite boundary. The cited inputs are prerequisites; the argument in the retained span is not replaced by any cited result.

Source reference. Original lemma 08MB: sources/stacks-defos/pinned-original-source.tex, source lines 2770–2772. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 08M3: sources/stacks-defos/pinned-original-source.tex, source lines 2836–2836. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Lemma 0.1. *Let $i : (Sh(\mathcal{C}), \mathcal{O}) \to (Sh(\mathcal{D}), \mathcal{O}')$ be a first order thickening of ringed topoi. Assume given extensions*

$$0 \to \mathcal{K} \to \mathcal{F}' \to \mathcal{F} \to 0 \quad \text{and} \quad 0 \to \mathcal{L} \to \mathcal{G}' \to \mathcal{G} \to 0$$

as in (08MB) and maps $\varphi : \mathcal{F} \to \mathcal{G}$ and $\psi : \mathcal{K} \to \mathcal{L}$.

(1) *If there exists an $\mathcal{O}'$-module map $\varphi' : \mathcal{F}' \to \mathcal{G}'$ compatible with φ and ψ, then the diagram*

$$\begin{array}{ccc} \mathcal{I} \otimes_{\mathcal{O}} \mathcal{F} & \xrightarrow[c_{\mathcal{F}'}]{} & \mathcal{K} \\ {\scriptstyle 1\otimes\varphi}\downarrow & & \downarrow{\scriptstyle \psi} \\ \mathcal{I} \otimes_{\mathcal{O}} \mathcal{G} & \xrightarrow{c_{\mathcal{G}'}} & \mathcal{L} \end{array}$$

is commutative.

(2) *The set of $\mathcal{O}'$-module maps $\varphi' : \mathcal{F}' \to \mathcal{G}'$ compatible with φ and ψ is, if nonempty, a principal homogeneous space under $\mathrm{Hom}_{\mathcal{O}}(\mathcal{F}, \mathcal{L})$.*

Date: stacks-project:defos.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

Proof. Part (1) is immediate from the description of the maps. For (2), if φ' and φ'' are two maps $\mathcal{F}' \to \mathcal{G}'$ compatible with φ and ψ, then $\varphi' - \varphi''$ factors as

$$\mathcal{F}' \to \mathcal{F} \to \mathcal{L} \to \mathcal{G}'$$

The map in the middle comes from a unique element of $\mathrm{Hom}_{\mathcal{O}}(\mathcal{F}, \mathcal{L})$ by Modules on Sites, Lemma 08M3. Conversely, given an element α of this group we can add the composition (as displayed above with α in the middle) to φ'. Some details omitted. $\square$