STACKS, DEFORMATION THEORY: LEMMA-INF-OBS-MAP-RINGED-TOPOI

THE STACKS PROJECT AUTHORS

Editorial note. Statement span is lines 2838-2868 of sources/stacks-defos/pinned-original-source.tex; proof span is lines 2870-2909, the complete author proof (38 body lines).

Editorial note. Navigation stubs: 1 cited-but-not-included labels are supplied as non-load-bearing pointers with their pinned tags where the snapshot tag index has them.

Editorial note. extra_packages is empty on purpose: the retained preamble already loads xy with options and a second declaration clashes in the assembler wrapper.

Editorial note. Text is retained verbatim, including the author's own typos and placeholders.

Editorial note. Queue index 26 of queue_defos.json (pb=1136, stmt 2838-2868, proof 2870-2909).

Allowed standard prerequisites (editorial).

- Explicit prerequisite boundary. The cited inputs are prerequisites; the argument in the retained span is not replaced by any cited result.

Source reference. Original lemma 08MB: sources/stacks-defos/pinned-original-source.tex, source lines 2770–2772. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Lemma 0.1. *Let $i : (Sh(\mathcal{C}), \mathcal{O}) \to (Sh(\mathcal{D}), \mathcal{O}')$ be a first order thickening of ringed topoi. Assume given extensions*

$$0 \to \mathcal{K} \to \mathcal{F}' \to \mathcal{F} \to 0 \quad \text{and} \quad 0 \to \mathcal{L} \to \mathcal{G}' \to \mathcal{G} \to 0$$

as in (08MB) and maps $\varphi : \mathcal{F} \to \mathcal{G}$ and $\psi : \mathcal{K} \to \mathcal{L}$. Assume the diagram

$$\begin{array}{ccc} \mathcal{I} \otimes_{\mathcal{O}} \mathcal{F} & \xrightarrow{c_{\mathcal{F}'}} & \mathcal{K} \\ {\scriptstyle 1\otimes\varphi}\downarrow & & \downarrow{\scriptstyle \psi} \\ \mathcal{I} \otimes_{\mathcal{O}} \mathcal{G} & \xrightarrow{c_{\mathcal{G}'}} & \mathcal{L} \end{array}$$

is commutative. Then there exists an element

$$o(\varphi, \psi) \in \mathrm{Ext}^1_{\mathcal{O}}(\mathcal{F}, \mathcal{L})$$

whose vanishing is a necessary and sufficient condition for the existence of a map $\varphi' : \mathcal{F}' \to \mathcal{G}'$ compatible with φ and ψ.

Proof. We can construct explicitly an extension

$$0 \to \mathcal{L} \to \mathcal{H} \to \mathcal{F} \to 0$$

Date: stacks-project:defos.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

by taking $\mathcal{H}$ to be the cohomology of the complex

$$\mathcal{K} \xrightarrow{1,-\psi} \mathcal{F}' \oplus \mathcal{G}' \xrightarrow{\varphi,1} \mathcal{G}$$

in the middle (with obvious notation). A calculation with local sections using the assumption that the diagram of the lemma commutes shows that $\mathcal{H}$ is annihilated by $\mathcal{I}$. Hence $\mathcal{H}$ defines a class in

$$\mathrm{Ext}^1_{\mathcal{O}}(\mathcal{F}, \mathcal{L}) \subset \mathrm{Ext}^1_{\mathcal{O}'}(\mathcal{F}, \mathcal{L})$$

Finally, the class of $\mathcal{H}$ is the difference of the pushout of the extension $\mathcal{F}'$ via ψ and the pullback of the extension $\mathcal{G}'$ via φ (calculations omitted). Thus the vanishing of the class of $\mathcal{H}$ is equivalent to the existence of a commutative diagram

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \mathcal{K} & \longrightarrow & \mathcal{F}' & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle \psi} & & \downarrow{\scriptstyle \varphi'} & & \downarrow{\scriptstyle \varphi} & & \\
0 & \longrightarrow & \mathcal{L} & \longrightarrow & \mathcal{G}' & \longrightarrow & \mathcal{G} & \longrightarrow & 0
\end{array}$$

as desired. □