A REPRESENTABLE MORPHISM IS UNIVERSALLY INJECTIVE EXACTLY WHEN ALL ITS BASE CHANGES ARE INJECTIVE ON SPACES

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied from the pinned Stacks Project chapter source sources/stacks-algebraic-stacks-morphisms/pinned-original-source.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. Preamble provenance: the Stacks Project keeps one single shared preamble for every chapter; this card takes it directly from the staged sources/stacks-algebraic-stacks-morphisms/preamble.tex as three verbatim spans (lines 8-12, 14-18 and 154-217), so no derived file was written and nothing was added or redefined. The wrapper class and the base packages are supplied by the assembler.

Editorial note. Bibliography provenance: neither the counted statement nor its proof contains a citation (this lane rescans the whole span before writing the card), so the card carries no bibliography and no entry was reconstructed.

Editorial note. Selection provenance: this card was opened under the file-based control contract (CONTROL.md), with the mechanical pre-card gate run on its statement and proof interval before the card was written.

Allowed standard prerequisites (editorial).

- The definition of a universally injective representable morphism of algebraic stacks, tested after base change by schemes. Referred to in the statement (source lines 2839-2840) as a pointer to the chapter on properties of stacks. Formulates the two conditions being compared.
- The description of the topological space of an algebraic stack through a smooth cover, with the induced maps open and surjective and the induced map to the fibre product of the spaces surjective. Invoked inside the counted proof (source lines 2852-2866) as a pointer to the same chapter. Descends injectivity from a smooth cover to the base change of the stack.

Source reference. Original reference stacks-properties-section-points: sources/stacks-algebraic-stacks-morphisms/pinned-original-source.tex, source lines 2488–2488. Navigation pointer only: the counted text refers to the Stacks item stacks-properties-section-points, which belongs to another chapter of the project and is not part of this card. The pointer carries the target's own label as its number because this chapter's staged snapshot does not contain the other chapter's tag table; its content is not reproduced here.

Source reference. Original reference stacks-properties-section-properties-morphisms: sources/stacks-algebraic-stacks-morphisms/pinned-original-source.tex, source lines 248–248. Navigation pointer only: the counted text refers to the Stacks item stacks-properties-section-properties-morphisms, which belongs to another chapter of the project and is not part of this card. The pointer carries the target's own label as its number because this chapter's staged snapshot does not contain the other chapter's tag table; its content is not reproduced here.

Date: stacks-project:stacks-morphisms.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

Lemma 0.1. *Let $f : \mathcal{X} \to \mathcal{Y}$ be a morphism of algebraic stacks which is representable by algebraic spaces. The following are equivalent*

(1) *f is universally injective (as in Properties of Stacks, Section stacks-properties-section-properties-morphisms), and*
(2) *for every morphism of algebraic stacks $\mathcal{Z} \to \mathcal{Y}$ the map $|\mathcal{Z} \times_{\mathcal{Y}} \mathcal{X}| \to |\mathcal{Z}|$ is injective.*

Proof. Assume (1), and let $\mathcal{Z} \to \mathcal{Y}$ be as in (2). Choose a scheme V and a surjective smooth morphism $V \to \mathcal{Z}$. By assumption the morphism $V \times_{\mathcal{Y}} \mathcal{X} \to V$ of algebraic spaces is universally injective, in particular the map $|V \times_{\mathcal{Y}} \mathcal{X}| \to |V|$ is injective. By Properties of Stacks, Section stacks-properties-section-points in the commutative diagram

$$\begin{array}{ccc} |V \times_{\mathcal{Y}} \mathcal{X}| & \longrightarrow & |\mathcal{Z} \times_{\mathcal{Y}} \mathcal{X}| \\ \downarrow & & \downarrow \\ |V| & \longrightarrow & |\mathcal{Z}| \end{array}$$

the horizontal arrows are open and surjective, and moreover

$$|V \times_{\mathcal{Y}} \mathcal{X}| \longrightarrow |V| \times_{|\mathcal{Z}|} |\mathcal{Z} \times_{\mathcal{Y}} \mathcal{X}|$$

is surjective. Hence as the left vertical arrow is injective it follows that the right vertical arrow is injective. This proves (2). The implication (2) $\Rightarrow$ (1) follows from the definitions. $\square$