FUNCTORIALITY OF THE CONORMAL SHEAF FOR IMMERSIONS OF ALGEBRAIC SPACES

THE STACKS PROJECT AUTHORS

Editorial note. Statement span is lines 454-471 of sources/stacks-spaces-more-morphisms/pinned-original-source.tex; proof span is lines 473-517, the complete author proof (43 body lines).

Editorial note. Navigation stubs: 1 cited-but-not-included labels are supplied as non-load-bearing pointers with their pinned tags where the snapshot tag index has them.

Editorial note. extra_packages is empty on purpose: the retained preamble already loads xy with options and a second declaration clashes in the assembler wrapper.

Editorial note. Text is retained verbatim, including the author's own typos and placeholders.

Allowed standard prerequisites (editorial).

- Explicit prerequisite boundary. Conormal sheaves of immersions, ideal sheaves and their square-zero quotients, and pullback along morphisms of algebraic spaces; the construction is retained.

Source reference. Original lemma spaces-morphisms-lemma-closed-immersion-ideals: sources/stacks-spaces-more-morphisms/pinned-original-source.tex, source lines 517–517. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Lemma 0.1. *Let S be a scheme. Let*

$$\begin{array}{ccc} Z & \xrightarrow{i} & X \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle g} \\ Z' & \xrightarrow{i'} & X' \end{array}$$

be a commutative diagram of algebraic spaces over S. Assume i, i' immersions. There is a canonical map of $\mathcal{O}_Z$-modules

$$f^*\mathcal{C}_{Z'/X'} \longrightarrow \mathcal{C}_{Z/X}$$

Proof. First find open subspaces $U' \subset X'$ and $U \subset X$ such that $g(U) \subset U'$ and such that $i(Z) \subset U$ and $i(Z') \subset U'$ are closed (proof existence omitted). Replacing X by U and X' by U' we may assume that i and i' are closed immersions. Let $\mathcal{I}' \subset \mathcal{O}_{X'}$ and $\mathcal{I} \subset \mathcal{O}_X$ be the quasi-coherent sheaves of ideals associated to i' and i, see Morphisms of Spaces, Lemma spaces-morphisms-lemma-closed-immersion-ideals. Consider the composition

$$g^{-1}\mathcal{I}' \to g^{-1}\mathcal{O}_{X'} \xrightarrow{g^\sharp} \mathcal{O}_X \to \mathcal{O}_X/\mathcal{I} = i_*\mathcal{O}_Z$$

Date: stacks-project:spaces-more-morphisms.tex:local-snapshot-20261011; GNU FDL 1.2 or later.

Since $g(i(Z)) \subset Z'$ we conclude this composition is zero (see statement on factorizations in Morphisms of Spaces, Lemma spaces-morphisms-lemma-closed-immersion-ideals). Thus we obtain a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{I} & \longrightarrow & \mathcal{O}_X & \longrightarrow & i_*\mathcal{O}_Z & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \uparrow & & \\ 0 & \longrightarrow & g^{-1}\mathcal{I}' & \longrightarrow & g^{-1}\mathcal{O}_{X'} & \longrightarrow & g^{-1}i'_*\mathcal{O}_{Z'} & \longrightarrow & 0 \end{array}$$

The lower row is exact since g^{-1} is an exact functor. By exactness we also see that $(g^{-1}\mathcal{I}')^2 = g^{-1}((\mathcal{I}')^2)$. Hence the diagram induces a map $g^{-1}(\mathcal{I}'/(\mathcal{I}')^2) \to \mathcal{I}/\mathcal{I}^2$. Pulling back (using i^{-1} for example) to Z we obtain $i^{-1}g^{-1}(\mathcal{I}'/(\mathcal{I}')^2) \to \mathcal{C}_{Z/X}$. Since $i^{-1}g^{-1} = f^{-1}(i')^{-1}$ this gives a map $f^{-1}\mathcal{C}_{Z'/X'} \to \mathcal{C}_{Z/X}$, which induces the desired map. $\square$