Bounded-rank tensor isomorphism has rank-dependent Nullstellensatz and finite-field polynomial-calculus degree bounds

Nicola Galesi, Joshua A. Grochow, Toniann Pitassi, Adrian She

doi:10.4230/LIPIcs.CCC.2023.4; CC BY 4.0

Editorial note. Public sources contain only the corresponding official published complete author material; native TeX served as editable transcription aid. Raw native source/tar remains private. Original theorem numbering preserved with section counters. Rank and inversion consequences counted as a single result, and the alternative Random3XOR tensor lower-bound proof is not a new record.

Editorial note. References to unselected original sections are printed with their accurate original article chapter numbers; selected section headers and original labels remain in place. No absent section is presented as an included proof.

Allowed standard prerequisites (editorial).

- Known graph-isomorphism and functional-pigeonhole polynomial-calculus lower bounds; effective Nullstellensatz and tensor-rank inheritance. Exact original cited statements retained where used; standard earlier literature not recursively reproved

Abstract

The Tensor Isomorphism problem (TI) has recently emerged as having connections to multiple areas of research within complexity and beyond, but the current best upper bound is essentially the brute force algorithm. Being an algebraic problem, TI (or rather, proving that two tensors are *non*-isomorphic) lends itself very naturally to algebraic and semi-algebraic proof systems, such as the Polynomial Calculus (PC) and Sum of Squares (SoS). For its combinatorial cousin Graph Isomorphism, essentially optimal lower bounds are known for approaches based on PC and SoS (Berkholz & Grohe, SODA '17). Our main results are an $\Omega(n)$ lower bound on PC degree or SoS degree for Tensor Isomorphism, and a nontrivial upper bound for testing isomorphism of tensors of bounded rank.

We also show that PC cannot perform basic linear algebra in sublinear degree, such as comparing the rank of two matrices, or deriving $BA = I$ from $AB = I$. As linear algebra is a key tool for understanding tensors, we introduce a strictly stronger proof system, PC+Inv, which allows as derivation rules all substitution instances of the implication $AB = I \to BA = I$. We conjecture that even PC+Inv cannot solve TI in polynomial time either, but leave open getting lower bounds on

PC+Inv for any system of equations, let alone those for TI. We also highlight many other open questions about proof complexity approaches to TI.

1 Introduction

Tensors have rapidly emerged as a fundamental data structure and key mathematical object of the 21st century. They play key roles in many different areas of science, engineering, and mathematics, from quantum mechanics and general relativity to neural networks [NPOV15] and mechanical engineering. They arise in theoretical computer science in many ways, including from (post-quantum) cryptography [Pat96, JQSY19], derandomization, Matrix Multiplication, Graph Isomorphism [GQ21b], and several different parts of Geometric Complexity Theory.

The fundamental notion of equivalence between tensors is that of isomorphism: two tensors are isomorphic if one can be transformed into the other by an invertible linear change of basis in each of the corresponding vector spaces. For example, two 2-tensors (=matrices) M, M' are equivalent under this notion if there are invertible matrices X, Y such that $XMY = M'$; similarly, two 3-tensors, represented by 3-way arrays T_{ijk}, T'_{ijk} are isomorphic if there are three invertible matrices X, Y, Z such that

$$\sum_{ijk} X_{ii'} Y_{jj'} Z_{kk'} T_{ijk} = T'_{i'j'k'} \tag{1}$$

for all i', j', k'. The problem of (3-)Tensor Isomorphism (TI) is: given two such 3-way arrays, to decide if they are isomorphic.

Over finite fields, two different versions of TI sandwich the complexity of its more famous cousin, Graph Isomorphism. Namely, as presented above, GI reduces to TI. In the other direction, over a finite field $\mathbb{F} = \mathbb{F}_{p^a}$, one can take an $n \times n \times n$ tensor and list it out "verbosely", as a set of p^{an} many $n \times n$ matrices over $\mathbb{F}$; the isomorphism problem for such verbosely given tensors is equivalent to Group Isomorphism for a certain class of p-groups, widely believed to be the hardest cases of Group Isomorphism in general. As such, this verbose version of TI reduces to GI. Furthermore, with Babai's quasi-polynomial-time algorithm [Bab16], the running times are quite close: $N^{O(\log N)}$ for VerboseTI and $N^{O(\log^2 N)}$ for GI (the exponent of the exponent was worked out by Helfgott [Hel19]). Thus TI stands as a key obstacle to putting GI into P.

In this paper, we initiate the study of (algebraic) proof complexity approaches to proving that two tensors are non-isomorphic. Lower bounds on the Polynomial Calculus proof system imply lower bounds on Gröbner basis techniques, and the latter are some of the leading methods for solving TI-complete problems in cryptanalysis, e.g., [TDJ+22, FP06]. In the context of GI, proof complexity plays an important role, through its connection with the Weisfeiler–Leman (WL) algorithm. Although this algorithm does not, on its own, solve GI in polynomial time [CFI92], it is a key subroutine in many of the best algorithms for GI, both in theory [Bab16] and in practice (see [McK81, MP14]). And the picture that has emerged is that some proof systems for GI are known to be equivalent in power to WL [AM13, BG15], and some lower bounds on proof systems are closely related to lower bounds for WL [SSC14, OWWZ14, BG15].

Versions of WL for groups, and in particular finite p-groups—and hence, by the connection above, tensors over finite fields—have only recently begun to be explored [BGL$^+$19, BS20, BS22, CL22].

1.1 Main results

We focus on the Polynomial Calculus (PC, or Gröbner) proof system [CEI96], though our results will also hold for semi-algebraic proof systems such as Sum-of-Squares [Las01] as well. PC is used to show that a system of polynomial equations over a field $\mathbb{F}$ is unsatisfiable over the algebraic closure $\overline{\mathbb{F}}$, by deriving from the system of equations, in a line-by-line fashion, the contradiction $1 = 0$. The degree of a PC proof is the maximum degree of any line appearing in the proof, and it is a fundamental result that PC proofs of constant degree can be found in polynomial time [CEI96]. Much as WL informally "captures all combinatorial approaches" to GI, PC informally "captures all approaches based on Gröbner bases" to showing that a system of polynomial equations is unsatisfiable.

The systems of equations we study are, for two non-isomorphic tensors T, T', the equations (1) along with new matrices X', Y', Z', and equations saying that these are the inverses of X, Y, Z, resp., viz.: $XX' = X'X = \mathrm{Id}$, and similarly for the others. The reason for introducing these new matrices, despite their not appearing in (1), is that these invertibility equations are only degree 2. In contrast, if we instead used the determinant to indicate that X was invertible, then our starting equations would have degree $n+1$, rather than constant degree ≤ 3. Since the main complexity measure we study on PC is degree, having starting equations of degree n would make it difficult to make meaningful lower bound statements.

Our first main result is (two proofs of) a lower bound on such techniques.

Theorem 1.1. *Over any field, there are instances of $n \times n \times n$ Tensor Isomorphism that require PC degree $\Omega(n)$ to refute. Over $\mathbb{R}$, they also require Sum-of-Squares degree $\Omega(n)$ to refute.*

The preceding goes by reduction from known lower bounds on PC for Graph Isomorphism [BG15, BG17], but has the disadvantage (from the tensor point of view) that the resulting tensors are quite sparse: in one direction, one of the slices is supported on an $\Omega(n) \times n$ matrix and all the others slices have support size 1. In a second proof (Section 6 (in the original article)), we get a polynomially worse lower bound $\Omega(\sqrt[4]{n})$, but with a reduction from Random 3XOR that is more direct. Indeed, we show that 3XOR itself can be viewed as a particular instance of a tensor problem *without* gadgets; gadgets are only then needed to reduce from that tensor problem to Tensor Isomorphism itself. In contrast, the lower bounds on PC for GI (*ibid.*) already use the Cai–Fürer–Immerman gadgets [CFI92] to reduce from XOR-SAT, and then even further gadgets are needed to reduce from GI to TI.

Our technical contributions in the above theorem are thus three-fold:

1. We show that the known reductions from GI to TI can be carried out in low-degree PC;

2. We realize 3XOR very naturally as a tensor problem; and

3. We give new reductions from 3XOR, through a series of tensor-related problems, to TI, that work as many-one reductions of the decision problems that can be carried out in low-degree PC.

Complementing our lower bound, we also show that tensors of low rank are comparatively easy to test for (non)-isomorphism. Here, one of our upper bounds is in the weaker Nullstellensatz proof system (giving a stronger upper bound than only a PC upper bound). In the Nullstellensatz proof system, a proof that a system of equations $f_1 = \cdots = f_m = 0$ is unsatisfiable consists of polynomials g_i such that $\sum g_i f_i = 1$, and the Nullstellensatz degree is the maximum degree of any $g_i f_i$. The PC degree is always at most the Nullstellensatz degree, and the gap between the two can be nearly maximal for Boolean equations: $O(1)$ versus $\Omega(n/\log n)$ [BCIP02]. (For Boolean equations, there is always an $O(n)$ upper bound, though this does not apply to TI, see Remark 1.3 below).

Theorem 1.2. *Over any field, the Nullstellensatz degree of refuting isomorphism of two $n \times n \times n$ tensors of tensor rank $\leq r$ is at most $2^{O(r^2)}$. If working over a finite field $\mathbb{F}_q$ and including the equations $x^q - x$, the PC degree is at most $O(qr^2)$.*

In particular, isomorphism of constant-rank tensors can be decided in polynomial time.

Remark 1.3. In many settings in proof complexity, Boolean axioms such as $x_i^2 = x_i$ or $x_i^2 = 1$ are included among the system of equations, and all such unsatisfiable systems of equations can be refuted in degree $O(n)$ ($n =$# variables). If this were the case here, the above would only be interesting for very small values of r. In contrast, the equations for TI do not include any such Boolean axioms, and as such the naive degree upper bound is exponential in the number of variables. For $n \times n \times n$ tensors, this gives an upper bound of $2^{O(n^2)}$ [Som99], and thus, Theorem 1.2 gives nontrivial upper bounds all the way up to $r \leq n$. (We note that $n \times n \times n$ tensors can have rank up to $\Theta(n^2)$ [Lic85].) The proof of Theorem 1.2 shows that for rank-r tensors, TI can essentially be reduced to a system of equations in only $O(r^2)$ variables.

Remark 1.4. For fixed r, testing if an $n \times n \times n$ tensor has rank $\leq r$ can be done in polynomial time, as follows. This will show that the algorithm of Theorem 1.2 genuinely solves the decision problem, and not just a promise problem. Given an $n \times n \times n$ tensor T, consider its three $n \times n^2$ flattenings. Use Gaussian elimination to put each such flattening, separately, into reduced row echelon form. If any of these flattenings has rank $> r$, reject. Otherwise, we get from this a list of $3r$ vectors $u_1, \ldots, u_r, v_1, \ldots, v_r, w_1, \ldots, w_r$, such that T lives in the $r \times r \times r$-dimensional space $\mathrm{Span}\{u_1, \ldots, u_r\} \otimes \mathrm{Span}\{v_1, \ldots, v_r\} \otimes \mathrm{Span}\{w_1, \ldots, w_r\}$. Now in this space we can write down the Brent equations [Bre70] for T to have rank $\leq r$, which will be r^3 cubic equations in $3r^2$ variables (Brent's equations [Bre70, (5.06)] were specifically for the matrix multiplication tensor, but analogous equations are easily constructed for arbitrary tensors using the same idea). Since r is constant, these equations may be solved in polynomial time (here we assume that we are either working over a finite field, a finite-degree extension of the rationals—see, for example, Grigoriev [Gri13]—or in the BSS model over an arbitrary field).

Lastly, one may wonder why we focus on 3-Tensor Isomorphism, and not some of its many related variants. Indeed, just as there are other equivalence notions for matrices—such as conjugacy XMX^{-1} and congruence XMX^T—there are many different kinds of multilinear objects that can be represented by multi-way arrays, including tensors, homogeneous polynomials (commutative or noncommutative), alternating matrix spaces, multilinear maps, and so on, each with their own corresponding notion of isomorphism. While these problems are indeed distinct, they are all equivalent under polynomial-time isomorphisms [FGS19, GQ21b]; such problems are called TI-complete. Even isomorphism of k-way tensors (for any fixed $k \geq 3$) is equivalent to isomorphism of 3-tensors [GQ21b]. This partially justifies our focus on 3-Tensor Isomorphism. In the course of proving our reductions for the results stated above, we use many of the gadgets from [FGS19, GQ21b], and show that such uses also often yield proof complexity reductions as well. Because of the variety of gadgets used in our reductions, we believe that many, if not all, of the gadgets from those results would also yield proof complexity reductions, so the proof complexity of all the known TI-complete problems should be polynomially related.

1.2 Comparison with linear algebra, a new proof system, and a conjecture

As linear algebra is part of the core toolkit for understanding tensors, it is natural to wonder how linear algebra can help in algebraic proof complexity approaches to TI. We believe that even if it had the "full power" of linear algebra at its disposal "for free," PC could still not solve TI efficiently. We begin to make this precise in this section.

Some basic derivations in linear algebra are to relate the ranks of two matrices and to derive $BA = I$ from $AB = I$ (the Inversion Principle, one of the so-called "hard matrix identities" [SC04], only recently shown to have short NC^2-Frege proofs [HT15]). Soltys [Sol01] and Soltys & Cook [SC04] discuss the relationship between these and other standard implications in linear algebra. We show that PC is not strong enough to prove these in low-degree:

Theorem 1.5. *The unsatisfiable system of equations $XY = \mathrm{Id}_n$ where X is $n \times r$ and Y is $r \times n$ with $1 \leq r < n$, requires degree $\geq r/2 + 1$ to refute in PC, over any field.*

We refer to this system of equations as the Rank Principle, as refuting them amounts to showing that $\mathrm{rk}\,\mathrm{Id}_n > r$.

Theorem 1.6. *Any PC derivation of $BA = I$ from $AB = I$, where A, B are $n \times n$ matrices with $\{0, 1\}$ entries, requires degree $\geq n/2 + 1$, over any field.*

We also observe that the Rank Principle can be derived in low degree from the Inversion Principle.

Although it remains open whether the Inversion Principle is "complete" for linear-algebraic reasoning (see [Sol01, SC04]), we introduce the proof system PC+Inv in an attempt to capture some linear-algebraic reasoning that seems potentially useful for TI. PC+Inv has all the same derivation rules as PC, but in addition, for any square matrices A, B (whose entries may themselves be

polynomials—that is, we allow substitution instances), we have the rule

$$\frac{AB = I}{BA = I}.$$

where the antecedent represents the set of n^2 equations corresponding to $AB = I$, and similarly the consequent denotes the set of n^2 equations $BA = I$ (see 2.3 for more details). Degree is still measured in the usual way, but this rule lets us "cut out" the high-degree proof that would usually be required to derive $BA = I$ from $AB = I$. We now formalize our intuition that linear algebra should not suffice to solve TI efficiently in the following:

Conjecture 1.7. Tensor Isomorphism for $n \times n \times n$ tensors requires degree $\Omega(n)$ in PC+Inv, over any field.

Despite the conjecture, we do not yet know how to prove lower bounds on PC+Inv for any unsatisfiable system of equations, let alone those coming from TI. Mod p counting principles (for p different from the characteristic of the field) strike us as potentially interesting instances to examine for PC+Inv lower bounds, before tackling a harder problem like TI. In the final section, we highlight many other open questions around the proof complexity of TI.

1.3 Organization

In Section 2 we cover preliminaries. In Section 3 (in the original article) we prove the lower bounds on linear algebraic principles just discussed. In Section 4 we prove the upper bound for isomorphism of bounded rank tensors (Theorem 1.2). In Section 5 (in the original article) we prove Theorem 1.1 by reduction from GI. In Section 6 (in the original article) we prove the polynomially related lower bound by direct reduction from Random 3XOR.

2 Preliminaries

2.1 Proof systems

All our rings are commutative and unital. *Polynomial calculus* (PC) is a proof system to prove that a given system of (multivariate) polynomial equations $\mathcal{P}$ over a field $\mathbb{F}$ of the form $p = 0$, has no solution over the algebraic closure $\overline{\mathbb{F}}$ (i.e. the system is unsolvable). We usually shorten the polynomial equation $p = 0$ to just p. The derivation rules of the system are the following one:

$$\frac{p}{xp} \text{ (multiplication)}, \quad \frac{p \quad q}{ap + bq} \text{ (linear combination)}$$

where x is any formal variable, $a, b \in \mathbb{F}$ and p, q are polynomials over $\mathbb{F}$.

When refuting Boolean systems of equations it is common to include the Boolean axioms $x_i^2 - x_i$. Because we do *not* always include these (esp. for TI) we are explicit about our use of these, but do not assume they are built into the proof system—that is, if we are assuming them as axioms, we say so.

A *PC derivation (or proof)* of a polynomial q from a set of polynomials $\mathcal{P}$ is a sequence of polynomial equations $p_1, \ldots, p_m$ ending with the polynomial q (so p_m is q) and where each p_i, $i \in [m]$, is either an *axiom* p for $p \in \mathcal{P}$, or is

obtained from previous equations in the refutation by multiplication or linear combination. We denote this by writing $\mathcal{P} \vdash q$. Observe that if p is derivable in PC and q is a polynomial then, by repeated applications of multiplication and linear combination rules, we can derive pq. We often use this generalization of the multiplication in our proofs without mention.

A *PC refutation* is just a PC proof of the polynomial 1. The *degree* of a PC derivation is the maximal degree of a polynomial used in the proof. The *size* of a polynomial p is the number of terms in p. The *size* a PC derivation $p_1, \ldots, p_m$ is the sum of the sizes of the polynomials $p_1, \ldots, p_m$.

For our upper bound in Theorem 1.2, we also consider another algebraic proof system, known as *Nullstellensatz* (NS), to certify unsolvability of sets of polynomial equations. Nullstellensatz is defined in a static form as follows: a refutation of a list $\mathcal{P} = (p_1, \ldots, p_m)$ of polynomial equations over variables $x_1 \ldots, x_n$ is given by the list of polynomials $\mathcal{Q} = (q_1, \ldots q_m)$ such that

$$\sum_{i \in [m]} p_i q_i = 1$$

The *degree* of a NS refutation is the maximal degree of a polynomial in $\mathcal{P} \cup \mathcal{Q}$. The size of NS proof is the sum of the number of monomials appearing in the polynomials $q_1, \ldots, q_m$.

Sum-of-Squares (SOS) is a static proof system for certifying the unsolvability of systems of polynomial equations and polynomial inequalities, where polynomials are usually over the ring $\mathbb{R}[x_1 \ldots, x_n]$.

A polynomial p is a *sum-of-squares* polynomial if it is in the form $p = \sum_i r_i^2$ and the r_i's are polynomials as well. Given a system made by a set of polynomial equations $\mathcal{P} = \{p_1 = 0, \ldots p_m = 0\}$ and a set $\mathcal{Q} = \{q_1 \geq 0, \ldots q_k \geq 0\}$ of polynomial inequalities, a *sum-of-squares* proof of the polynomial inequality $p \geq 0$ from $\mathcal{P} \cup \mathcal{Q}$ is given by the formal identity

$$p = s_0 + \sum_{i \in [k]} s_i q_i + \sum_{j \in [m]} t_j p_j$$

where $s_0, s_1, \ldots, s_k$ are sum-of-squares polynomials, while $t_1, \ldots, t_m$ are arbitrary polynomials. When the system $\mathcal{P} \cup \mathcal{Q}$ is unsatisfiable, a *refutation* of $\mathcal{P} \cup \mathcal{Q}$ is a proof of the inequality $-1 \geq 0$, that is for p the constant polynomial -1. The *degree* of the proof is the $\max\{\deg(p), \deg(s_0), \deg(s_i) + \deg(q_i), \deg(t_j) + \deg(p_j) | i \in [k], j \in [m]\}$.

Definition 2.1 (PC reduction between systems of polynomials, cf. [BGIP01, Sec. 3])**.** Let $P(x_1, \ldots, x_n)$ and $Q(y_1, \ldots, y_m)$ be two sets of polynomials over a field $\mathbb{F}$. P is (d_1, d_2)*-reducible* to Q if:

1. For each $i \in [m]$ there is a polynomial $r_i(\mathbf{x})$ of degree at most d_1 (which we think of as defining y_i in terms of the $\mathbf{x}$ variables);
2. There exists a degree d_2 PC derivation of $Q(r_1(\mathbf{x}), \ldots, r_m(\mathbf{x}))$ from polynomials $P(\mathbf{x})$.

Lemma 2.2 ([BGIP01, Lem. 1])**.** *If $P(\mathbf{x})$ is (d_1, d_2)-reducible to $Q(\mathbf{y})$ and there is a degree d PC refutation of $Q(\mathbf{y})$, then there is a degree* $\max(d_2, d_1 d)$ *refutation of $P(\mathbf{x})$.*

In their paper, they typically only applied this to systems of equations which were known to be unsatisfiable (such as PHP and Tseitin tautologies), whereas in our paper we have several situations we want to combine the above notion together with the usual notion of many-one reduction. We encapsulate this in the following definition. We say a decision problem Π is a *polynomial solvability problem* over a field $\mathbb{F}$ if all valid instances of the problem are systems of polynomial equations over $\mathbb{F}$, and the problem is to decide whether such a system of equations has solutions over the algebraic closure $\overline{\mathbb{F}}$. Thus, the difference between multiple polynomial solvability problems is just *which* systems of equations are valid inputs.

Definition 2.3 (PC many-one reduction)**.** Let Π_1, Π_2 be two polynomial solvability problems over a field $\mathbb{F}$. We say that Π_1 (d_1, d_2)*-many-one reduces* to Π_2 if there is a polynomial-time many-one reduction ρ from Π_1 to Π_2, such that for all unsatisfiable instances $\mathcal{F}$ of Π_1, $\mathcal{F}$ (d_1, d_2)-reduces to $\rho(\mathcal{F})$. When this occurs with $d_1, d_2 = O(1)$, we write

$$\Pi_1 \leq_m^{PC} \Pi_2.$$

2.2 Linear algebra and tensors

Given three vector spaces U, V, W over a field $\mathbb{F}$, a 3-tensor is an element of the vector space $U \otimes V \otimes W$, whose dimension is $(\dim U)(\dim V)(\dim W)$. If e_i is the i-th standard basis vector, then a basis for $U \otimes V \otimes W$ is given by the vectors $\{e_i \otimes e_j \otimes e_k\}$. One may also interpret the symbol $\otimes$ more concretely as the Kronecker product, in which $e_i \otimes e_j \otimes e_k$ represents a 3-way array whose only nonzero entry is in the (i, j, k) position. The vector space of such 3-way arrays (with coordinate-wise addition) is isomorphic to $U \otimes V \otimes W$.

The rank of a tensor $T \in U \otimes V \otimes W$ is the minimum r such that $T = \sum_{i=1}^{r} u_i \otimes v_i \otimes w_i$ for some vectors u_i, v_i, w_i.

Two $n \times m \times p$ 3-tensors $T, T' \in U \otimes V \otimes W$ are isomorphic if there exist matrices $X \in \mathrm{GL}(U), Y \in \mathrm{GL}(V), Z \in \mathrm{GL}(W)$ such that $(X, Y, Z) \cdot T = T'$, where the latter is shorthand for (1). If we treat T, T' as given non-isomorphic tensors, then we may treat (1) as a system of equations in the $n^2 + m^2 + p^2$ variables $X_{ii'}, Y_{jj'}, Z_{kk'}$. To enforce that these variable matrices are invertible, we furthermore introduce three additional sets of variables X', Y', Z' meant to be the inverse matrices, and include also the equations

$$XX' = X'X = I_n \qquad YY' = Y'Y = I_m \qquad ZZ' = Z'Z = I_p,$$

where I_n denotes the $n \times n$ identity matrix, which is Id_U in any basis. (We could have instead introduced new variables such as δ and the equation $\det(X)\delta = 1$, however, the latter equation is degree n, whereas the above equations all have degree $O(1)$, which is more desirable from the point of view of algebraic proof complexity.)

2.3 Polynomial encodings and the inversion principle

Some principles of linear algebra can be formulated as tautologies in propositional logic and therefore also as a set of polynomial equations. In this paper we preliminarily consider two such principles.

Rank Principle. As a first example we consider a set of unsatisfiable polynomials encoding the principle that the product of a $n \times r$ matrix X by a $r \times n$ matrix Y cannot be the identity matrix whenever $r < n$. We consider variables $x_{i,k}, y_{j,k}$ for $i, j \in [n]$ and $k \in [r]$, where $r < n$ to encode X and Y. Then the polynomial encoding is:

$$\mathbb{I}(r,n) := \sum_{k \in [r]} x_{i,k} y_{j,k} - \delta_{i,j} \qquad i,j \in [n]$$

where $\delta_{i,j} = 1$ if $i = j$ and 0 otherwise. This set of polynomials is clearly unsatisfiable as long as $r < n$.

Inversion Principle. The second principle encodes the invertibility of a square $n \times n$ matrix A, expressing the tautology that $AB = I \to BA = I$ where A, B are $n \times n$ matrices and I is the identity matrix. Stephen A. Cook suggested this principle as a tautology that may be hard to prove in several proof systems.

Let $a_{i,j}, b_{i,j}$ be formal variables encoding respectively the (i,j)-th entries of A and B. We represent the fact that $AB = I$ as the set of degree 2 polynomials

$$\sum_{k \in [n]} a_{i,k} b_{k,j} - \delta_{i,j} \qquad i,j \in [n],$$

where $\delta_{i,j} = 1$ if $i = j$ and 0 otherwise. We denote this set of polynomials by $AB = I$. In Section 3 (in the original article), we study the degree complexity of $AB = I \vdash BA = I$, that is of PC derivations of the polynomials $BA = I$ from the polynomials $AB = I$.

In view of the results we obtain in Section 3 (in the original article), in Section 1.2 we considered a *polynomial rule schema* of the form

$$\frac{AB = I}{BA = I}$$

which we call the *Inversion Rule* (INV) meant to be added to PC as an extra rule. We make this slightly more precise here.

A polynomial instantiation τ of the polynomials $AB = I$ is a substitution of polynomials $p_{i,j}, q_{i,j}$ to variables $a_{i,j}$ and $b_{i,j}$. In PC+INV a polynomial p is derivable from a set of polynomials $\mathcal{P}$ if

1. p is an axiom, or $p \in \mathcal{P}$;
2. p is obtained by multiplication or linear combination from previous polynomials in the proof;
3. p is a polynomial among a polynomial instantiation τ of $BA = I$, given that among the polynomials previously derived in the proof there are all the polynomials forming the instantiation τ of $AB = I$.

Pigeonhole Principle. An important role in proving the results in Section 3 (in the original article) is played by the well-known *Pigeonhole principle* stating that any function f from $[n]$ to $[r]$ with $r < n$ has a collision, that is there are $i \neq i' \in [n]$ and a $j \in [r]$ such that $f(i) = f(i') = j$. PHP_r^n is the set of polynomials:

$$\sum_{k\in[r]} p_{i,k}-1, \text{ for } i\in[n], \qquad p_{i,k}p_{j,k}, \text{ for } i\neq j\in[n], k\in[r] \qquad p_{ij}^2-p_{ij}, \text{ for } i\in[n], j\in[r]$$

Razborov [Raz98] additionally included the "functional equations" (encoding that each pigeon cannot be matched to more than one hole):

$$p_{i,k}p_{i,k'}, \text{ for } i\in[n], k\neq k'\in[r].$$

4 Upper bound for non-isomorphism of bounded-rank tensors

Theorem 4.1. *Over any algebraically closed field, there is a function $f(r) \leq 2^{O(r^2)}$, depending only on r, such that, given two non-isomorphic tensors M, M' of tensor rank $\leq r$, the Nullstellensatz degree of refuting isomorphism is at most $f(r)$.*

If working over a finite field $GF(q)$ and including the equations $x^q - x = 0$ for all variables x, then the PC degree is at most $12qr^2$.

Proof. The proof is based mainly on the so-called inheritance property of tensor rank.

Let $M = \sum_{i=1}^r u_i \otimes v_i \otimes w_i$ and let $M' = \sum_{i=1}^r u'_i \otimes v'_i \otimes w'_i$ be our two tensors of format $n_1 \times n_2 \times n_3$. Let $d_1 = \dim \mathrm{Span}\{u_1, u_2, \ldots, u_r, u'_1, u'_2, \ldots, u'_r\}$, d_2 similarly for the v's and d_3 for the w's. Choose a basis $e_1, e_2, \ldots, e_{n_1}$ for $\mathbb{F}^{n_1}$ such that $\mathrm{Span}\{e_1, \ldots, e_{d_1}\} = \mathrm{Span}\{u_1, \ldots, u_r, u'_1, \ldots, u'_r\}$. Let $f_1, \ldots, f_{n_2}$ be a similar basis for $\mathbb{F}^{n_2}$ (with the first d_2 vectors a basis for $\mathrm{Span}\{v_1, \ldots, v_r, v'_1, \ldots, v'_r\}$), and similarly $g_1, \ldots, g_{n_3}$. Changing everything in sight into the $e_\bullet \otimes f_\bullet \otimes g_\bullet$ basis, we find that M, M' are both supported in the upper-left $d_1 \times d_2 \times d_3$ sub-tensors, with all zeros outside of this. Call the corresponding $d_1 \times d_2 \times d_3$ tensors $\overline{M}, \overline{M}'$. Because all the entries outside this box are zero, it is not difficult to show that $M \cong M'$ iff $\overline{M} \cong \overline{M}'$ (the so-called "Inheritance Theorem,", see, e. g., [Lan12, §3.7.1]); note that isomorphism of $\overline{M}$ with $\overline{M}'$ is via the much smaller group $\mathrm{GL}_{d_1} \times \mathrm{GL}_{d_2} \times \mathrm{GL}_{d_3}$, rather than $\mathrm{GL}_n \times \mathrm{GL}_n \times \mathrm{GL}_n$ (the latter of which is used to determine isomorphism of M with M').

In this basis, isomorphism of $\overline{M}, \overline{M}'$ is solely determined by the upper-left $d_1 \times d_1$ sub-matrix of X, X', the upper-left $d_2 \times d_2$ submatrix of Y, Y', and the upper-left $d_3 \times d_3$ sub-matrix of Z, Z'. So we now only need to deal with equations in $d_1^2 + d_2^2 + d_3^2$ variables. Since each $d_i \leq 2r$, this is at most $12r^2$ variables.

Since we have $\leq 12r^2$ variables, $d_1 d_2 d_3$ cubic equations, and $6n^2$ quadratic equations ($XX' = I = X'X = YY' = \cdots$), over an algebraically closed field Sombra's Effective Nullstellensatz [Som99] implies that the Nullstellensatz degree of refuting our equations is then at most $4 \cdot 3^{\Theta(r^2)}$.

Over a finite field with the extra equations $x^q = x$, we may reduce degrees so that the degree of each variable is never more than q, the size of the field. In this case, the PC degree is at most q times the number of variables, i. e., at most $12qr^2$. □

References

[ABRW04] Michael Alekhnovich, Eli Ben-Sasson, Alexander A. Razborov, and Avi Wigderson. Pseudorandom generators in propositional proof complexity. *SIAM J. Comput.*, 34(1):67–88, 2004. doi:10.1137/S0097539701389944.

[AM13] Albert Atserias and Elitza N. Maneva. Sherali–Adams relaxations and indistinguishability in counting logics. *SIAM J. Comput.*, 42(1):112–137, 2013. doi:10.1137/120867834.

[AS05] Manindra Agrawal and Nitin Saxena. Automorphisms of finite rings and applications to complexity of problems. In *STACS 2005, 22nd Annual Symposium on Theoretical Aspects of Computer Science, Proceedings*, pages 1–17, 2005. doi:10.1007/978-3-540-31856-9_1.

[Bab16] László Babai. Graph isomorphism in quasipolynomial time [extended abstract]. In *STOC'16—Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing*, pages 684–697. ACM, New York, 2016. doi:10.1145/2897518.2897542.

[BCIP02] Josh Buresh-Oppenheim, Matthew Clegg, Russell Impagliazzo, and Toniann Pitassi. Homogenization and the polynomial calculus. *Comput. Complex.*, 11(3-4):91–108, 2002. doi:10.1007/s00037-002-0171-6.

[Ber18] Christoph Berkholz. The relation between polynomial calculus, sherali-adams, and sum-of-squares proofs. In *35th Symposium on Theoretical Aspects of Computer Science (STACS 2018)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2018.

[BG15] Christoph Berkholz and Martin Grohe. Limitations of algebraic approaches to graph isomorphism testing. In Magnús M. Halldórsson, Kazuo Iwama, Naoki Kobayashi, and Bettina Speckmann, editors, *Automata, Languages, and Programming - 42nd International Colloquium, ICALP 2015, Kyoto, Japan, July 6-10, 2015, Proceedings, Part I*, volume 9134 of *Lecture Notes in Computer Science*, pages 155–166. Springer, 2015.

[BG17] Christoph Berkholz and Martin Grohe. Linear diophantine equations, group csps, and graph isomorphism. In Philip N. Klein, editor, *Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2017, Barcelona, Spain, Hotel Porta Fira, January 16-19*, pages 327–339. SIAM, 2017. Preprint arXiv:1607.04287 [cs.CC]. doi:10.1137/1.9781611974782.21.

[BGIP01] Samuel R. Buss, Dima Grigoriev, Russell Impagliazzo, and Toniann Pitassi. Linear gaps between degrees for the polynomial calculus modulo distinct primes. *J. Comput. Syst. Sci.*, 62(2):267–289, 2001. doi:10.1006/jcss.2000.1726.

[BGL+19] Peter A. Brooksbank, Joshua A. Grochow, Yinan Li, Youming Qiao, and James B. Wilson. Incorporating Weisfeiler–Leman into algorithms for group isomorphism. arXiv:1905.02518 [cs.CC], 2019.

[BI99] Eli Ben-Sasson and Russell Impagliazzo. Random CNF's are hard for the Polynomial Calculus. In *40th Annual Symposium on Foundations of Computer Science, FOCS '99, 17-18 October, 1999, New York, NY, USA*, pages 415–421. IEEE Computer Society, 1999. (Journal version in *Comput. Complex.* 2010, doi:10.1007/s00037-010-0293-1). doi:10.1109/SFFCS.1999.814613.

[Bre70] R. P. Brent. Algorithms for matrix multiplication. Stanford Computer Science Dept. Tech. Report STAN-CS-70-157, available online at https://apps.dtic.mil/sti/pdfs/AD0705509.pdf, 1970.

[BS20] Jendrik Brachter and Pascal Schweitzer. On the Weisfeiler–Leman dimension of finite groups. In Holger Hermanns, Lijun Zhang, Naoki Kobayashi, and Dale Miller, editors, *LICS '20: 35th Annual ACM/IEEE Symposium on Logic in Computer Science, Saarbrücken, Germany, July 8-11, 2020*, pages 287–300. ACM, 2020. doi:10.1145/3373718.3394786.

[BS22] Jendrik Brachter and Pascal Schweitzer. A systematic study of isomorphism invariants of finite groups via the Weisfeiler–Leman dimension. In Shiri Chechik, Gonzalo Navarro, Eva Rotenberg, and Grzegorz Herman, editors, *30th Annual European Symposium on Algorithms, ESA 2022, September 5-9, 2022, Berlin/Potsdam, Germany*, volume 244 of *LIPIcs*, pages 27:1–27:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022. doi:10.4230/LIPIcs.ESA.2022.27.

[CEI96] Matthew Clegg, Jeffery Edmonds, and Russell Impagliazzo. Using the Groebner basis algorithm to find proofs of unsatisfiability. In *Proceedings of the Twenty-eighth Annual ACM Symposium on the Theory of Computing (Philadelphia, PA, 1996)*, pages 174–183. ACM, New York, 1996. doi:10.1145/237814.237860.

[CFI92] Jin-Yi Cai, Martin Fürer, and Neil Immerman. An optimal lower bound on the number of variables for graph identification. *Combinatorica*, 12(4):389–410, 1992. doi:10.1007/BF01305232.

[CL22] Nathaniel A. Collins and Michael Levet. Count-free Weisfeiler–Leman and group isomorphism. arXiv:2212.11247 [cs.DS], 2022.

[FGS19] Vyacheslav Futorny, Joshua A. Grochow, and Vladimir V. Sergeichuk. Wildness for tensors. *Linear Algebra Appl.*, 566:212–244, 2019. doi:10.1016/j.laa.2018.12.022.

[FP06] Jean-Charles Faugère and Ludovic Perret. Polynomial equivalence problems: Algorithmic and theoretical aspects. In Serge Vaudenay, editor, *Advances in Cryptology - EUROCRYPT 2006, 25th Annual International Conference on the Theory and Applications of Cryptographic Techniques, St. Petersburg, Russia, May 28 - June*

1, 2006, Proceedings, volume 4004 of *Lecture Notes in Computer Science*, pages 30–47. Springer, 2006. doi:10.1007/11761679_3.

[GQ21a] Joshua A. Grochow and Youming Qiao. On p-group isomorphism: Search-to-decision, counting-to-decision, and nilpotency class reductions via tensors. In Valentine Kabanets, editor, *36th Computational Complexity Conference, CCC 2021, July 20-23, 2021, Toronto, Ontario, Canada (Virtual Conference)*, volume 200 of *LIPIcs*, pages 16:1–16:38. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021. doi:10.4230/LIPIcs.CCC.2021.16.

[GQ21b] Joshua A. Grochow and Youming Qiao. On the complexity of isomorphism problems for tensors, groups, and polynomials I: tensor isomorphism-completeness. In James R. Lee, editor, *12th Innovations in Theoretical Computer Science Conference, ITCS 2021, January 6-8, 2021, Virtual Conference*, volume 185 of *LIPIcs*, pages 31:1–31:19. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021. doi:10.4230/LIPIcs.ITCS.2021.31.

[Gri81] D. Ju. Grigoriev. Complexity of "wild" matrix problems and of the isomorphism of algebras and graphs. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 105:10–17, 198, 1981. Theoretical applications of the methods of mathematical logic, III. doi:10.1007/BF01084390.

[Gri01] Dima Grigoriev. Linear lower bound on degrees of positivstellensatz calculus proofs for the parity. *Theoretical Computer Science*, 259(1-2):613–622, 2001.

[Gri13] Dima Grigoriev. Polynomial complexity of solving systems of few algebraic equations with small degrees. In Vladimir P. Gerdt, Wolfram Koepf, Ernst W. Mayr, and Evgenii V. Vorozhtsov, editors, *Computer Algebra in Scientific Computing - 15th International Workshop, CASC 2013, Berlin, Germany, September 9-13, 2013. Proceedings*, volume 8136 of *Lecture Notes in Computer Science*, pages 136–139. Springer, 2013. doi:10.1007/978-3-319-02297-0_11.

[Gro12] Joshua A. Grochow. Matrix Lie algebra isomorphism. In *IEEE Conference on Computational Complexity (CCC12)*, pages 203–213, 2012. Also available as arXiv:1112.2012 [cs.CC] and ECCC Technical Report TR11-168. doi:10.1109/CCC.2012.34.

[Gro13] Joshua A. Grochow. Answer to "deciding bound on tensor rank for a fixed value". CSTheory StackExchange, https://cstheory.stackexchange.com/a/19518/129, 2013.

[Hel19] Harald Andrés Helfgott. Isomorphismes de graphes en temps quasi-polynomial [d'après Babai et Luks, Weisfeiler-Leman,...]. *Astérisque*, (407):Exp. No. 1125, 135–182, 2019. Séminaire Bourbaki. Vol. 2016/2017. Exposés 1120–1135. English translation with appendices by Jitendra Bajpai and Daniele Dona available at arXiv:17010.04574 [math.GR]. doi:10.24033/ast.

[HQ21] Xiaoyu He and Youming Qiao. On the Baer-Lovász-Tutte construction of groups from graphs: isomorphism types and homomorphism notions. *European J. Combin.*, 98:Paper No. 103404, 12, 2021. doi:10.1016/j.ejc.2021.103404.

[HT15] Pavel Hrubes and Iddo Tzameret. Short proofs for the determinant identities. *SIAM J. Comput.*, 44(2):340–383, 2015. doi:10.1137/130917788.

[JQSY19] Zhengfeng Ji, Youming Qiao, Fang Song, and Aaram Yun. General linear group action on tensors: A candidate for post-quantum cryptography. In Dennis Hofheinz and Alon Rosen, editors, *Theory of Cryptography - 17th International Conference, TCC 2019, Nuremberg, Germany, December 1-5, 2019, Proceedings, Part I*, volume 11891 of *Lecture Notes in Computer Science*, pages 251–281. Springer, 2019. Preprint arXiv:1906.04330 [cs.CR]. doi:10.1007/978-3-030-36030-6_11.

[Lan12] J. M. Landsberg. *Tensors: geometry and applications*, volume 128 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012. doi:10.1090/gsm/128.

[Las01] Jean B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM J. Optim.*, 11(3):796–817, 2000/01. doi:10.1137/S1052623400366802.

[Lic85] Thomas Lickteig. Typical tensorial rank. *Linear Algebra Appl.*, 69:95–120, 1985. doi:10.1016/0024-3795(85)90070-9.

[LL89] Thomas Lehmkuhl and Thomas Lickteig. On the order of approximation in approximative triadic decompositions of tensors. *Theoret. Comput. Sci.*, 66(1):1–14, 1989. doi:10.1016/0304-3975(89)90141-2.

[Luk93] Eugene M. Luks. Permutation groups and polynomial-time computation. In *Groups and computation (New Brunswick, NJ, 1991)*, volume 11 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 139–175. Amer. Math. Soc., Providence, RI, 1993.

[McK81] Brendan D. McKay. Practical graph isomorphism. *Congr. Numer.*, 30:45–87, 1981.

[Miy96] Takunari Miyazaki. Luks's reduction of graph isomorphism to code equivalence. Comment to E. W. Clark, https://groups.google.com/forum/#!msg/sci.math.research/puZxGj9HXKI/CeyH2yyyNFUJ, 1996.

[MP14] Brendan D. McKay and Adolfo Piperno. Practical graph isomorphism, II. *J. Symbolic Comput.*, 60:94–112, 2014. doi:10.1016/j.jsc.2013.09.003.

[NPOV15] Alexander Novikov, Dmitry Podoprikhin, Anton Osokin, and Dmitry Vetrov. Tensorizing neural networks. In *Proceedings of the 28th International Conference on Neural Information Processing Systems - Volume 1*, NIPS'15, pages 442–450. MIT Press, 2015.

[OWWZ14] Ryan O'Donnell, John Wright, Chenggang Wu, and Yuan Zhou. Hardness of robust graph isomorphism, lasserre gaps, and asymmetry of random graphs. In Chandra Chekuri, editor, *Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2014, Portland, Oregon, USA, January 5-7, 2014*, pages 1659–1677. SIAM, 2014. Preprint available as arXiv:1401.2436 [cs.CC]. doi:10.1137/1.9781611973402.120.

[Pat96] Jacques Patarin. Hidden fields equations (HFE) and isomorphisms of polynomials (IP): two new families of asymmetric algorithms. In *Advances in Cryptology - EUROCRYPT '96, International Conference on the Theory and Application of Cryptographic Techniques, Saragossa, Spain, May 12-16, 1996, Proceeding*, pages 33–48, 1996. doi:10.1007/3-540-68339-9_4.

[PR97] Erez Petrank and Ron M. Roth. Is code equivalence easy to decide? *IEEE Trans. Inf. Theory*, 43(5):1602–1604, 1997. doi:10.1109/18.623157.

[Raz98] Alexander A. Razborov. Lower bounds for the polynomial calculus. *Comput. Complex.*, 7(4):291–324, 1998. doi:10.1007/s000370050013.

[SC04] Michael Soltys and Stephen Cook. The proof complexity of linear algebra. *Ann. Pure Appl. Logic*, 130(1-3):277–323, 2004. doi:10.1016/j.apal.2003.10.018.

[Sch08] Grant Schoenebeck. Linear level lasserre lower bounds for certain k-csps. In *2008 49th Annual IEEE Symposium on Foundations of Computer Science*, pages 593–602. IEEE, 2008.

[Sol01] Michael Soltys. *The complexity of derivations of matrix identities.* PhD thesis, University of Toronto, 2001. Availalble on ECCC at https://eccc.weizmann.ac.il/resources/pdf/soltys.pdf.

[Som99] Martín Sombra. A sparse effective Nullstellensatz. *Adv. in Appl. Math.*, 22(2):271–295, 1999. doi:10.1006/aama.1998.0633.

[SSC14] Aaron Snook, Grant Schoenebeck, and Paolo Codenotti. Graph Isomorphism and the Lasserre hierarchy. arXiv:1401.0758 [cs.CC], 2014.

[TDJ+22] Gang Tang, Dung Hoang Duong, Antoine Joux, Thomas Plantard, Youming Qiao, and Willy Susilo. Practical post-quantum signature schemes from isomorphism problems of trilinear forms. In Orr Dunkelman and Stefan Dziembowski, editors, *Advances in Cryptology - EUROCRYPT 2022 - 41st Annual International Conference on the Theory and Applications of Cryptographic Techniques, Trondheim, Norway, May 30 - June 3, 2022, Proceedings, Part III*, volume 13277 of *Lecture Notes in Computer Science*, pages 582–612. Springer, 2022. doi:10.1007/978-3-031-07082-2_21.

[Zui17] Jeroen Zuiddam. A note on the gap between rank and border rank. *Linear Algebra Appl.*, 525:33–44, 2017. doi:10.1016/j.laa.2017.03.015.