Stanley Supplementary Exercise 88
The power-sum image of the hook-content transformation

Statement: Richard P. Stanley
Paired original proof: Richard P. Stanley

Statement version 4 June 2023; author proof arXiv:0807.0383v3

Original exercise

Define a $\mathbb{Q}$-linear transformation $\varphi : \Lambda_{\mathbb{Q}} \to \mathbb{Q}[t]$ by

$$\varphi(s_\lambda) = \frac{\prod_{i=1}^{n}(t+\lambda_i+n-i)}{H_\lambda},$$

where $\lambda = (\lambda_1, \ldots, \lambda_n) \vdash n$ and H_λ denotes the product of the hook lengths of λ. Show that for any $\mu \vdash n$ with $\ell(\mu) = \ell$ and $m_1(\mu) = m$ (the number of parts of μ equal to 1), we have

$$\varphi(p_\mu) = (-1)^{n-\ell} \sum_{i=0}^{m} \binom{m}{i} t(t+1)\cdots(t+i-1).$$

Paired author proof

The official solutions explicitly cite Lemma 3.2 of R. Stanley, *Some combinatorial properties of hook lengths, contents, and parts of partitions*, Ramanujan J. 23 (2010), 91–105; arXiv:0807.0383v3. The lemma as printed there is exactly the displayed identity, with the paper's variable v in place of t (the exercise writes $\varphi : \Lambda_{\mathbb{Q}} \to \mathbb{Q}[t]$, the paper writes $\varphi : R \to \mathbb{Q}[v_1, \ldots, v_k]$ and sets $v = v_1$). The complete native LaTeX source and the author PDF of version 3 are retained with this statement. The paired source uses the paper's definition of φ (Section 3) and the full proof of Lemma 3.2: expanding e_{n-i} in power sums gives the coefficient of p_μ in equation (16), and applying φ to the Cauchy identity (17) followed by linear independence of the p_μ closes the argument. Lemma 3.1 supplies the expansion used at equation (11).

Sources

Original statement: Stanley, exercise 88, physical page 27, `https://math.mit.edu/~rstan/ec/ch7supp.pdf`. Official solution pointer: version 24 March 2023, physical page 40, `https://math.mit.edu/~rstan/ec/ch7suppsol.pdf`. Original author proof: `https://arxiv.org/abs/0807.0383v3`.