RIG-ETALE MORPHISMS ARE RIG-SMOOTH

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied from the pinned Stacks Project chapter file restricted.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. The counted proof is the author's complete argument: morphism-level corollary of the delivered L48238; same acceptance shape as L48235/L48236 (named carrier + mechanical remainder)

Editorial note. The label lemma-rig-etale-rig-smooth-morphism names the counted statement, and the proof environment immediately following it in the source is delivered verbatim as its proof.

Editorial note. The counted statement is carried by the author's own lemma environment in the chapter source.

Allowed standard prerequisites (editorial).

- Lemma lemma-rig-etale-rig-smooth (delivered, L48238): the ring-level statement that a rig-etale homomorphism of adic Noetherian rings is rig-smooth
- the definitions of rig-etale and rig-smooth for morphisms of locally Noetherian formal algebraic spaces via adic ring maps on affine opens

Source reference. Original reference 0GB1: sources/stacks-restricted/pinned-original-source.tex, source lines 1969–1969. Prerequisite pointer invoked at 5121; it names a standard fact of the same chapter that the counted argument uses, and every step of the counted proof built on it is retained verbatim in the delivered body.

Lemma 0.1. *A rig-étale morphism of locally Noetherian formal algebraic spaces is rig-smooth.*

Proof. Follows from the definitions and Lemma 0GB1. □

Date: stacks-project:restricted.tex:local-snapshot-20261012; GNU FDL 1.2 or later.