SEMINORMALITY AND ABSOLUTE WEAK NORMALITY ARE LOCAL PROPERTIES OF RINGS

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied from the pinned Stacks Project chapter file morphisms.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. The chapter file pinned-original-source.tex carries only \input{preamble} on its first line, so no preamble text lives inside the chapter. The retained file sources/stacks-morphisms/retained-preamble.tex is a mechanical, line-for-line copy of the project's single shared preamble as staged in the snapshot. Nothing was added, removed or redefined, and the excerpt serves only as preamble, never as a span coordinate.

Editorial note. Both counted spans start at the \begin line of their environment, so the statement and the author's proof are delivered whole; surrounding prose of the section is deliberately outside the counted spans and is carried as standard prerequisites instead.

Editorial note. The citation fields (source_id, source_version, authors, source_url, license.name, source_title) are identical to those of every other card this lane has produced from this chapter, so the batch selector treats them as one source.

Editorial note. Screened under the Lead's 12:4x rule (prescreen hit, proof body 1136 bytes): read in full and accepted, not refused. The seminar-normal half is proved out in detail; the absolutely weakly normal half is dispatched by the author's own words 'In exactly the same manner the reader shows that if A is absolutely weakly normal, then A_f is absolutely weakly normal' and 'then the exact same argument shows that A is absolutely weakly normal' – a self-description of a routine repetition of the two arguments given just above, which is the class the Lead's rule says to accept with the wording quoted. It is deliberately NOT in the class of the three refusals in this same window (lemma-seminormalization-ring, lemma-seminormalization, lemma-rational-map-from-reduced-to-separated), where a numbered statement item is left with no argument whatsoever ('we omit the proof of (3) and (4)'). Both verbatim phrases are quoted here so that a reviewer can re-judge the call without re-reading the source.

Allowed standard prerequisites (editorial).

- The definition of a local property of rings, in terms of localisations at a finite family generating the unit ideal. Properties, Definition \ref{properties-definition-property-local}, cited at proof lines 12023-12026. Fixes what has to be proved for the converse direction.
- Glueing of elements given over a standard covering of the spectrum. Algebra, Lemma \ref{algebra-lemma-standard-covering}, cited at proof lines 12047-12050. Produces the single element inducing all the local ones.
- The definition of seminormal and absolutely weakly normal rings. Definition \ref{definition-seminormal-ring}, the definition that this lemma is

Date: stacks-project:morphisms.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

about. Supplies the equations and the uniqueness clause used in both directions.

Source reference. Original lemma 0EUM: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 18557–18557. The lemma's own canonical tag, quoted so that readers can resolve the label inside the delivered text; it points at the counted statement itself and carries no proof content.

Source reference. Original lemma 01OP: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 2202–2202. Navigation pointer to a statement the counted proof refers to but does not contain; nothing load-bearing is delegated to it.

Source reference. Original lemma 00EJ: focused-seeds/AGP/sources/stacks-snapshot-20261011/tags.txt, source lines 536–536. Navigation pointer to a statement the counted proof refers to but does not contain; nothing load-bearing is delegated to it.

Lemma 0.2. *Being seminormal or being absolutely weakly normal is a local property of rings, see Properties, Definition 01OP.*

Proof. Suppose that A is seminormal and $f \in A$. Let $x', y' \in A_f$ with $(x')^3 = (y')^2$. Write $x' = x/f^{2n}$ and $y' = y/f^{3n}$ for some $n \geq 0$ and $x, y \in A$. After replacing x, y by $f^{2m}x, f^{3m}y$ and n by $n + m$, we see that $x^3 = y^2$ in A. Then we find a unique $a \in A$ with $x = a^2$ and $y = a^3$. Setting $a' = a/f^n$ we get $x' = (a')^2$ and $y' = (a')^3$ as desired. Uniqueness of a' follows from uniqueness of a. In exactly the same manner the reader shows that if A is absolutely weakly normal, then A_f is absolutely weakly normal.

Assume A is a ring and $f_1, \ldots, f_n \in A$ generate the unit ideal. Assume A_{f_i} is seminormal for each i. Let $x, y \in A$ with $x^3 = y^2$. For each i we find a unique $a_i \in A_{f_i}$ with $x = a_i^2$ and $y = a_i^3$ in A_{f_i}. By the uniqueness and the result of the first paragraph (which tells us that $A_{f_i f_j}$ is seminormal) we see that a_i and a_j map to the same element of $A_{f_i f_j}$. By Algebra, Lemma 00EJ we find a unique $a \in A$ mapping to a_i in A_{f_i} for all i. Then $x = a^2$ and $y = a^3$ by the same token. Clearly this a is unique. Thus A is seminormal. If we assume A_{f_i} is absolutely weakly normal, then the exact same argument shows that A is absolutely weakly normal. □