Uniform semiclassical nonlinear Hartree phase mixing and weighted scattering for small perturbations

Marnie Smith

doi:10.5802/jep.353; CC BY4.0

Editorial note. Only the full nonlinear main theorem is counted. Bootstrap improvements, Green-function bounds, resonance integrals and the scattering deduction are solution components, not separate new tasks.

Editorial note. Local well-posedness/persistence and its continuation criterion are explicitly permitted standard inputs, following the authors declaration at761 and860. The uniform nonlinear a-priori closure is fully supplied.

Editorial note. The Penrose boundary criterion and sufficient radial/smallness cases are outside this task; uniform Penrose stability is an assumption.

Editorial note. The original bootstrap dependency diagram is native TikZ in the selected proof and needs no external raster asset. Bibliography keys are mechanically mapped to the publishers exact references; the BMM and Nguyen-You attribution is retained.

Editorial note. 2026-10-11 材料收口：复用既有 H2、来源许可和近邻判断；核对原命题与作者证明起止、所指上下文及装配范围齐备。未重证作者结论；沿用既有非重复依据。

Editorial scope. One author theorem: global nonlinear phase mixing and scattering with constants independent of hbar in(0,1]. All load-bearing source-local analytical estimates are in the solution; no bootstrap bound is assumed for free.

Allowed standard prerequisites (editorial).

- Classical Fourier and Sobolev analysis. Plancherel, Fourier/Laplace inversion, Sobolev embedding and trace on a line, weighted-moment equivalence, Young and Schur estimates, interpolation, and elementary commutator/Leibniz formulas.

- Local Hartree well-posedness and continuation. For each fixed hbar in(0,1], the weighted quantum Sobolev data have a unique maximal self-adjoint solution with persistence of regularity; bounded lower Sobolev control on each finite interval gives continuation. This is the standard local input explicitly invoked by the source at761 and860, not a claimed new local well-posedness proof.

0.1 The Hartree equation

The *Hartree equation*, a mean-field model describing the dynamics of infinitely many interacting fermions, is expressed as:

$$\begin{cases} i\hbar\partial_t\gamma = \Big[-\dfrac{\hbar^2}{2}\Delta + w * \rho_\gamma, \gamma\Big], \\ \gamma(0) = \gamma_{\text{in}}, \end{cases} \tag{1}$$

where $\gamma(t)$ is a self-adjoint operator on $L^2(\mathbb{R}^d;\mathbb{C})$. The function $\rho_\gamma(t,x) = \gamma(t,x,x)$ represents the density, with $\gamma(t,x,y)$ denoting the integral kernel of $\gamma(t)$. The nonlinear convolution term $w*\rho_\gamma$ describes the mean-field interaction between the particles, where $w\colon \mathbb{R}^d \to \mathbb{R}$ is a specified two-body interaction kernel. The parameter $\hbar > 0$ represents the Planck constant, and the semiclassical limit involves taking $\hbar \to 0$.

For any nonnegative function $g \in L^\infty(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$, define the translation-invariant operator

$$\gamma_g := (2\pi\hbar)^d g(-i\hbar\nabla)$$

on $L^2(\mathbb{R}^d;\mathbb{C})$. The integral kernel of γ_g is

$$\gamma_g(x,y) = \widehat{g}\Big(\frac{y-x}{\hbar}\Big),$$

and its constant density is given by

$$\rho_{\gamma_g} = \widehat{g}(0) = \int_{\mathbb{R}^d} g(k)\,dk.$$

These operators represent a class of steady-state solutions to the Hartree equation (1), as they satisfy the commutation relations $[-\Delta, \gamma_g] = 0$ and $[w * \rho_{\gamma_g}, \gamma_g] = 0$, where $w * \rho_{\gamma_g}$ is a constant multiple of the identity operator. The trace of a solution to (1) determines the total number of particles. For the steady states γ_g with $g \not\equiv 0$, this number is infinite due to the constant positive density: $\operatorname{Tr}\gamma_g = \int_{\mathbb{R}^d} \rho_{\gamma_g}\,dx = +\infty$.

The interaction kernel and steady state are assumed to satisfy a linear stability criterion, known in kinetic theory as the *Penrose condition*. In the

present setting, the condition must hold uniformly for all $\hbar \in (0,1]$ in order to remain valid in the semiclassical limit $\hbar \to 0$. This motivates the following formulation:

Definition 0.1 (Uniform Penrose condition)**.** *An interaction kernel $w\colon \mathbb{R}^d \to \mathbb{R}$ and steady state $\gamma_g = (2\pi\hbar)^d g(-i\hbar\nabla)$ satisfy the uniform Penrose condition if there exists a constant $c_{\mathrm{P}} > 0$ such that the inequality*

$$\inf_{\substack{\operatorname{Re}\lambda \geq 0 \\ k \in \mathbb{R}^d}} \left| 1 + \widetilde{\mathscr{L}}(\lambda, k) \right| \geq c_{\mathrm{P}} \tag{2}$$

holds uniformly for all $\hbar \in (0,1]$, where

$$\widetilde{\mathscr{L}}(\lambda, k) = \frac{2}{\hbar}\, \widehat{w}(k) \int_0^\infty e^{-\lambda t} \sin\!\left(\tfrac{1}{2}\hbar t |k|^2\right) \widehat{g}\,(kt)\; dt.$$

Define the perturbation $Q := \gamma - \gamma_g$, where γ is a solution of (1) and $\gamma_g = (2\pi\hbar)^d g(-i\hbar\nabla)$ is a translation-invariant steady state. The evolution of Q is governed by the nonlinear Hartree equation

$$\begin{cases} i\hbar\partial_t Q = \left[-\dfrac{\hbar^2}{2}\Delta + w * \rho_Q, Q + \gamma_g\right], \\ Q(0) = Q_{\mathrm{in}}. \end{cases} \tag{3}$$

0.2 Quantum operators and norms

To formulate phase-mixing and scattering estimates in the quantum setting, quantum analogues of classical derivatives and weighted Sobolev norms are introduced. The standard and double Japanese brackets are defined by $\langle x \rangle := (1 + |x|^2)^{1/2}$ and $\langle x, y \rangle := (1 + |x|^2 + |y|^2)^{1/2}$, respectively. For a multi-index $\alpha = (\alpha_1, \ldots, \alpha_d) \in \mathbb{N}^d$, write $|\alpha| = \alpha_1 + \cdots + \alpha_d$ and $D^\alpha = \partial_1^{\alpha_1} \cdots \partial_d^{\alpha_d}$.

Definition 0.2 (Quantum derivatives and weights)**.** *The quantum analogues of spatial and velocity derivatives are*

$$\begin{aligned} \boldsymbol{\nabla}_x Q &:= [\nabla, Q], \quad & (\boldsymbol{\nabla}_x Q)(x,y) &= (\nabla_x + \nabla_y) Q(x,y), \\ \boldsymbol{\nabla}_\xi Q &:= [\tfrac{x}{i\hbar}, Q], \quad & (\boldsymbol{\nabla}_\xi Q)(x,y) &= (\tfrac{x-y}{i\hbar}) Q(x,y). \end{aligned}$$

The quantum analogues of spatial and velocity weights are

$$\begin{aligned} \boldsymbol{x} Q &:= \tfrac{1}{2}(xQ + Qx), & (\boldsymbol{x}Q)(x,y) &= (\tfrac{x+y}{2}) Q(x,y), \\ \boldsymbol{\xi} Q &:= -\tfrac{i\hbar}{2}(\nabla Q + Q\nabla), & (\boldsymbol{\xi}Q)(x,y) &= -\tfrac{i\hbar}{2}(\nabla_x - \nabla_y) Q(x,y). \end{aligned}$$

Remark 0.3 (Commutation of the quantum operators). The operators of Definition 0.2 satisfy the commutation relations

$$[\boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi] = [\boldsymbol{x}, \boldsymbol{\xi}] = [\boldsymbol{\nabla}_x, \boldsymbol{\xi}] = [\boldsymbol{x}, \boldsymbol{\nabla}_\xi] = 0,$$

in exact analogy with the classical operators ∇_x, ∇_ξ, x, ξ acting on functions $f(x,\xi)$. Indeed, on the level of integral kernels, $\boldsymbol{x}$ and $\boldsymbol{\nabla}_x$ involve only the sum $x+y$, while $\boldsymbol{\xi}$ and $\boldsymbol{\nabla}_\xi$ involve only the difference $x-y$: for instance,

$$(\nabla_x - \nabla_y)\left[\left(\tfrac{x+y}{2}\right) Q(x,y)\right] = \left(\tfrac{x+y}{2}\right)(\nabla_x - \nabla_y)Q(x,y),$$

which gives $[\boldsymbol{x}, \boldsymbol{\xi}] = 0$, and similarly $(\nabla_x + \nabla_y)(x-y) = 0$ gives $[\boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi] = 0$. The remaining relations hold since $\boldsymbol{\nabla}_x$ and $\boldsymbol{\xi}$ are both constant-coefficient differential operators, and $\boldsymbol{x}$ and $\boldsymbol{\nabla}_\xi$ are both multiplication operators. In particular, products such as $\boldsymbol{\xi}^\alpha \boldsymbol{x}^\beta$ are independent of the order of the factors. By contrast, $[\boldsymbol{\nabla}_x, \boldsymbol{x}] \neq 0$ and $[\boldsymbol{\nabla}_\xi, \boldsymbol{\xi}] \neq 0$, again as in the classical case.

The classical weighted Sobolev norm $\|\cdot\|_{H^\sigma_M}$ is now recalled.

Definition 0.4 (Weighted Sobolev norm)**.** *Let $\sigma \geq 0$ and let $M \geq 0$ be an integer. For a function $g = g(v)$, the weighted Sobolev norm H^σ_M is defined as*

$$\|g\|_{H^\sigma_M} := \sum_{|\alpha| \leq M} \|v^\alpha \langle \nabla \rangle^\sigma g\|_{L^2}.$$

For a function $f = f(x,v)$, the weighted Sobolev norm H^σ_M is defined as

$$\|f\|_{H^\sigma_M} := \sum_{|\alpha| \leq M} \|v^\alpha \langle \nabla_x, \nabla_v \rangle^\sigma f\|_{L^2},$$

where the weight applies only to the velocity variable.

The quantum Sobolev norms $\mathcal{H}^\sigma_M$ and $\mathcal{H}^\sigma_{N,M}$ are now defined. The former includes weights in $\boldsymbol{\xi}$, while the latter includes weights in both $\boldsymbol{x}$ and $\boldsymbol{\xi}$. The norms depend on $\hbar$ both through a prefactor and through the definitions of the quantum derivative and weight operators $\boldsymbol{\nabla}_\xi$ and $\boldsymbol{\xi}$.

Definition 0.5 (Quantum norms)**.** *For an operator Q on $L^2(\mathbb{R}^d;\mathbb{C})$ with integral kernel $Q = Q(x,y)$, the following norms are defined for $N, M \in \mathbb{N}$:*

a. Quantum Lebesgue norm $\mathcal{L}^2$:

$$\|Q\|_{\mathcal{L}^2} := (2\pi\hbar)^{-d/2} \|Q(x,y)\|_{L^2_{x,y}}.$$

b. $\boldsymbol{\xi}$-weighted quantum Sobolev norm $\mathcal{H}^\sigma_M$:

$$\|Q\|_{\mathcal{H}^\sigma_M} := \sum_{|\alpha| \leq M} \|\boldsymbol{\xi}^\alpha \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi \rangle^\sigma Q\|_{\mathcal{L}^2}.$$

c. $\boldsymbol{x}$–$\boldsymbol{\xi}$-weighted quantum Sobolev norm $\mathcal{H}^\sigma_{N,M}$:

$$\|Q\|_{\mathcal{H}^\sigma_{N,M}} := \sum_{\substack{|\alpha| \leq M \\ |\beta| \leq N}} \left\|\boldsymbol{x}^\beta \boldsymbol{\xi}^\alpha \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi \rangle^\sigma Q\right\|_{\mathcal{L}^2}.$$

0.3 Fourier and Wigner preliminaries

This subsection collects the Fourier conventions and basic properties of the Wigner transform used throughout the paper.

Throughout, integrals such as $\int_x dx$ denote integration over $\mathbb{R}^d$ in the x variable. The following conventions are adopted for the Fourier transform of functions $g = g(v)$ and $f = f(x, v)$:

$$\widehat{g}(p) := \int_v e^{-ip\cdot v} g(v)\, dv, \qquad \widehat{f}(k,p) := \int_x \int_v e^{-ik\cdot x} e^{-ip\cdot v} f(x,v)\, dvdx.$$

On the Fourier side, the operators defined in Definition 0.2 obey the relations:

$$\begin{aligned}
\widehat{\boldsymbol{\nabla}_x Q}(k,p) &= i(k+p)\widehat{Q}(k,p), & \widehat{\boldsymbol{\nabla}_x Q}(\tfrac{k}{2}+p, \tfrac{k}{2}-p) &= ik\widehat{Q}(\tfrac{k}{2}+p, \tfrac{k}{2}-p),\\
\widehat{\boldsymbol{\nabla}_\xi Q}(k,p) &= \tfrac{1}{\hbar}(\nabla_k - \nabla_p)\widehat{Q}(k,p), & \widehat{\boldsymbol{\nabla}_\xi Q}(\tfrac{k}{2}+p, \tfrac{k}{2}-p) &= \tfrac{1}{\hbar}\nabla_p\Big[\widehat{Q}(\tfrac{k}{2}+p, \tfrac{k}{2}-p)\Big],\\
\widehat{\boldsymbol{x} Q}(k,p) &= \tfrac{i}{2}(\nabla_k + \nabla_p)\widehat{Q}(k,p), & \widehat{\boldsymbol{x} Q}(\tfrac{k}{2}+p, \tfrac{k}{2}-p) &= i\nabla_k\Big[\widehat{Q}(\tfrac{k}{2}+p, \tfrac{k}{2}-p)\Big],\\
\widehat{\boldsymbol{\xi} Q}(k,p) &= \tfrac{\hbar}{2}(k-p)\widehat{Q}(k,p), & \widehat{\boldsymbol{\xi} Q}(\tfrac{k}{2}+p, \tfrac{k}{2}-p) &= \hbar p\widehat{Q}(\tfrac{k}{2}+p, \tfrac{k}{2}-p).
\end{aligned}$$

The Wigner transform facilitates the representation of the density matrix in phase space, revealing structural parallels with classical kinetic equations.

Definition 0.6 (Wigner transform)**.** *For an operator Q on $L^2(\mathbb{R}^d;\mathbb{C})$ with integral kernel $Q(x,y)$, define the Wigner transform of Q, $W[Q]\colon \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{C}$, by*

$$W[Q](x,\xi) = \frac{1}{(2\pi)^d}\frac{1}{\hbar^d}\int_{\mathbb{R}^d} e^{-i\xi\cdot y/\hbar} Q\left(x+\tfrac{y}{2}, x-\tfrac{y}{2}\right)\, dy.$$

The Fourier transform of the Wigner transform of Q satisfies

$$\widehat{W[Q]}(k,\eta) = \int_x e^{-ik\cdot x} Q\big(x-\tfrac{\hbar\eta}{2}, x+\tfrac{\hbar\eta}{2}\big)\, dx = \frac{1}{(2\pi)^d}\int_p e^{-i\hbar\eta\cdot p}\widehat{Q}\left(\tfrac{k}{2}+p, \tfrac{k}{2}-p\right)\, dp,$$

and can be controlled by the $\mathcal{H}^0_{M,M}$ norm of Q via the Sobolev embedding.

0.4 Free phase mixing

Consider the linear equation obtained by removing the interaction term from the Hartree equation, referred to here as the *free Schrödinger equation*:

$$\begin{cases} i\hbar\partial_t Q_{\mathrm{FS}} = \Big[-\dfrac{\hbar^2}{2}\Delta, Q_{\mathrm{FS}}\Big], \\ Q_{\mathrm{FS}}(0) = Q_{\mathrm{in}}, \end{cases} \tag{4}$$

where $Q_{\mathrm{FS}}(t)$ is an operator on $L^2(\mathbb{R}^d;\mathbb{C})$. The solution is given explicitly by

$$Q_{\mathrm{FS}}(t) = e^{i(\hbar/2)t\Delta} Q_{\mathrm{in}} e^{-i(\hbar/2)t\Delta}, \tag{5}$$

with integral kernel and Fourier transform

$$Q_{\mathrm{FS}}(t,x,y) = e^{i(\hbar/2)t(\Delta_x-\Delta_y)}Q_{\mathrm{in}}(x,y), \quad \widehat{Q_{\mathrm{FS}}}(t,k,p) = e^{-i(\hbar/2)t(|k|^2-|p|^2)}\widehat{Q_{\mathrm{in}}}(k,p).$$

By Lemma 1.8, the density of Q_{FS} has Fourier transform

$$\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k) = \frac{1}{(2\pi)^d}\int_p e^{-i\hbar p\cdot kt}\widehat{Q_{\mathrm{in}}}\left(\tfrac{k}{2}+p, \tfrac{k}{2}-p\right)\,dp = \widehat{W[Q_{\mathrm{in}}]}(k,kt).$$

The next result gives decay estimates for each Fourier mode of the density under free evolution.

0.5 Bootstrap scheme

This subsection introduces the bootstrap scheme used to prove the nonlinear phase-mixing estimates, based on propagating a set of weighted quantum Sobolev norms.

0.5.1 Bootstrap assumptions

Fix positive constants $\sigma_4 > \sigma_3 > \sigma_2 > \sigma_1$ such that

$$\begin{gathered}\sigma_1 \geq d+8, \qquad \sigma_2-\sigma_1 > 3/2, \qquad \sigma_3-\sigma_2 > d/2,\\ \sigma_4-\sigma_3 > (d+1)/2, \qquad \sigma_4 < N_0-(d+3)/2\end{gathered} \tag{6}$$

and constants $K_i \geq 1$ determined by the proof. Set $0 < \delta \ll 1/2$ such that $\sigma_4-\sigma_3 > (d+1)/2+\delta$. For each fixed $\hbar \in (0,1]$, standard local well-posedness and persistence of regularity provide a unique maximal self-adjoint solution Q to (3) on an interval $[0,T_{\max})$. Fix $T^* \in (0,T_{\max})$ and suppose that the following bootstrap controls hold on $[0,T^*]$:

$$\sum_{|\alpha|\leq M}\left\|\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_4}\langle t\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle\left(\boldsymbol{\xi}^\alpha P(t)\right)\right\|_{\mathcal{L}^2}^2 \leq 4K_1\langle t\rangle^{2d-1}\varepsilon^2, \tag{B1}$$

$$\left\|\langle \hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|_{L_t^2([0,T^*],L_k^2)}^2 \leq 4K_2\varepsilon^2, \tag{B2}$$

$$\sum_{|\alpha|\leq M}\left\|\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_3}|\boldsymbol{\nabla}_x|^\delta\left(\boldsymbol{\xi}^\alpha P(t)\right)\right\|_{\mathcal{L}^2}^2 \leq 4K_3\varepsilon^2, \tag{B3}$$

$$\left\||k|^{1/2}\langle k,kt\rangle^{\sigma_2}\widehat{\rho_Q}(t,k)\right\|_{L_k^\infty L_t^2([0,T^*])}^2 \leq 4K_4\varepsilon^2, \tag{B4}$$

$$\langle k,\eta\rangle^{2\sigma_1}\left|\widehat{W[P]}(t,k,\eta)\right|^2 \leq 4K_5\varepsilon^2. \tag{B5}$$

Remark 0.7. The bootstrap controls (B1), (B3), and (B5) are analogous to those used by Bedrossian, Masmoudi and Mouhot [2], but adapted to the quantum

setting. A key refinement in the present setting is the inclusion of the weight $\langle \hbar k \rangle$ inside the norm in (B2), which has no analogue in the classical framework. Precisely, the additional norm

$$\big\| \hbar k|k|^{1/2} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|^2_{L^2_t([0,T^*],L^2_k)} \leq 4K_2 \varepsilon^2$$

is propagated, reflecting a structural difference between the Hartree and Vlasov equations. While the quantum setting allows control of one more derivative, uniformity in the semiclassical limit requires pairing it with a factor of $\hbar$. This refinement enables the result in Theorem 1.3 to cover the screened Coulomb kernel. As $\hbar \to 0$, the norm in (B2) reduces to the classical counterpart used in [2], and the norm in (B4) already coincides with its classical analogue.

Remark 0.8. The decay statement (17) in Theorem 1.3 follows solely from the control (B5), and the scattering statement (18) follows from (B3) and (B5), as shown in Lemmas 1.14 and 1.15.

0.5.2 Justification of the bootstrap norms

The bootstrap assumptions (B1), (B3), and (B5) are initially satisfied under the smallness condition $\|Q_{\mathrm{in}}\|_{\mathcal{H}^{N_0}_{M,M}} \leq \varepsilon$ in Theorem 1.3. For instance, at $t = 0$, the left-hand side of (B5) satisfies

$$\langle k, \eta \rangle^{2\sigma_1} \Big| \widehat{W[Q_{\mathrm{in}}]}(k,\eta) \Big|^2 = \Big| \widehat{W[\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi \rangle^{\sigma_1} Q_{\mathrm{in}}]}(k,\eta) \Big|^2 \lesssim \|Q_{\mathrm{in}}\|^2_{\mathcal{H}^{\sigma_1}_{M,M}} \leq \varepsilon^2,$$

where Lemma 1.4 has been used. The density estimates (B2) and (B4) are trivially satisfied at $t = 0$, but their long-time structure is less obvious. The next lemma shows that these norms are natural: for the free Schrödinger evolution, they are directly controlled by the initial data Q_{in}. This motivates their appearance in the bootstrap framework.

1 Linearised phase mixing

The *linearised Hartree equation* is obtained by removing the quadratic-in-Q term $[w * \rho_Q, Q]$ from (3), which is expected to be small:

$$\begin{cases} i\hbar \partial_t Q_{\mathrm{L}} = \Big[-\dfrac{\hbar^2}{2} \Delta, Q_{\mathrm{L}} \Big] + [w * \rho_{Q_{\mathrm{L}}}, \gamma_g] \,, \\ Q_{\mathrm{L}}(0) = Q_{\mathrm{in}}. \end{cases} \tag{7}$$

1.1 Fourier–Laplace transform

Writing (7) in kernel form gives

$$i\hbar \partial_t Q_{\mathrm{L}}(t,x,y) = -\frac{\hbar^2}{2} \left(\Delta_x - \Delta_y \right) Q_{\mathrm{L}}(t,x,y) + \left(w * \rho_{Q_{\mathrm{L}}}(t,x) - w * \rho_{Q_{\mathrm{L}}}(t,y) \right) \widehat{g}\Big(\frac{y-x}{\hbar} \Big).$$

By Duhamel's formula, the solution of the linearised Hartree equation can be written as

$$Q_{\mathrm{L}}(t,x,y) = e^{i(\hbar/2)(\Delta_x-\Delta_y)t}Q_{\mathrm{in}}(x,y) \\ -\frac{i}{\hbar}\int_0^t e^{i(\hbar/2)(\Delta_x-\Delta_y)(t-s)}\left[w*\rho_{Q_{\mathrm{L}}}(s,x)-w*\rho_{Q_{\mathrm{L}}}(s,y)\right]\widehat{g}\Big(\frac{y-x}{\hbar}\Big)\,ds. \quad (8)$$

Taking the Fourier transform of (8) yields

$$\widehat{Q_{\mathrm{L}}}(t,k,p) = e^{-i(\hbar/2)(|k|^2-|p|^2)t}\widehat{Q_{\mathrm{in}}}(k,p) \\ -i\hbar^{d-1}(2\pi)^d\widehat{w}(k+p)\left[g(-\hbar p)-g(\hbar k)\right]\int_0^t e^{-i(\hbar/2)(|k|^2-|p|^2)(t-s)}\widehat{\rho_{Q_{\mathrm{L}}}}(s,k+p)\,ds. \quad (9)$$

Using Lemma 1.8, the corresponding expression for the density is

$$\widehat{\rho_{Q_{\mathrm{L}}}}(t,k) = \frac{1}{(2\pi)^d}\int_p e^{-i\hbar kt\cdot p}\widehat{Q_{\mathrm{in}}}\left(\tfrac{k}{2}+p,\tfrac{k}{2}-p\right)\,dp \\ -i\hbar^{d-1}\widehat{w}(k)\int_p\int_0^t e^{-i\hbar(t-s)k\cdot p}\left[g\left(\hbar(p-\tfrac{k}{2})\right)-g\left(\hbar(p+\tfrac{k}{2})\right)\right]\widehat{\rho_{Q_{\mathrm{L}}}}(s,k)\,dsdp.$$

This is a Volterra equation of the form

$$\widehat{\rho_{Q_{\mathrm{L}}}}(t,k) = \widehat{\rho_{Q_{\mathrm{FS}}}}(t,k) - \widehat{\mathscr{L}} *_t \widehat{\rho_{Q_{\mathrm{L}}}}(t,k), \quad (10)$$

where $\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k)$ is the spatial Fourier transform of the density of the free Schrödinger equation (4), and the kernel $\widehat{\mathscr{L}}$ is given by

$$\begin{aligned}\widehat{\mathscr{L}}(t,k) &:= i\hbar^{d-1}\widehat{w}(k)\int_p e^{-i\hbar kt\cdot p}\left[g\left(\hbar(p-\tfrac{k}{2})\right)-g\left(\hbar(p+\tfrac{k}{2})\right)\right]\,dp \\ &= i\hbar^{d-1}\widehat{w}(k)\mathcal{F}_p\left[g\left(\hbar(p-\tfrac{k}{2})\right)-g\left(\hbar(p+\tfrac{k}{2})\right)\right](\hbar kt) \\ &= \frac{i}{\hbar}\widehat{w}(k)\big(e^{-i(\hbar/2)|k|^2t}-e^{i(\hbar/2)|k|^2t}\big)\widehat{g}\,(kt) \\ &= \frac{2}{\hbar}\widehat{w}(k)\sin\big(\tfrac{1}{2}\hbar t|k|^2\big)\widehat{g}\,(kt)\,.\end{aligned}$$

Observe that since $w, g \in L^1(\mathbb{R}^d)$, the zero mode vanishes: $\widehat{\mathscr{L}}(t,k=0)=0$ for all $t \geq 0$.

In order to find an explicit equation for $\widehat{\rho_{Q_{\mathrm{L}}}}$, the Laplace transform in time is applied to (10). Denoting, for $\lambda \in \mathbb{C}$ with $\operatorname{Re}\lambda > 0$, $\widetilde{f}(\lambda,k) := \int_0^\infty e^{-\lambda t}\widehat{f}(t,k)\,dt$, the transformed equation reads

$$\big(1+\widetilde{\mathscr{L}}(\lambda,k)\big)\widetilde{\rho_{Q_{\mathrm{L}}}}(\lambda,k) = \widetilde{\rho_{Q_{\mathrm{FS}}}}(\lambda,k), \quad (11)$$

where

$$\widetilde{\mathscr{L}}(\lambda,k) = \frac{2}{\hbar}\widehat{w}(k)\int_0^\infty e^{-\lambda t}\sin(\tfrac{1}{2}\hbar t|k|^2)\widehat{g}(kt)\,dt. \quad (12)$$

Lemma 1.1 (Ordering of moments)**.** *Let $\sigma \geq 0$, let $M \in \mathbb{N}$, and $g \in H^\sigma_M(\mathbb{R}^d)$. Then the following norms are equivalent:*

$$\|g\|_{H^\sigma_M} := \sum_{|\alpha|\leq M} \|\langle\nabla_k\rangle^\sigma(k^\alpha g)\|_{L^2} \sim \sum_{|\alpha|\leq M} \|k^\alpha\langle\nabla_k\rangle^\sigma g\|_{L^2}.$$

The Sobolev L^2 trace lemma is also recalled.

Lemma 1.2 (L^2 trace [2, Lem. 2.15])**.** *Let $g \in H^s(\mathbb{R}^d)$ with $s > {}^{(d-1)}/_2$ and $C \subset \mathbb{R}^d$ be an arbitrary straight line. Then, there holds*

$$\|g\|_{L^2(C)} \lesssim_{s,d} \|g\|_{H^s(\mathbb{R}^d)}.$$

1.2 Inverse Laplace transform of Green function

Throughout the remainder of the paper, the interaction kernel w and the steady state g are assumed to satisfy the uniform Penrose condition (2) with constant $c_{\mathrm{P}} > 0$. Under this assumption, the formula

$$\widetilde{\rho_{Q_{\mathrm{L}}}}(\lambda,k) = \widetilde{\mathscr{G}}(\lambda,k)\,\widetilde{\rho_{Q_{\mathrm{FS}}}}(\lambda,k) \tag{13}$$

is justified, where

$$\widetilde{\mathscr{G}}(\lambda,k) := \frac{1}{1+\widetilde{\mathscr{L}}(\lambda,k)}. \tag{14}$$

The analysis of the inverse linearised operator, referred to as the *Green function*, follows the approach of Nguyen and You [4]. Specifically, this includes Lemma 1.17, Lemma 1.18, and Proposition 1.19.

For $k \neq 0$, define the *Lindhard* function m_g by

$$\begin{aligned} m_g(\lambda,k) &= \frac{2}{\hbar}\int_0^\infty e^{-\lambda t}\sin(\tfrac{\hbar}{2}t|k|^2)\widehat{g}(kt)\,dt \\ &= \frac{2}{\hbar|k|}\int_0^\infty e^{-\lambda^{s/|k|}}\sin(\tfrac{\hbar}{2}|k|s)\widehat{g}(\tfrac{k}{|k|}s)\,ds, \end{aligned} \tag{15}$$

so that $\widetilde{\mathscr{L}}(\lambda,k) = \widehat{w}(k)m_g(\lambda,k)$. The Lindhard function plays a central role in the physics literature; see [3].

Theorem 1.3 (Nonlinear phase mixing)**.** *Fix dimension $d \geq 3$ and $M := \lceil (d+1)/2 \rceil$. Choose constants $N_0 > \sigma_1 > \sigma_0$ satisfying*

$$\sigma_1 \geq d+8, \quad N_0 - \sigma_1 > 3d/2 + 7/2 \quad \text{and} \quad \sigma_1 - \sigma_0 > d+1.$$

Let $g \in H^{N_0+7/2+\delta_0}_{2+\lceil d/2\rceil}(\mathbb{R}^d;\mathbb{R}_+)$ for some constant $\delta_0 \in (0,1]$, and let $w \in L^1(\mathbb{R}^d,\mathbb{R})$ satisfy

$$|\widehat{w}(k)| \lesssim \frac{1}{\langle k\rangle^{M-1/2}} \tag{16}$$

for all $k \in \mathbb{R}^d$. Assume g and w satisfy the uniform Penrose condition (2) with constant c_{P}. Then there exist constants $C > 0$ and $\varepsilon_0 > 0$, depending only on $N_0, \sigma_1, \sigma_0, w, g, c_{\mathrm{P}}, d, \delta_0$, and independent of $\hbar \in (0,1]$, such that the following holds.

For any $\varepsilon \in (0, \varepsilon_0]$ and any self-adjoint initial data $Q_{\mathrm{in}} \in \mathcal{H}^{N_0}_{M,M}$ with

$$\|Q_{\mathrm{in}}\|_{\mathcal{H}^{N_0}_{M,M}} \leq \varepsilon,$$

the nonlinear Hartree equation (3) admits a unique global self-adjoint solution $Q(t)$, which satisfies:

- *Density decay: for all $k \in \mathbb{R}^d$ and $t \geq 0$,*

$$|\widehat{\rho_Q}(t,k)| \leq \frac{C\varepsilon}{\langle k, kt\rangle^{\sigma_1}}. \tag{17}$$

- *Scattering: there exists an operator $Q_\infty \in \mathcal{H}^{\sigma_0}_M(\mathbb{R}^d;\mathbb{C})$ such that, for all $t \geq 0$,*

$$\left\| e^{-i(\hbar/2)t\Delta} Q(t) e^{i(\hbar/2)t\Delta} - Q_\infty \right\|_{\mathcal{H}^{\sigma_0}_M} \leq \frac{C\varepsilon}{\langle t\rangle^{d/2}}. \tag{18}$$

Lemma 1.4. *Suppose $Q \in \mathcal{H}^0_{M,M}(\mathbb{R}^d;\mathbb{C})$. Then the Wigner transform of Q satisfies, for all $k, \eta \in \mathbb{R}^d$,*

$$\left|\widehat{W[Q]}(k,\eta)\right| \lesssim_d \|Q\|_{\mathcal{H}^0_{M,M}}.$$

证明. For $0 < \widetilde{\delta} \leq 1/2$, Definition 0.2, Cauchy–Schwarz and Sobolev embedding give

$$\begin{aligned}
\left|\widehat{W[Q]}(k,\eta)\right| &= \left| \int_x e^{-ik\cdot x} Q\left(x - \tfrac{\hbar}{2}\eta, x + \tfrac{\hbar}{2}\eta\right) dx \right| \\
&\lesssim \left(\int_x \left| \langle x\rangle^{d/2+\widetilde{\delta}} Q\left(x - \tfrac{\hbar}{2}\eta, x + \tfrac{\hbar}{2}\eta\right) \right|^2 dx \right)^{1/2} \\
&\lesssim \left(\int_{x,y} \left| \langle \nabla_y\rangle^{d/2+\widetilde{\delta}} \left[\langle x\rangle^{d/2+\widetilde{\delta}} Q\left(x - \tfrac{\hbar}{2}y, x + \tfrac{\hbar}{2}y\right) \right] \right|^2 dxdy \right)^{1/2} \\
&= \left(\int_{x,y} \left| \langle \tfrac{\hbar}{2}(\nabla_x - \nabla_y)\rangle^{d/2+\widetilde{\delta}} \langle x\rangle^{d/2+\widetilde{\delta}} Q\left(x - \tfrac{\hbar}{2}y, x + \tfrac{\hbar}{2}y\right) \right|^2 dxdy \right)^{1/2} \\
&= \hbar^{-d/2} \left\| \langle \boldsymbol{\xi}\rangle^{d/2+\widetilde{\delta}} \langle \boldsymbol{x}\rangle^{d/2+\widetilde{\delta}} Q(x,y) \right\|_{L^2_{x,y}} \\
&\lesssim \hbar^{-d/2} \sum_{|\alpha|,|\beta| \leq M} \left\| \boldsymbol{\xi}^\alpha \boldsymbol{x}^\beta Q(x,y) \right\|_{L^2_{x,y}},
\end{aligned}$$

where the final inequality uses that $M := \lceil (d+1)/2 \rceil$ is an integer with $M \geq d/2 + \widetilde{\delta}$, together with the commutation of $\boldsymbol{\xi}$ and $\boldsymbol{x}$ (Remark 0.3). The lemma then follows from the definition of the quantum Sobolev norm. □

The operators defined in Definition 0.2 interact nicely with the Wigner transform:

Lemma 1.5. *Suppose $Q \in \mathcal{L}^2(\mathbb{R}^d;\mathbb{C})$. Then the Wigner transform of Q satisfies the following:*

$$\begin{aligned}
\nabla_x W[Q](x,\xi) &= W[\boldsymbol{\nabla}_x Q](x,\xi), & ik\widehat{W[Q]}(k,\eta) &= \widehat{W[\boldsymbol{\nabla}_x Q]}(k,\eta),\\
\nabla_\xi W[Q](x,\xi) &= W[\boldsymbol{\nabla}_\xi Q](x,\xi), & i\eta\widehat{W[Q]}(k,\eta) &= \widehat{W[\boldsymbol{\nabla}_\xi Q]}(k,\eta),\\
xW[Q](x,\xi) &= W[\boldsymbol{x}Q](x,\xi), & i\nabla_k\widehat{W[Q]}(k,\eta) &= \widehat{W[\boldsymbol{x}Q]}(k,\eta),\\
\xi W[Q](x,\xi) &= W[\boldsymbol{\xi}Q](x,\xi), & i\nabla_\eta\widehat{W[Q]}(k,\eta) &= \widehat{W[\boldsymbol{\xi}Q]}(k,\eta).
\end{aligned}$$

证明. These statements follow directly from the definitions of the Wigner transform (Definition 0.6) and the quantum operators (Definition 0.2). □

The L^2 norm of $W[Q]$ is related to the $\mathcal{L}^2$ norm of Q via the following relation:

Lemma 1.6. *Suppose $Q \in \mathcal{L}^2(\mathbb{R}^d;\mathbb{C})$. Then the Wigner transform of Q satisfies*

$$\|W[Q](x,\xi)\|_{L^2_{x,\xi}} = (2\pi\hbar)^{-d/2}\|Q(x,y)\|_{L^2_{x,y}} = \|Q\|_{\mathcal{L}^2}.$$

证明. The result follows from the observation that

$$W[Q](x,\xi) = (2\pi\hbar)^{-d}\mathcal{F}\left[Q\left(x+\tfrac{\cdot}{2}, x-\tfrac{\cdot}{2}\right)\right](\xi/\hbar),$$

combined with Plancherel's theorem. □

Remark 1.7. Combining Lemma 1.5 with Lemma 1.6 shows that the Wigner transform is an isometry between the quantum and classical weighted Sobolev scales: for every $\sigma \geq 0$, every integer $M \geq 0$ and every $Q \in \mathcal{H}^\sigma_M(\mathbb{R}^d;\mathbb{C})$,

$$\|W[Q]\|_{H^\sigma_M} = \|Q\|_{\mathcal{H}^\sigma_M}.$$

Indeed, the Wigner transform maps the quantum operators $\boldsymbol{\nabla}_x$, $\boldsymbol{\nabla}_\xi$ and $\boldsymbol{\xi}$ to ∇_x, ∇_ξ and multiplication by ξ, respectively, so that $\xi^\alpha\langle\nabla_x,\nabla_\xi\rangle^\sigma W[Q] = W\left[\boldsymbol{\xi}^\alpha\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^\sigma Q\right]$; the identity then follows term by term from Lemma 1.6.

For an operator Q, the density $\rho_Q(x) := Q(x,x)$ is, formally, the velocity marginal of the Wigner transform, $\rho_Q(x) = \int_\xi W[Q](x,\xi)\,d\xi$: on the Fourier side, setting $\eta = 0$ in the Fourier representation of the Wigner transform above,

$$\widehat{\rho_Q}(k) = \widehat{W[Q]}(k,0) = \frac{1}{(2\pi)^d}\int_p \widehat{Q}\left(\tfrac{k}{2}+p, \tfrac{k}{2}-p\right)\,dp.$$

The following lemma makes this representation rigorous.

Lemma 1.8. *Suppose $Q \in \mathcal{H}^0_{M,M}(\mathbb{R}^d;\mathbb{C})$, where $M = \lceil (d+1)/2 \rceil$, identified with its integral kernel $Q = Q(x,y)$. Then the density $\rho_Q(x) := Q(x,x)$ satisfies, for all $k \in \mathbb{R}^d$,*

$$\widehat{\rho_Q}(k) = \frac{1}{(2\pi)^d} \int_p \widehat{Q}(k-p,p)\, dp = \frac{1}{(2\pi)^d} \int_p \widehat{Q}\left(\tfrac{k}{2}+p, \tfrac{k}{2}-p\right)\, dp,$$

both integrals converging absolutely.

证明. First suppose that the kernel $Q(x,y)$ is a Schwartz function on $\mathbb{R}^d \times \mathbb{R}^d$. By Fourier inversion and Fubini's theorem,

$$\begin{aligned}\rho_Q(x) = Q(x,x) &= \frac{1}{(2\pi)^{2d}} \int_{k'} \int_p e^{i(k'+p)\cdot x} \widehat{Q}(k',p)\, dp\, dk' \\ &= \frac{1}{(2\pi)^d} \int_k e^{ik\cdot x} \left[\frac{1}{(2\pi)^d} \int_p \widehat{Q}(k-p,p)\, dp \right] dk,\end{aligned}$$

where the final equality follows from the substitution $k' = k - p$. Hence ρ_Q is the inverse Fourier transform of the bracketed function, which gives the first representation; the second follows from the change of variables $p \mapsto \frac{k}{2} - p$.

It remains to extend the identity to $Q \in \mathcal{H}^0_{M,M}$. Set $F_Q(k,p) := \widehat{Q}\left(\frac{k}{2}+p, \frac{k}{2}-p\right)$. Since $M > d/2$, the Cauchy–Schwarz inequality in p and the Sobolev embedding in k give

$$\begin{aligned}\sup_k \int_p |F_Q(k,p)|\, dp &\lesssim_d \hbar^{-d/2} \sup_k \left\| \langle \hbar p \rangle^M F_Q(k,\cdot) \right\|_{L^2_p} \\ &\lesssim_d \hbar^{-d/2} \sum_{|\beta| \le M} \left\| \partial_k^\beta \left(\langle \hbar p \rangle^M F_Q \right) \right\|_{L^2_{k,p}} \\ &\lesssim_d \|Q\|_{\mathcal{H}^0_{M,M}},\end{aligned}$$

where the final inequality bounds $\langle \hbar p \rangle^M$ by a finite sum of absolute monomials (M being an integer) and uses the relations

$$\widehat{\boldsymbol{x} Q}\left(\tfrac{k}{2}+p, \tfrac{k}{2}-p\right) = i \nabla_k F_Q(k,p), \qquad \widehat{\boldsymbol{\xi} Q}\left(\tfrac{k}{2}+p, \tfrac{k}{2}-p\right) = \hbar p\, F_Q(k,p),$$

iterated, together with Plancherel's theorem and the normalisation of the quantum Lebesgue norm. Thus the integrals in the statement converge absolutely, uniformly in k.

For each fixed k, the estimate above shows that the right-hand side of the stated identity depends continuously on $Q \in \mathcal{H}^0_{M,M}$, while, since $\widehat{\rho_Q}(k) = \widehat{W[Q]}(k,0)$ by the Fourier representation of the Wigner transform, Lemma 1.4 shows the same for the left-hand side. The identity therefore extends to all of $\mathcal{H}^0_{M,M}$ by density of Schwartz kernels. □

Proposition 1.9 (Phase mixing for the free Schrödinger equation)**.** *Fix $N_0 \geq 0$. Then for any $t \geq 0$ and $k \in \mathbb{R}^d$, the density of the solution Q_{FS} to (4) satisfies*

$$\left|\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k)\right| \lesssim_d \frac{\|Q_{\mathrm{in}}\|_{\mathcal{H}^{N_0}_{M,M}}}{\langle k, kt\rangle^{N_0}}.$$

证明. First recall that $\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k) = \widehat{W[Q_{\mathrm{in}}]}(k,kt)$. Since $\boldsymbol{\nabla}_x$ and $\boldsymbol{\nabla}_\xi$ commute (Remark 0.3), the relations of Lemma 1.5 extend from the individual operators to the joint weight $\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{N_0}$, whence

$$\langle k, kt\rangle^{N_0}\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k) = \langle k,\eta\rangle^{N_0}\widehat{W[Q_{\mathrm{in}}]}(k,\eta)\Big|_{\eta=kt} = \widehat{W\left[\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{N_0} Q_{\mathrm{in}}\right]}(k,kt).$$

The result then follows from Lemma 1.4. □

1.3 Fourier and Wigner formulations of the Hartree equation

This subsection rewrites the nonlinear Hartree equation in both Fourier and Wigner form, making the structure of the linear and nonlinear terms explicit.

1.3.1 Fourier formulation

The nonlinear Hartree equation (3) takes the form

$$i\hbar\partial_t Q(t,x,y) = -\frac{\hbar^2}{2}\left(\Delta_x - \Delta_y\right) Q(t,x,y) + \big(V(t,x) - V(t,y)\big)\big(\gamma_g(x,y) + Q(t,x,y)\big),$$

where $V(s,x) = w * \rho_Q(s,x)$. Applying Duhamel's formula yields an implicit expression for Q:

$$\begin{aligned} Q(t,x,y) &= e^{i(\hbar/2)(\Delta_x-\Delta_y)t}Q_{\mathrm{in}}(x,y) \\ &\quad - \frac{i}{\hbar}\int_0^t e^{i(\hbar/2)(\Delta_x-\Delta_y)(t-s)}\left[V(s,x) - V(s,y)\right]\left(\gamma_g(x,y) + Q(s,x,y)\right)\, ds. \end{aligned}$$

Taking the Fourier transform in both spatial variables gives

$$\begin{aligned} &\widehat{Q}(t,k,p) = e^{-i(\hbar/2)(|k|^2-|p|^2)t}\widehat{Q_{\mathrm{in}}}(k,p) \\ &-\frac{i}{\hbar}\frac{1}{(2\pi)^d}\int_0^t e^{-i(\hbar/2)(|k|^2-|p|^2)(t-s)}\left[\widehat{V}(s,k)*_k - \widehat{V}(s,p)*_p\right]\left(\widehat{\gamma_g}(k,p) + \widehat{Q}(s,k,p)\right) ds. \end{aligned}$$

Recalling that $\widehat{\gamma_g}(k,p) = (2\pi)^{2d}\hbar^d g(\hbar k)\,\delta(p+k)$, where δ denotes the Dirac delta, and $\widehat{V}(t,k) = \widehat{w}(k)\widehat{\rho_Q}(t,k)$ gives

$$\begin{aligned} &\widehat{Q}(t,k,p) = e^{-i(\hbar/2)(|k|^2-|p|^2)t}\widehat{Q_{\mathrm{in}}}(k,p) \\ &- i\hbar^{d-1}(2\pi)^d\widehat{w}(k+p)\left[g(-\hbar p) - g(\hbar k)\right]\int_0^t e^{-i(\hbar/2)(|k|^2-|p|^2)(t-s)}\widehat{\rho_Q}(s,k+p)\, ds \\ &-\frac{i}{\hbar}\frac{1}{(2\pi)^d}\int_0^t\int_\ell e^{-i(\hbar/2)(|k|^2-|p|^2)(t-s)}\widehat{w}(\ell)\widehat{\rho_Q}(s,\ell)\big(\widehat{Q}(s,k-\ell,p) - \widehat{Q}(s,k,p-\ell)\big)\, d\ell ds. \end{aligned}$$

1.3.2 Conjugation by the free Schrödinger flow

Recall the operator P obtained by conjugating Q along the free evolution:

$$Q(t,x,y) = e^{i(\hbar/2)(\Delta_x - \Delta_y)t} P(t,x,y), \quad \widehat{Q}(t,k,p) = e^{-i(\hbar/2)(|k|^2-|p|^2)t}\widehat{P}(t,k,p). \tag{19}$$

This conjugation reverses Q along the free flow, removing the dominant free dynamics and allowing P to encode the interaction-driven dynamics alone. The operator P thus provides a natural change of variables in which the regularity of the initial data is expected to be largely preserved, in contrast to Q, which is subject to the mixing effects of the free evolution. At the level of the Wigner transform, this conjugation is a free-transport change of variables. The conjugation relation (19), together with $|k/2 + p|^2 - |k/2 - p|^2 = 2k \cdot p$, gives

$$\begin{aligned}\widehat{W[Q]}(t,k,\eta) &= \frac{1}{(2\pi)^d}\int_p e^{-i\hbar\eta\cdot p}\widehat{Q}\left(t, \tfrac{k}{2}+p, \tfrac{k}{2}-p\right)\, dp \\ &= \frac{1}{(2\pi)^d}\int_p e^{-i\hbar(\eta+kt)\cdot p}\widehat{P}\left(t, \tfrac{k}{2}+p, \tfrac{k}{2}-p\right)\, dp = \widehat{W[P]}(t,k,\eta+kt).\end{aligned}$$

Writing z for the spatial variable carried along the free flow and v for the velocity, this is equivalent to

$$W[P](t,z,v) = W[Q](t, z+tv, v).$$

Thus $W[P]$ is $W[Q]$ read along the free characteristics $x = z + tv$, the exact quantum analogue of the classical relation $g(t,z,v) = f(t, z+tv, v)$ between a profile g and a solution f of the Vlasov equation.

The conjugated perturbation P satisfies the integral equation

$$\widehat{P}(t,k,p) = \widehat{Q_{\mathrm{in}}}(k,p) + \int_0^t \widehat{\mathrm{L}_P}(s,k,p)\, ds + \int_0^t \widehat{\mathrm{N}_P}(s,k,p)\, ds. \tag{20}$$

Here L_P denotes the linearised contribution, arising from the interaction of the perturbation with the homogeneous equilibrium, and is defined by

$$\widehat{\mathrm{L}_P}(s,k,p) := -i\hbar^{d-1}(2\pi)^d \widehat{w}(k+p)\,[g(-\hbar p) - g(\hbar k)]\, e^{i(\hbar/2)(|k|^2-|p|^2)s}\widehat{\rho_Q}(s,k+p).$$

The term N_P denotes the nonlinear self-interaction of the perturbation and is defined by

$$\begin{aligned}\widehat{\mathrm{N}_P}(s,k,p) := -\frac{i}{\hbar}\frac{1}{(2\pi)^d}\int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(s,\ell)\Big(&e^{-i(\hbar/2)(|k-\ell|^2-|k|^2)s}\widehat{P}(s,k-\ell,p) \\ &- e^{-i(\hbar/2)(|p|^2-|p-\ell|^2)s}\widehat{P}(s,k,p-\ell)\Big)\, d\ell.\end{aligned}$$

By Lemma 1.8 and the Fourier conjugation relation (19), the density $\widehat{\rho_Q}$ can be expressed in terms of P as

$$\widehat{\rho_Q}(t,k) = \frac{1}{(2\pi)^d}\int_p e^{-i\hbar p\cdot kt}\widehat{P}(t,\tfrac{k}{2}+p,\tfrac{k}{2}-p)\,dp. \tag{21}$$

Substituting the representation (20) into this formula and applying the changes of variables $p \mapsto p-\ell/2$ and $p\mapsto p+\ell/2$ in the two nonlinear terms leads to

$$\begin{aligned}\widehat{\rho_Q}(t,k) &= \widehat{\rho_{Q_{\mathrm{FS}}}}(t,k)\\ &\quad - i\hbar^{d-1}\widehat{w}(k)\int_p \left[g\left(\hbar(p-\tfrac{k}{2})\right) - g\left(\hbar(p+\tfrac{k}{2})\right)\right]\int_0^t e^{-i\hbar p\cdot k(t-s)}\widehat{\rho_Q}(s,k)\,dsdp\\ &\qquad - \frac{2\hbar^{-1}}{(2\pi)^{2d}}\int_0^t\int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(s,\ell)\sin\left(\tfrac{\hbar}{2}\ell\cdot k(t-s)\right)\\ &\qquad\qquad \times\int_p e^{-i\hbar p\cdot(kt-\ell s)}\,\widehat{P}\left(s,\tfrac{k-\ell}{2}+p,\tfrac{k-\ell}{2}-p\right)\,dpd\ell ds. \end{aligned}\tag{22}$$

The sine factor in the nonlinear term follows from the identity

$$e^{-i(\hbar/2)\ell\cdot k(t-s)} - e^{i(\hbar/2)\ell\cdot k(t-s)} \equiv -2i\sin\big(\tfrac{\hbar}{2}\ell\cdot k(t-s)\big).$$

1.3.3 Wigner formulation and derivative bound

Taking the Wigner transform of equation (20), as defined in Definition 0.6, yields the following evolution equation for $\widehat{W[P]}$:

$$\begin{cases}\partial_t\widehat{W[P]}(t,k,\eta) = \widehat{W[\mathrm{L}_P]}(t,k,\eta) + \widehat{W[\mathrm{N}_P]}(t,k,\eta),\\ \widehat{W[\mathrm{L}_P]}(t,k,\eta) = -2\hbar^{-1}\widehat{w}(k)\widehat{\rho_Q}(t,k)\sin\left(\tfrac{\hbar}{2}k\cdot(\eta-kt)\right)\widehat{g}(\eta-kt),\\ \widehat{W[\mathrm{N}_P]}(t,k,\eta) = -\dfrac{2\hbar^{-1}}{(2\pi)^d}\displaystyle\int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(t,\ell)\sin\left(\tfrac{\hbar}{2}\ell\cdot(\eta-kt)\right)\\ \qquad\qquad\qquad\qquad\qquad\qquad \times\widehat{W[P]}(t,k-\ell,\eta-\ell t)\,d\ell,\end{cases}\tag{23}$$

where L_P and N_P are defined in (20).

Essential to the analysis are bounds on derivatives of the Wigner transform in the η variable, $D_\eta^\alpha\widehat{W[P]}$, corresponding to velocity weights $\boldsymbol{\xi}$ on P. For multi-indices α with $|\alpha|\geq 2$, additional terms of order $O(\hbar)$ arise from differentiation of the oscillatory phase, in stark contrast to the classical case.

Lemma 1.10. *Let α be a multi-index. The Wigner transform of L_P satisfies*

$$\begin{aligned}\left|D_\eta^\alpha\widehat{W[\mathrm{L}_P]}(t,k,\eta)\right| &\lesssim |k||\widehat{w}(k)||\widehat{\rho_Q}(t,k)||\eta-kt|\left|D_\eta^\alpha\widehat{g}(\eta-kt)\right|\\ &\quad + |k|\langle\hbar k\rangle^{|\alpha|-1}|\widehat{w}(k)||\widehat{\rho_Q}(t,k)|\sum_{|\beta|\leq|\alpha|-1}\left|D_\eta^\beta\widehat{g}(\eta-kt)\right|.\end{aligned}$$

The Wigner transform of N_P *satisfies*

$$\begin{aligned}\big|D_\eta^\alpha \widehat{W[\mathrm{N}_P]}(t,k,\eta)\big| &\lesssim \int_\ell |\ell|\,|\widehat{w}(\ell)|\,|\widehat{\rho_Q}(t,\ell)|\,|\eta-kt|\,\big|D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t)\big|\,d\ell \\ &+ \sum_{|\beta|\le|\alpha|-1}\int_\ell |\ell|\,\langle\hbar\ell\rangle^{|\alpha|-1}\,|\widehat{w}(\ell)|\,|\widehat{\rho_Q}(t,\ell)|\,\big|D_\eta^\beta \widehat{W[P]}(t,k-\ell,\eta-\ell t)\big|\,d\ell.\end{aligned}$$

When $|\alpha|=0$*, sums over* $|\beta|\le|\alpha|-1$ *are understood to be empty.*

证明. Applying the Leibniz rule to the linear term gives

$$D_\eta^\alpha\left[\sin\left(\tfrac{\hbar}{2}k\cdot(\eta-kt)\right)\widehat{g}(\eta-kt)\right]=\sum_{\beta\le\alpha}\binom{\alpha}{\beta}D_\eta^{\alpha-\beta}\left[\sin\left(\tfrac{\hbar}{2}k\cdot(\eta-kt)\right)\right]D_\eta^\beta\widehat{g}(\eta-kt).$$

For the term with $\beta=\alpha$, use $|\sin x|\le|x|$ to get

$$\left|\sin\left(\tfrac{\hbar}{2}k\cdot(\eta-kt)\right)\right|\le\hbar|k||\eta-kt|.$$

For terms with $\beta<\alpha$, differentiate sin directly and use $|\sin x|,|\cos x|\le 1$ to obtain

$$\left|D_\eta^{\alpha-\beta}\left[\sin\left(\tfrac{\hbar}{2}k\cdot(\eta-kt)\right)\right]\right|\le(\hbar|k|)^{|\alpha-\beta|}.$$

This leads to the estimate

$$\begin{aligned}&\left|D_\eta^\alpha\left[\left(e^{-i(\hbar/2)k\cdot(\eta-kt)}-e^{i(\hbar/2)k\cdot(\eta-kt)}\right)\widehat{g}(\eta-kt)\right]\right| \\ &\qquad\lesssim\hbar|k||\eta-kt|\big|D_\eta^\alpha\widehat{g}(\eta-kt)\big|+\sum_{\beta<\alpha}(\hbar|k|)^{|\alpha-\beta|}\big|D_\eta^\beta\widehat{g}(\eta-kt)\big|.\end{aligned}$$

Since for any $\beta<\alpha$, $(\hbar|k|)^{|\alpha-\beta|}\lesssim\hbar|k|\langle\hbar k\rangle^{|\alpha|-1}$, this gives the bound for $\widehat{W[\mathrm{L}_P]}$. The estimate for $\widehat{W[\mathrm{N}_P]}$ follows similarly. □

Lemma 1.11. *Fix* $\sigma>0$ *and let* $\rho_{Q_{\mathrm{FS}}}$ *be the density of the solution to the free Schrödinger equation* (4)*, defined in* (5)*. Then, for* $s_1>(d+1)/2$ *and* $s_2>1/2$*,*

$$\begin{aligned}\big\||k|^{1/2}\langle k,kt\rangle^\sigma\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k)\big\|_{L_t^2([0,\infty),L_k^2)}&\lesssim_{s_1}\|Q_{\mathrm{in}}\|_{\mathfrak{H}^{\sigma+s_1}_{M,M}}\\ \big\||k|^{1/2}\langle k,kt\rangle^\sigma\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k)\big\|_{L_k^\infty L_t^2([0,\infty))}&\lesssim_{s_2}\|Q_{\mathrm{in}}\|_{\mathfrak{H}^{\sigma+s_2}_{M,M}}.\end{aligned}$$

证明. Recalling that $\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k)=\widehat{W[Q_{\mathrm{in}}]}(k,kt)$,

$$\begin{aligned}&\big\||k|^{1/2}\langle k,kt\rangle^\sigma\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k)\big\|_{L_t^2([0,\infty),L_k^2)}\\ &\qquad=\left(\int_{t=0}^\infty\int_k|k|\langle k,kt\rangle^{2\sigma}\left|\widehat{W[Q_{\mathrm{in}}]}(k,kt)\right|^2dkdt\right)^{1/2}\\ &\qquad\le\sup_{k,\eta}\left|\langle k,\eta\rangle^{\sigma+s_1}\widehat{W[Q_{\mathrm{in}}]}(k,\eta)\right|\left(\int_k\int_{t=0}^\infty\frac{|k|}{\langle k,kt\rangle^{2s_1}}\,dtdk\right)^{1/2}.\end{aligned}$$

Then, by making the change of variables $t \mapsto t/|k|$, and denoting $\overline{\varepsilon} = s_1 - \frac{d+1}{2}$,

$$\int_k \int_{t=0}^{\infty} \frac{|k|}{\langle k, kt\rangle^{2s_1}}\, dtdk = \int_k \int_{t=0}^{\infty} \frac{1}{\langle k, t\rangle^{d+1+2\overline{\varepsilon}}}\, dtdk \lesssim \int_k \frac{1}{\langle k\rangle^{d+\overline{\varepsilon}}}\, dk \int_{\mathbb{R}} \frac{1}{\langle t\rangle^{1+\overline{\varepsilon}}}\, dt \lesssim 1.$$

The bound then follows by applying Lemma 1.4. The second estimate is proved similarly. □

1.3.4 Bootstrap proposition and strategy

The following proposition forms the backbone of the nonlinear analysis. The inequalities (I1)–(I5) below are referred to as the *bootstrap improvements*: they coincide with the bootstrap controls (B1)–(B5), but with each constant $4K_i$ halved to $2K_i$. The proposition thus asserts that, on any interval on which the controls hold, the solution in fact obeys strictly stronger bounds. By continuity, these strict improvements allow the bootstrap controls to be propagated throughout the maximal lifespan. The continuation argument is given at the end of the proof.

Proposition 1.12 (Bootstrap)**.** *Let the assumptions on w, g and Q_{in} be as in Theorem 1.3. There exist $\varepsilon_0 = \varepsilon_0(N_0, w, g, c_{\mathrm{P}}, d, \delta_0, \sigma_i)$ and $K_i = K_i(N_0, w, g, c_{\mathrm{P}}, d, \delta_0, \sigma_i)$ such that, for every $T^* \in (0, T_{\max})$, if the bootstrap controls* (B1)–(B5) *hold on $[0, T^*]$ and $\varepsilon < \varepsilon_0$, then*

$$\sum_{|\alpha|\le M} \left\|\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_4} \langle t\boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle \left(\boldsymbol{\xi}^\alpha P(t)\right)\right\|_{\mathcal{L}^2}^2 \le 2K_1 \langle t\rangle^{2d-1}\varepsilon^2, \tag{I1}$$

$$\left\|\langle \hbar k\rangle |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k)\right\|_{L_t^2([0,T^*],L_k^2)}^2 \le 2K_2\varepsilon^2, \tag{I2}$$

$$\sum_{|\alpha|\le M} \left\|\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_3} |\boldsymbol{\nabla}_x|^{\delta} \left(\boldsymbol{\xi}^\alpha P(t)\right)\right\|_{\mathcal{L}^2}^2 \le 2K_3\varepsilon^2, \tag{I3}$$

$$\left\||k|^{1/2} \langle k, kt\rangle^{\sigma_2} \widehat{\rho_Q}(t,k)\right\|_{L_k^\infty L_t^2([0,T^*])}^2 \le 2K_4\varepsilon^2, \tag{I4}$$

$$\langle k, \eta\rangle^{2\sigma_1} \left|\widehat{W[P]}(t,k,\eta)\right|^2 \le 2K_5\varepsilon^2, \tag{I5}$$

for all $t \in [0, T^*]$. Consequently, the maximal solution is global and the bootstrap controls hold for all $t \ge 0$.

Remark 1.13. Among the improvements, (I2) presents the main analytical difficulty, as it permits neither regularity loss nor time growth. Precise control is required over the nonlinear interaction term in (22), which includes a bilinear contribution in the perturbation and allows earlier density modes $\widehat{\rho_Q}(s,\ell)$ to influence the evolution of $\widehat{\rho_Q}(t,k)$. In certain regimes, this interaction may

accumulate and produce *resonances* at large times, see e.g. [1]. In the Hartree equation on $\mathbb{R}^d$ with $d \geq 3$ and sufficiently regular w, such resonances are controlled by a dispersive mechanism. The mechanism relies on separating the nearly collinear interactions, for which

$$|\ell_\perp| \lesssim \frac{|k|^b}{\langle s\rangle^\zeta}, \qquad \ell_\perp := \ell - \frac{k\cdot\ell}{|k|^2}k,$$

from the non-resonant interactions, for suitable exponents $b, \zeta \in (0,1)$. The resonant set is a shrinking cylinder around the line spanned by k, whose decreasing cross-section provides sufficient smallness. On its complement, the bound $|kt - \ell s| \geq s|\ell_\perp|$ yields additional decay. The resulting time-response kernel matches that arising in the Vlasov–Poisson system and is handled similarly in Section 2.2.3.

The proposition is proved below using the estimates established in Sections 2 and 3.

Proof of Proposition 1.12. Fix dimension $d \geq 3$, constants σ_0, σ_1, N_0, kernel w, steady state g and initial perturbation Q_{in} as in Theorem 1.3 and choose parameters σ_2, σ_3, σ_4 satisfying (6). Section 2 establishes that the constants K_2 and K_4 appearing in the bootstrap assumptions (B2) and (B4) may be chosen depending only on the fixed setup. Section 3 then shows that the remaining constants K_1, K_3, and K_5 can also be selected depending only on the setup and on K_2, K_4. If $\varepsilon_0 > 0$ satisfies

$$\varepsilon_0 \leq \min\Big\{\frac{1}{\sqrt{K_3K_5 + K_2K_5}}, \frac{1}{\sqrt{K_4K_5} + \sqrt{K_3K_5}}, \frac{1}{\sqrt{K_2K_3} + \sqrt{K_1K_5} + \sqrt{K_3K_5}}\Big\},$$

then, for any $0 < \varepsilon \leq \varepsilon_0$, all bootstrap improvements (I1)–(I5) hold by Sections 2.2, 2.3, 3.3, 3.1 and 3.2.

The proof is concluded by a standard continuity and continuation argument. Local well-posedness and the initial estimates imply that the bootstrap controls hold for sufficiently small times. Proposition 1.12 gives strict improvements on every interval on which the controls hold. By continuity, the bootstrap estimates therefore persist throughout $[0, T_{\max})$. In particular, (I1) bounds the lower-order Sobolev norm entering the local continuation criterion on every finite interval. Hence $T_{\max} = \infty$. Persistence of regularity retains the regularity required above on every finite time interval. □

Lemma 1.14. *The global bootstrap control* (B5) *implies that, for all* $t \geq 0$ *and* $k \in \mathbb{R}^d$,

$$|\widehat{\rho_Q}(t,k)| \leq \frac{2\sqrt{K_5}\,\varepsilon}{\langle k, kt\rangle^{\sigma_1}}.$$




图 1: Bootstrap dependencies: each arrow indicates an assumption used to establish an improvement, e.g. (B1) $\to$ (B2) reflects the use of (B1) in proving (I2).

In particular, Proposition 1.12 implies the damping estimate (17) *in Theorem 1.3.*

证明. The Fourier transform of the Wigner transform gives

$$\widehat{W[Q]}(t,k,0) = \widehat{\rho_Q}(t,k),$$

while the conjugation identity of Section 1.3.2 gives

$$\widehat{W[Q]}(t,k,0) = \widehat{W[P]}(t,k,kt).$$

Hence

$$|\widehat{\rho_Q}(t,k)| = \left|\widehat{W[P]}(t,k,kt)\right| \le \frac{2\sqrt{K_5}\,\varepsilon}{\langle k,kt\rangle^{\sigma_1}},$$

by bootstrap assumption (B5) evaluated at $\eta = kt$. Since Proposition 1.12 shows that the bootstrap control (B5) holds globally, the estimate (17) follows with $C = 2\sqrt{K_5}$. □

Bootstrap assumptions (B3) and (B5) suffice to deduce the scattering statement (18).

Lemma 1.15. *Proposition 1.12 implies the scattering statement* (18) *in Theorem 1.3.*

证明. Let $|\alpha| \le M$. To prove convergence of $P(t)$ in $\mathcal{H}^{\sigma_0}_M$, the Duhamel formula (20) is analysed in Fourier–Wigner variables. The aim is to show that both the linear and nonlinear contributions decay sufficiently fast to be absolutely

integrable in time in the $\mathcal{H}_M^{\sigma_0}$ norm, yielding the existence of an asymptotic state $Q_\infty := \lim_{t\to\infty} P(t)$.

For the linear term, apply Lemma 1.6 and Plancherel's theorem to obtain

$$\|\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_0}(\boldsymbol{\xi}^\alpha \mathrm{L}_P(t))\|_{\mathcal{L}^2}^2 \le \big\|\langle k,\eta\rangle^{\sigma_0} D_\eta^\alpha(\widehat{W[\mathrm{L}_P]}(t))\big\|_{L^2_{k,\eta}}^2.$$

Lemma 1.10 is used to extract the highest-order D_η^α derivative falling on $\widehat{W[\mathrm{L}_P]}$. Using assumption (16) on w, which gives $\langle \hbar k\rangle^{|\alpha|-1}|\widehat{w}(k)| \lesssim 1$, and the inequality $\langle k,\eta\rangle^{\sigma_0} \lesssim \langle k, kt\rangle^{\sigma_0}\langle \eta - kt\rangle^{\sigma_0}$, one obtains

$$\begin{aligned}
&\big\|\langle k,\eta\rangle^{\sigma_0} D_\eta^\alpha(\widehat{W[\mathrm{L}_P]}(t))\big\|_{L^2_{k,\eta}}^2 \\
&\lesssim \int_{k,\eta} |\widehat{w}(k)|^2|k|^2\langle k,kt\rangle^{2\sigma_0}|\widehat{\rho_Q}(t,k)|^2\langle \eta-kt\rangle^{2\sigma_0}|\eta-kt|^2\big|D_\eta^\alpha \widehat{g}(\eta-kt)\big|^2\,dkd\eta \\
&+ \sum_{|\beta|\le|\alpha|-1}\int_{k,\eta}\langle \hbar k\rangle^{2|\alpha|-2}|\widehat{w}(k)|^2|k|^2\langle k,kt\rangle^{2\sigma_0}|\widehat{\rho_Q}(t,k)|^2\langle \eta-kt\rangle^{2\sigma_0}\big|D_\eta^\beta \widehat{g}(\eta-kt)\big|^2\,dkd\eta \\
&\lesssim \|g\|_{H_M^{\sigma_0+1}}^2\int_k |k|^2\langle k,kt\rangle^{2\sigma_0}|\widehat{\rho_Q}(t,k)|^2\,dk \\
&\lesssim K_5\varepsilon^2\|g\|_{H_M^{\sigma_0+1}}^2\int_k \frac{|k|^2}{\langle k,kt\rangle^{2\sigma_1-2\sigma_0}}\,dk,
\end{aligned}$$

where the final bound follows from the bootstrap estimate (B5) (via Lemma 1.14). Provided $2\sigma_1 - 2\sigma_0 > d+2$, this integral is uniformly bounded in time after the change of variables $k \mapsto k/\langle t\rangle$, leading to

$$\|\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_0}(\boldsymbol{\xi}^\alpha \mathrm{L}_P(t))\|_{\mathcal{L}^2} \lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d/2+1}}.$$

The nonlinear term is handled similarly. Lemma 1.10 extracts the highest-order D_η^α derivative falling on $\widehat{W[P]}$, and the factor $\langle \hbar\ell\rangle^{|\alpha|-1}|\widehat{w}(\ell)| \lesssim 1$ on the lower-order terms is again controlled by (16). Using the inequalities $|\eta - kt| \lesssim \langle t\rangle\langle k-\ell, \eta-\ell t\rangle$ and $\langle k,\eta\rangle^{\sigma_0} \lesssim \langle \ell,\ell t\rangle^{\sigma_0}\langle k-\ell,\eta-\ell t\rangle^{\sigma_0}$, together with the bootstrap estimate (B5) (via Lemma 1.14), one obtains

$$\begin{aligned}
&\big\|\langle k,\eta\rangle^{\sigma_0} D_\eta^\alpha(\widehat{W[\mathrm{N}_P]}(t))\big\|_{L^2_{k,\eta}}^2 \\
&\qquad\lesssim K_5\varepsilon^2\langle t\rangle^2\sum_{|\beta|\le M}\int_{k,\eta}\Big(\int_\ell \frac{|\ell|}{\langle \ell,\ell t\rangle^{\sigma_1-\sigma_0}}\langle k-\ell,\eta-\ell t\rangle^{\sigma_0+1} \\
&\qquad\qquad\times\big|D_\eta^\beta\widehat{W[P]}(t,k-\ell,\eta-\ell t)\big|\,d\ell\Big)^2\,dkd\eta.
\end{aligned}$$

By Lemma 1.5, multipliers in the Fourier variables (k,η) and derivatives in η correspond, respectively, to the quantum derivatives $\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle$ and the velocity weights $\boldsymbol{\xi}$ acting on P. In particular,

$$\begin{aligned}
&\langle k-\ell,\eta-\ell t\rangle^{\sigma_0+1} D_\eta^\beta\widehat{W[P]}(t,k-\ell,\eta-\ell t) \\
&\qquad = (-i)^{|\beta|}\widehat{W[\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_0+1}(\boldsymbol{\xi}^\beta P)]}(t,k-\ell,\eta-\ell t).
\end{aligned}$$

In order to apply bootstrap assumption (B3), Cauchy–Schwarz in ℓ is applied with weight $|k-\ell|^{\delta}$, for $0<\delta\ll 1/2$ as introduced in Section 0.5.1, yielding

$$\begin{aligned}
&\Big(\int_\ell \frac{|\ell|}{\langle \ell,\ell t\rangle^{\sigma_1-\sigma_0}}\Big|\widehat{W[\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_0+1}(\boldsymbol{\xi}^\beta P)]}(t,k-\ell,\eta-\ell t)\Big|\,d\ell\Big)^2\\
&\qquad\le \Big[\int_\ell \frac{1}{|k-\ell|^{2\delta}}\,\frac{|\ell|}{\langle \ell,\ell t\rangle^{\sigma_1-\sigma_0}}\,d\ell\Big]\\
&\qquad\times\int_\ell \frac{|\ell|}{\langle \ell,\ell t\rangle^{\sigma_1-\sigma_0}}|k-\ell|^{2\delta}\Big|\widehat{W[\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_0+1}(\boldsymbol{\xi}^\beta P)]}(t,k-\ell,\eta-\ell t)\Big|^2 d\ell.
\end{aligned}$$

To bound the first factor by its supremum over k, set $q:=\langle t\rangle k$. The change of variables $\ell\mapsto \ell/\langle t\rangle$ sends $\langle \ell,\ell t\rangle$ to $\langle \ell\rangle$, so that

$$\int_\ell \frac{1}{|k-\ell|^{2\delta}}\,\frac{|\ell|}{\langle \ell,\ell t\rangle^{\sigma_1-\sigma_0}}\,d\ell=\frac{1}{\langle t\rangle^{d+1-2\delta}}\int_\ell \frac{1}{|q-\ell|^{2\delta}}\,\frac{|\ell|}{\langle \ell\rangle^{\sigma_1-\sigma_0}}\,d\ell. \tag{24}$$

Splitting the remaining integral into the regions $|q-\ell|\ge 1$, where $|q-\ell|^{-2\delta}\le 1$, and $|q-\ell|<1$, where the singularity is integrable as $2\delta<d$, shows that

$$\sup_k\int_\ell \frac{1}{|k-\ell|^{2\delta}}\,\frac{|\ell|}{\langle \ell,\ell t\rangle^{\sigma_1-\sigma_0}}\,d\ell\lesssim\frac{1}{\langle t\rangle^{d+1-2\delta}},$$

provided $\sigma_1-\sigma_0>d+1$. Inserting this bound yields

$$\begin{aligned}
&\big\|\langle k,\eta\rangle^{\sigma_0}D_\eta^\alpha(\widehat{W[\mathrm{N}_P]}(t))\big\|_{L^2_{k,\eta}}^2\\
&\lesssim K_5\varepsilon^2\langle t\rangle^2\frac{1}{\langle t\rangle^{d+1-2\delta}}\int_\ell\frac{|\ell|}{\langle \ell,\ell t\rangle^{\sigma_1-\sigma_0}}d\ell\sum_{|\beta|\le M}\Big\|\widehat{W[|\boldsymbol{\nabla}_x|^\delta\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_0+1}(\boldsymbol{\xi}^\beta P)]}(t)\Big\|_{L^2_{k,\eta}}^2\\
&\lesssim K_5\varepsilon^2\langle t\rangle^2\frac{1}{\langle t\rangle^{d+1-2\delta}}\frac{1}{\langle t\rangle^{d+1}}K_3\varepsilon^2,
\end{aligned}$$

where the final step uses bootstrap assumption (B3), together with Plancherel and Lemma 1.6. Hence, for the nonlinear term,

$$\|\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_0}(\boldsymbol{\xi}^\alpha\mathrm{N}_P(t))\|_{\mathcal{L}^2}\lesssim\frac{\sqrt{K_5K_3}\varepsilon^2}{\langle t\rangle^{d-\delta}}.$$

Consequently, define

$$Q_\infty:=Q_{\mathrm{in}}+\int_0^\infty \mathrm{L}_P(s)\,ds+\int_0^\infty \mathrm{N}_P(s)\,ds\in\mathcal{H}_M^{\sigma_0}.$$

Inequality (18) then follows from

$$\begin{aligned}
&\|P(t)-Q_\infty\|_{\mathcal{H}_M^{\sigma_0}}\\
&\lesssim\sum_{|\alpha|\le M}\Big(\int_t^\infty\|\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_0}(\boldsymbol{\xi}^\alpha\mathrm{L}_P(s))\|_{\mathcal{L}^2_{x,y}}\,ds+\int_t^\infty\|\langle \boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_0}(\boldsymbol{\xi}^\alpha\mathrm{N}_P(s))\|_{\mathcal{L}^2_{x,y}}\,ds\Big)\\
&\lesssim\sqrt{K_5}\varepsilon\int_t^\infty\frac{1}{\langle s\rangle^{d/2+1}}\,ds+\sqrt{K_5K_3}\varepsilon^2\int_t^\infty\frac{1}{\langle s\rangle^{d-\delta}}\,ds\\
&\lesssim\frac{\varepsilon}{\langle t\rangle^{d/2}},
\end{aligned}$$

as the contribution from the nonlinear term satisfies

$$\int_t^\infty \langle s\rangle^{-(d-\delta)}\,ds \lesssim \langle t\rangle^{-d+\delta+1} \le \langle t\rangle^{-d/2}$$

for $d \ge 3$ and $0 < \delta \ll 1/2$, ensuring the overall decay rate $\langle t\rangle^{-d/2}$. □

This completes the proof of the scattering statement (18), and hence of Theorem 1.3, subject to the bootstrap estimates established in the following sections.

Lemma 1.16. *Let $n \ge 0$, and let α be a multi-index. Then, for every $k \neq 0$,*

$$\int_0^\infty \langle s\rangle^n \big|\partial_k^\alpha \widehat{g}\big(\tfrac{k}{|k|}s\big)\big|\,ds \lesssim_{d,\delta_0,n,|\alpha|} \|g\|_{H^{n+1/2+\delta_0}_{|\alpha|+\lceil d/2\rceil}}.$$

证明. Applying Cauchy–Schwarz followed by Lemma 1.2,

$$\begin{aligned}\int_0^\infty \langle s\rangle^n \big|\partial_k^\alpha \widehat{g}\big(\tfrac{k}{|k|}s\big)\big|\,ds &\lesssim_{\delta_0} \left(\int_0^\infty \langle s\rangle^{2n+1+2\delta_0}\big|\partial_k^\alpha \widehat{g}\big(\tfrac{k}{|k|}s\big)\big|^2\,ds\right)^{1/2}\\ &\lesssim_d \left(\int_{\mathbb{R}^d_k}\Big|\langle\nabla_k\rangle^{\lceil d/2\rceil}\big(\langle k\rangle^{n+1/2+\delta_0}\partial_k^\alpha \widehat{g}(k)\big)\Big|^2\,dk\right)^{1/2}.\end{aligned}$$

Then, by Plancherel and Lemma 1.1:

$$\begin{aligned}\int_0^\infty \langle s\rangle^n \big|\partial_k^\alpha \widehat{g}\big(\tfrac{k}{|k|}s\big)\big|\,ds &\lesssim_d \big\|\langle x\rangle^{\lceil d/2\rceil}\langle\nabla_x\rangle^{n+1/2+\delta_0}\,(x^\alpha g(x))\big\|_{L^2}\\ &\lesssim \sum_{|\beta|\le\lceil d/2\rceil}\big\|x^\beta\langle\nabla_x\rangle^{n+1/2+\delta_0}\,(x^\alpha g(x))\big\|_{L^2}\\ &\lesssim_n \sum_{|\beta|\le\lceil d/2\rceil}\big\|\langle\nabla_x\rangle^{n+1/2+\delta_0}\,\big(x^{\beta+\alpha}g(x)\big)\big\|_{L^2}\\ &\lesssim_\alpha \|g\|_{H^{n+1/2+\delta_0}_{|\alpha|+\lceil d/2\rceil}}.\end{aligned}$$

□

Lemma 1.17 (Decay of $m_g(\lambda,k)$)**.** *For any integer $n \ge 0$, if $g \in H^{n+3/2+\delta_0}_{2+\lceil d/2\rceil}(\mathbb{R}^d)$, then*

i. $|\partial_\lambda^n m_g(i\tau,k)| \lesssim_{n,d,\delta_0} |k|^{-n}\|g\|_{H^{n+3/2+\delta_0}_{\lceil d/2\rceil}}$, *and*

ii. $|\partial_\lambda^n m_g(i\tau,k)| \lesssim_{n,d,\delta_0} \dfrac{\langle\hbar k\rangle^2|k|^{2-n}}{\langle\hbar k\rangle^2|k|^2+\tau^2}\|g\|_{H^{n+3/2+\delta_0}_{2+\lceil d/2\rceil}}$.

证明. Differentiating (15) n times in λ yields

$$(-1)^n\partial_\lambda^n m_g(\lambda,k) = \frac{1}{|k|^n}\frac{2}{\hbar|k|}\int_0^\infty e^{-\lambda s/|k|}s^n\sin(\tfrac{\hbar}{2}|k|s)\widehat{g}(\tfrac{k}{|k|}s)\,ds.$$

Hence, for any $\operatorname{Re}\lambda \ge 0$,

$$|\partial_\lambda^n m_g(\lambda,k)| \lesssim_n \frac{1}{|k|^n}\int_0^\infty s^{n+1}\big|\widehat{g}(\tfrac{k}{|k|}s)\big|\,ds \lesssim_{n,d,\delta_0} \frac{1}{|k|^n}\|g\|_{H^{n+3/2+\delta_0}_{\lceil d/2\rceil}}$$

using Lemma 1.16, which proves part (i).

To establish part (ii), consider $\lambda = i\tau \in i\mathbb{R}$ and $n \geq 2$. Multiply both sides by τ^2 and integrate by parts twice:

$$
\begin{aligned}
&\tau^2(-1)^n \partial_\lambda^n m_g(i\tau, k) \\
&= \frac{1}{|k|^n} \frac{2}{\hbar|k|} \int_0^\infty \tau^2 e^{-i\tau^s/|k|} s^n \sin(\tfrac{\hbar}{2}|k|s) \widehat{g}(\tfrac{k}{|k|}s)\, ds \\
&= \frac{|k|^2}{|k|^n} \frac{2}{\hbar|k|} \int_0^\infty (-\partial_s^2)\big(e^{-i\tau^s/|k|}\big) s^n \sin(\tfrac{\hbar}{2}|k|s) \widehat{g}(\tfrac{k}{|k|}s)\, ds \\
&= \frac{|k|^2}{|k|^n} \frac{2}{\hbar|k|} \int_0^\infty e^{-i\tau^s/|k|} \Big[\tfrac{\hbar^2}{4}|k|^2 s^n \sin(\tfrac{\hbar}{2}|k|s) \widehat{g}(\tfrac{k}{|k|}s) - n(n-1)s^{n-2} \sin(\tfrac{\hbar}{2}|k|s) \widehat{g}(\tfrac{k}{|k|}s) \\
&\qquad\qquad - 2ns^{n-1} \sin(\tfrac{\hbar}{2}|k|s) \frac{k}{|k|} \cdot \nabla_k \widehat{g}(\tfrac{k}{|k|}s) - \hbar|k| n s^{n-1} \cos(\tfrac{\hbar}{2}|k|s) \widehat{g}(\tfrac{k}{|k|}s) \\
&\qquad\qquad - \hbar|k| s^n \cos(\tfrac{\hbar}{2}|k|s) \tfrac{k}{|k|} \cdot \nabla_k \widehat{g}(\tfrac{k}{|k|}s) - s^n \sin(\tfrac{\hbar}{2}|k|s) (\tfrac{k}{|k|} \cdot \nabla_k)^2 \widehat{g}(\tfrac{k}{|k|}s) \Big] ds.
\end{aligned}
$$

The first term is exactly $(-1)^n(\hbar^2/4)|k|^4 \partial_\lambda^n m_g(i\tau, k)$. Moreover, all the other terms in the integrand are $O(\hbar|k|)$, and hence

$$
\begin{aligned}
&\tau^2 |\partial_\lambda^n m_g(i\tau, k)| \\
&\lesssim \hbar^2 |k|^4 |\partial_\lambda^n m_g(i\tau, k)| + |k|^{2-n} \int_0^\infty \langle s \rangle^{n+1} \Big(\big|\widehat{g}(\tfrac{k}{|k|}s)\big| + \big|\nabla_k \widehat{g}(\tfrac{k}{|k|}s)\big| + \big|\nabla_k^2 \widehat{g}(\tfrac{k}{|k|}s)\big| \Big) ds \\
&\lesssim \hbar^2 |k|^{4-n} \|g\|_{H^{n+3/2+\delta_0}_{\lceil d/2 \rceil}} + |k|^{2-n} \|g\|_{H^{n+3/2+\delta_0}_{2+\lceil d/2 \rceil}},
\end{aligned}
$$

again using Lemma 1.16. The cases $n = 0$ and $n = 1$ give rise to fewer terms upon the integration by parts and are treated similarly. Finally, combining this with the bound from part (i) gives

$$
(\langle \hbar k \rangle^2 |k|^2 + \tau^2) |\partial_\lambda^n m_g(i\tau, k)| \lesssim \langle \hbar k \rangle^2 |k|^{2-n} \|g\|_{H^{n+3/2+\delta_0}_{2+\lceil d/2 \rceil}},
$$

which proves part (ii). □

To quantify the deviation from the identity in the Green function, define the remainder

$$
\widetilde{\mathscr{G}^r}(\lambda, k) := \frac{\widetilde{\mathscr{L}}(\lambda, k)}{1 + \widetilde{\mathscr{L}}(\lambda, k)} = \frac{\widehat{w}(k) m_g(\lambda, k)}{1 + \widehat{w}(k) m_g(\lambda, k)},
$$

so that $\widetilde{\mathscr{G}}(\lambda, k) = 1 - \widetilde{\mathscr{G}^r}(\lambda, k)$.

Lemma 1.18 (Decay of $\widetilde{\mathscr{G}^r}(\lambda, k)$)**.** *Let $n \geq 0$ be an integer and suppose $w \in L^1(\mathbb{R}^d)$ and $g \in H^{n+3/2+\delta_0}_{2+\lceil d/2 \rceil}(\mathbb{R}^d)$ satisfy the uniform Penrose condition* (2) *with constant c_{P}. Then, there exists a constant $C = C(n, w, g, c_{\mathrm{P}}, d, \delta_0)$ such that*

$$
\big|\partial_\lambda^n \widetilde{\mathscr{G}^r}(i\tau, k)\big| \leq C \frac{\langle \hbar k \rangle^2 |k|^{2-n}}{\langle \hbar k \rangle^2 |k|^2 + \tau^2}.
$$

证明. For $n = 0$, the estimate follows directly from Lemma 1.17(ii) and the uniform Penrose condition. Using the Faà di Bruno formula, for any $n \geq 1$,

$$\frac{d^n}{dx^n}\Big[\frac{f(x)}{1+f(x)}\Big] = \sum_{\pi\in\Pi_n} \frac{(-1)^{|\pi|+1}|\pi|!}{(1+f(x))^{|\pi|+1}} \prod_{B\in\pi} f^{(|B|)}(x),$$

where π runs through the set Π_n of all the partitions of the set $\{1,\dots,n\}$. So,

$$\begin{aligned}\partial_\lambda^n \widetilde{\mathscr{G}^r}(i\tau,k) &= \frac{d^n}{d\lambda^n}\left[\frac{\widehat{w}(k)m_g(\lambda,k)}{1+\widehat{w}(k)m_g(\lambda,k)}\right]\Big|_{\lambda=i\tau} \\ &= \sum_{\pi\in\Pi_n} \frac{(-1)^{|\pi|+1}|\pi|!}{(1+\widehat{w}(k)m_g(i\tau,k))^{|\pi|+1}} \prod_{B\in\pi} \widehat{w}(k)\partial_\lambda^{|B|} m_g(i\tau,k).\end{aligned}$$

Moreover, writing $\pi = \{B_0\}\cup(\pi\setminus\{B_0\})$ and using Lemma 1.17(ii) to bound the term $\partial_\lambda^{|B_0|}m_g(i\tau,k)$ and Lemma 1.17(i) to bound the $\partial_\lambda^{|B|}m_g(i\tau,k)$ terms in the (possibly empty) $\pi\setminus\{B_0\}$,

$$\begin{aligned}\big|\partial_\lambda^n \widetilde{\mathscr{G}^r}(i\tau,k)\big| &\lesssim \sum_{\pi\in\Pi_n}\Bigg(\frac{1}{{c_{\mathrm{P}}}^{|\pi|+1}}\frac{\langle\hbar k\rangle^2|k|^{2-|B_0|}}{\langle\hbar k\rangle^2|k|^2+\tau^2}|\widehat{w}(k)|\|g\|_{H^{|B_0|+3/2+\delta_0}_{2+\lceil d/2\rceil}} \\ &\qquad\qquad \times \prod_{B\in\pi\setminus\{B_0\}} |\widehat{w}(k)||k|^{-|B|}\|g\|_{H^{|B|+3/2+\delta_0}_{\lceil d/2\rceil}}\Bigg) \\ &\leq \frac{\langle\hbar k\rangle^2|k|^{2-n}}{\langle\hbar k\rangle^2|k|^2+\tau^2}\sum_{\pi\in\Pi_n}\frac{|\widehat{w}(k)|^{|\pi|}}{{c_{\mathrm{P}}}^{|\pi|+1}}\prod_{B\in\pi}\|g\|_{H^{|B|+3/2+\delta_0}_{2+\lceil d/2\rceil}}.\end{aligned}$$

□

The decay of the Green function on the Fourier side is now obtained by taking the inverse Laplace transform.

Proposition 1.19 (Decay of $\widehat{\mathscr{G}^r}(t,k)$)**.** *Let $n \geq 0$ be an integer and assume $w \in L^1(\mathbb{R}^d)$ and $g \in H^{n+3/2+\delta_0}_{2+\lceil d/2\rceil}(\mathbb{R}^d)$ satisfy the uniform Penrose condition* (2) *with constant c_{P}. Then, there exists a constant $C = C(n,w,g,c_{\mathrm{P}},d,\delta_0)$ such that*

$$\big|\widehat{\mathscr{G}^r}(t,k)\big| \leq C\frac{\langle\hbar k\rangle|k|}{\langle kt\rangle^n}.$$

证明. By the uniform Penrose condition (2), the map $\lambda \mapsto \widetilde{\mathscr{G}^r}(\lambda,k)$ is analytic in $\operatorname{Re}\lambda > 0$ and extends continuously to $\operatorname{Re}\lambda = 0$. Moreover, Lemma 1.18 ensures sufficient decay in $\operatorname{Im}\lambda$, so the inverse Laplace contour can be deformed to the imaginary axis:

$$\widehat{\mathscr{G}^r}(t,k) = \frac{1}{2\pi i}\int_{\operatorname{Re}\lambda=0} e^{\lambda t}\widetilde{\mathscr{G}^r}(\lambda,k)\,d\lambda = \frac{1}{2\pi}\int_{\mathbb{R}} e^{i\tau t}\widetilde{\mathscr{G}^r}(i\tau,k)\,d\tau.$$

Assume first $|k|t \geq 1$. Integrating by parts n times and applying Lemma 1.18,

$$\begin{aligned}\left|\widehat{\mathscr{G}^r}(t,k)\right| = \frac{1}{2\pi t^n}\left|\int_{\mathbb{R}} \partial_\tau^n \left(e^{i\tau t}\right)\widetilde{\mathscr{G}^r}(i\tau,k)\,d\tau\right| &= \frac{1}{2\pi t^n}\left|\int_{\mathbb{R}} e^{i\tau t}\partial_\tau^n\widetilde{\mathscr{G}^r}(i\tau,k)\,d\tau\right| \\ &\lesssim \frac{1}{(|k|t)^n}\int_{\mathbb{R}} \frac{\langle\hbar k\rangle^2|k|^2}{\langle\hbar k\rangle^2|k|^2+\tau^2}\,d\tau \\ &= \pi\frac{\langle\hbar k\rangle|k|}{(|k|t)^n},\end{aligned}$$

where it was used that $\int_{\mathbb{R}} \frac{a^2}{a^2+x^2}\,dx = \pi a$ for $a > 0$. No boundary terms arise by Lemma 1.18. For $0 \leq |k|t \leq 1$, the estimate follows from Lemma 1.18 with $n = 0$, without integration by parts. □

2 Energy estimates on the density

This section initiates the nonlinear analysis by establishing the bootstrap improvements (I2) and (I4) on the density ρ_Q. The approach applies part of the linear theory from Section 1 to the nonlinear equation.

2.1 Linear damping in L_t^2

The first step is to derive an estimate for a general Volterra equation of the form

$$\phi(t,k) = H(t,k) - \int_0^t \widehat{\mathscr{L}}(t-s,k)\phi(s,k)\,ds, \quad \widehat{\mathscr{L}}(t,k) = \frac{2}{\hbar}\widehat{w}(k)\sin(\tfrac{\hbar}{2}t|k|^2)\widehat{g}\,(kt)\,. \tag{25}$$

Proposition 2.1. *Fix $T^* > 0$. Let $m \in \mathbb{R}$ and $\sigma \geq 0$. Suppose $w \in L^1(\mathbb{R}^d)$ and $g \in H^{\lceil\sigma\rceil+3/2+\delta_0}_{2+\lceil d/2\rceil}(\mathbb{R}^d)$ satisfy the uniform Penrose condition (2) with constant c_{P}. Assume*

$$\||k|^m\langle k,kt\rangle^\sigma H(t,k)\|_{L^2_t([0,T^*])} < \infty.$$

Then there exists a constant $C_{\mathrm{LD}} = C_{\mathrm{LD}}(\sigma,d,\delta_0,w,g,c_{\mathrm{P}})$ such that the solution $\phi(t,k)$ to the system (25) satisfies the pointwise-in-k estimate

$$\||k|^m\langle k,kt\rangle^\sigma\phi(t,k)\|_{L^2_t([0,T^*])} \leq C_{\mathrm{LD}}\||k|^m\langle k,kt\rangle^\sigma H(t,k)\|_{L^2_t([0,T^*])}.$$

证明. The result follows by taking Laplace transforms and applying Plancherel's theorem. The key point is to control the resolvent $\widetilde{\mathscr{G}}(\lambda,k)$ uniformly in k along the imaginary axis.

Taking the Laplace transform of (25) yields

$$\widetilde{\phi}(\lambda,k) = \widetilde{\mathscr{G}}(\lambda,k)\widetilde{H}(\lambda,k), \tag{26}$$

where $\widetilde{\mathscr{G}}$ is defined in (14). Assume first that σ is an integer. For any $0 \leq n \leq \sigma$,

$$|k|^n \partial_\lambda^n \widetilde{\phi}(\lambda, k) = \sum_{j \leq n} \binom{n}{j} |k|^j \partial_\lambda^j \widetilde{\mathscr{G}}(\lambda, k) |k|^{n-j} \partial_\lambda^{n-j} \widetilde{H}(\lambda, k)$$
$$= \widetilde{\mathscr{G}}(\lambda, k) |k|^n \partial_\lambda^n \widetilde{H}(\lambda, k) - \sum_{1 \leq j \leq n} \binom{n}{j} |k|^j \partial_\lambda^j \widetilde{\mathscr{G}^r}(\lambda, k) |k|^{n-j} \partial_\lambda^{n-j} \widetilde{H}(\lambda, k), \tag{27}$$

recalling that $\widetilde{\mathscr{G}}(\lambda, k) = 1 - \widetilde{\mathscr{G}^r}(\lambda, k)$. For the first term on the right-hand side, note that $\big|\widetilde{\mathscr{G}}(i\tau, k)\big| \leq {c_{\mathrm{P}}}^{-1}$, which follows from w and g satisfying the uniform Penrose condition (2). Along the imaginary axis $\lambda = i\tau$, Lemma 1.18 can be applied to the other terms in (27) to bound $|k|^j \big|\partial_\lambda^j \widetilde{\mathscr{G}^r}(i\tau, k)\big| \lesssim 1$. Therefore,

$$|k|^n \big|\partial_\lambda^n \widetilde{\phi}(i\tau, k)\big| \lesssim \sum_{j \leq n} |k|^j \big|\partial_\lambda^j \widetilde{H}(i\tau, k)\big|.$$

Taking the L^2 norm in τ and applying Plancherel's theorem yields

$$\| |kt|^n \phi(t, k) \|_{L_t^2([0, T^*])} \lesssim \| \langle kt \rangle^n H(t, k) \|_{L_t^2([0, T^*])}.$$

Moreover,

$$\langle k \rangle^\sigma \big|\widetilde{\phi}(i\tau, k)\big| = \langle k \rangle^\sigma \big|\widetilde{\mathscr{G}}(i\tau, k) \widetilde{H}(i\tau, k)\big| \lesssim {c_{\mathrm{P}}}^{-1} \langle k \rangle^\sigma \big|\widetilde{H}(i\tau, k)\big|.$$

Using $\langle k, kt \rangle^\sigma \sim \langle k \rangle^\sigma + |kt|^\sigma$, it follows that

$$\| \langle k, kt \rangle^\sigma \phi(t, k) \|_{L_t^2([0, T^*])} \lesssim \| \langle k, kt \rangle^\sigma H(t, k) \|_{L_t^2([0, T^*])}.$$

Multiplying the preceding estimate by $|k|^m$ yields the result for integer σ. For non-integer σ, the result follows by interpolation between the integer cases $\lfloor \sigma \rfloor$ and $\lceil \sigma \rceil$ [5]. □

2.2 Bootstrap improvement (I2)

To prove the bootstrap improvement (I2), the weighted norm is split into two contributions according to $\langle \hbar k \rangle^2 = 1 + \hbar^2 |k|^2$. Each contribution is treated separately. First, it is shown that under the bootstrap hypotheses (B1) and (B5), there exists a constant $K_2' > 0$ depending only on the fixed parameters in Proposition 1.12 and independent of $\hbar$, T^*, ε, Q_{in} and the bootstrap constants K_i, such that, for ε sufficiently small,

$$\big\| |k|^{1/2} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t, k) \big\|^2_{L_t^2([0, T^*], L_k^2)} \leq 2 K_2' \varepsilon^2. \tag{28}$$

Following this, it will be shown that there exists $K_2'' > 0$ such that, for ε sufficiently small,

$$\big\|\hbar|k|^{3/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|^2_{L^2_t([0,T^*],L^2_k)} \le 2K_2''\varepsilon^2. \tag{29}$$

Choosing $K_2 := K_2' + K_2''$ establishes the bootstrap improvement (I2).

2.2.1 Estimate for (28)

From (22), the density satisfies

$$\widehat{\rho_Q}(t,k) = H(t,k) - \int_0^t \widehat{\mathscr{L}}(t-s,k)\widehat{\rho_Q}(s,k)\,ds, \quad \widehat{\mathscr{L}}(t,k) = \frac{2}{\hbar}\widehat{w}(k)\sin(\tfrac{\hbar}{2}t|k|^2)\widehat{g}\,(kt)\,,$$

with

$$H(t,k) = \widehat{\rho_{Q_{\rm FS}}}(t,k) \\ - \frac{2\hbar^{-1}}{(2\pi)^d}\int_0^t\int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(s,\ell)\sin\big(\tfrac{\hbar}{2}\ell\cdot k(t-s)\big)\,\widehat{W[P]}(s,k-\ell,kt-\ell s)\,d\ell ds.$$

Under the assumptions of Theorem 1.3, Proposition 2.1 implies

$$\big\||k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|^2_{L^2_t([0,T^*],L^2_k)} \\ \le C_{\rm LD}\big(\big\||k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_{Q_{\rm FS}}}(t,k)\big\|^2_{L^2_t([0,T^*],L^2_k)} + \mathcal{N}^{(\mathrm{B2}),1}\big),$$

where

$$\mathcal{N}^{(\mathrm{B2}),1} = \int_{t=0}^{T^*}\int_k\bigg||k|^{1/2}\langle k,kt\rangle^{\sigma_4}\int_0^t\int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(s,\ell)\hbar^{-1} \\ \times\sin\big(\tfrac{\hbar}{2}\ell\cdot k(t-s)\big)\,\widehat{W[P]}\,(s,k-\ell,kt-\ell s)\;d\ell ds\bigg|^2 dkdt.$$

The initial data term is controlled by Lemma 1.11 and the assumption $\|Q_{\rm in}\|_{\mathcal{H}^{N_0}_{M,M}} \le \varepsilon$,

$$\big\||k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_{Q_{\rm FS}}}(t,k)\big\|^2_{L^2_t([0,T^*],L^2_k)} \lesssim \sum_{|\alpha|,|\beta|\le M}\big\|\boldsymbol{\xi}^\alpha\boldsymbol{x}^\beta Q_{\rm in}\big\|^2_{\mathcal{H}^{N_0}} \lesssim \varepsilon^2,$$

provided that $\sigma_4 < N_0 - (d+1)/2$. For the nonlinear term $\mathcal{N}^{(\mathrm{B2}),1}$, the triangle inequality $\langle k,kt\rangle^{\sigma_4} \lesssim \langle k-\ell,kt-\ell s\rangle^{\sigma_4} + \langle \ell,\ell s\rangle^{\sigma_4}$ gives the decomposition into *transport* and *reaction* parts $\mathcal{N}^{(\mathrm{B2}),1} \lesssim \mathcal{T}^{(\mathrm{B2}),1} + \mathcal{R}^{(\mathrm{B2}),1}$, where

$$\mathcal{T}^{(\mathrm{B2}),1} \lesssim \int_{t=0}^{T^*}\int_k\bigg[\int_0^t\int_\ell |k|^{1/2}\langle k-\ell,kt-\ell s\rangle^{\sigma_4}|\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)||\ell| \\ \times|k(t-s)|\big|\widehat{W[P]}\,(s,k-\ell,kt-\ell s)\big|\,d\ell ds\bigg]^2 dkdt,$$

$$\mathcal{R}^{(\mathrm{B2}),1} \lesssim \int_{t=0}^{T^*}\int_k\bigg[\int_0^t\int_\ell |k|^{1/2}\langle \ell,\ell s\rangle^{\sigma_4}|\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)||\ell| \\ \times|k(t-s)|\big|\widehat{W[P]}\,(s,k-\ell,kt-\ell s)\big|\,d\ell ds\bigg]^2 dkdt.$$

The regularity of the solution moves along the flow, allowing control of the transport term through the bootstrap assumption (B1). The reaction term, however, introduces the *time response kernel* and requires careful handling of resonances.

2.2.2 Control of transport term

For $\mathfrak{T}^{(\mathrm{B2}),1}$, since $k(t-s) = kt-\ell s-s(k-\ell)$, the factor $|k(t-s)|$ is bounded by a constant multiple of $\langle s(k-\ell), kt-\ell s\rangle$, and the weights are absorbed onto P using Definition 0.2. Writing $P_s := \langle s\boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_4} P(s)$ for $s \in [0, T^*]$,

$$\langle s(k-\ell), kt-\ell s\rangle\langle k-\ell, kt-\ell s\rangle^{\sigma_4}\left|\widehat{W[P]}\,(s, k-\ell, kt-\ell s)\right| = \left|\widehat{W[P_s]}\,(k-\ell, kt-\ell s)\right|.$$

This leads to

$$\mathfrak{T}^{(\mathrm{B2}),1} \lesssim \int_{t=0}^{T^*}\int_k \left[\int_0^t \int_\ell |k|^{1/2}|\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)||\ell|\left|\widehat{W[P_s]}\,(k-\ell, kt-\ell s)\right| d\ell ds\right]^2 dk dt. \tag{30}$$

Applying Cauchy–Schwarz in s and ℓ and the bound $|\widehat{w}(\ell)| \lesssim 1$,

$$\begin{aligned}\mathfrak{T}^{(\mathrm{B2}),1} \lesssim \int_{t=0}^{T^*}\int_k &\left[\int_{s=0}^t \int_\ell \langle s\rangle^{d-1/2}|\ell||\widehat{\rho_Q}(s,\ell)|\, d\ell ds\right] \\ &\times \left[\int_{s=0}^t \int_\ell \langle s\rangle^{-(d-1/2)}|\widehat{\rho_Q}(s,\ell)||\ell||k|\left|\widehat{W[P_s]}\,(k-\ell, kt-\ell s)\right|^2 d\ell ds\right] dk dt.\end{aligned}$$

Bootstrap assumption (B5) (via Lemma 1.14) implies

$$\int_{s=0}^t \langle s\rangle^{d-1/2}\int_\ell |\ell||\widehat{\rho_Q}(s,\ell)|\, d\ell ds \lesssim \sqrt{K_5}\varepsilon \int_0^\infty \frac{\langle s\rangle^{d-1/2}}{\langle s\rangle^{d+1}}\, ds \lesssim \sqrt{K_5}\varepsilon$$

provided that $\sigma_1 > d+1$, which allows control over the first s-ℓ integral. For the second, Fubini's theorem is applied:

$$\begin{aligned}\mathfrak{T}^{(\mathrm{B2}),1} \lesssim \sqrt{K_5}\varepsilon &\int_{s=0}^{T^*}\int_\ell \langle s\rangle^{d-1/2}|\ell||\widehat{\rho_Q}(s,\ell)| d\ell ds \\ &\times \sup_{s\in[0,T^*],\ell\in\mathbb{R}^d}\langle s\rangle^{-2d+1}\int_k\int_{t=-\infty}^\infty |k|\left|\widehat{W[P_s]}\,(k-\ell, kt-\ell s)\right|^2 dt dk.\end{aligned}$$

The s-ℓ integral is controlled again using (B5). Finally, the change of variable $t \mapsto t/|k|$ is made, and the L^2 trace lemma (Lemma 1.2) is applied to the

η-variable:

$$\begin{aligned}
\mathcal{T}^{(\mathrm{B2}),1} &\lesssim K_5\varepsilon^2 \sup_{s\in[0,T^*],\ell\in\mathbb{R}^d}\langle s\rangle^{-2d+1}\int_k\int_{t=-\infty}^{\infty}\left|\widehat{W[P_s]}\big(k-\ell,\tfrac{k}{|k|}t-\ell s\big)\right|^2 dtdk\\
&\lesssim K_5\varepsilon^2 \sup_{s\in[0,T^*]}\langle s\rangle^{-2d+1}\int_k\sup_{\omega\in\mathbb{S}^{d-1},x\in\mathbb{R}^d}\int_{t=-\infty}^{\infty}\left|\widehat{W[P_s]}\,(k,\omega t-x)\right|^2 dtdk\\
&\lesssim K_5\varepsilon^2 \sup_{s\in[0,T^*]}\langle s\rangle^{-2d+1}\int_k\left\|\langle\nabla_\eta\rangle^q\widehat{W[P_s]}\,(k,\eta)\right\|_{L^2_\eta}^2 dk\\
&\lesssim K_5\varepsilon^2 \sup_{s\in[0,T^*]}\langle s\rangle^{-2d+1}\sum_{|\alpha|\le M}\|v^\alpha W[P_s](z,v)\|_{L^2_{z,v}}^2,
\end{aligned}$$

where $(d-1)/2 < q \le M$. Using Definition 0.2 and Lemma 1.6, the resulting norm coincides with the quantity controlled by the bootstrap assumption (B1). Hence, it follows that the transport term satisfies

$$\mathcal{T}^{(\mathrm{B2}),1} \lesssim K_1K_5\varepsilon^4.$$

2.2.3 Control of resonances

For the reaction term $\mathcal{R}^{(\mathrm{B2}),1}$, the bound $|k(t-s)| \lesssim \langle s\rangle\langle k-\ell, kt-\ell s\rangle$ is utilised, and bootstrap assumption (B5) is used to control

$$\mathcal{R}^{(\mathrm{B2}),1} \lesssim K_5\varepsilon^2\int_{t=0}^{T^*}\int_k\left[\int_0^t\int_\ell \frac{\langle s\rangle|\widehat{w}(\ell)||k|^{1/2}|\ell|^{1/2}}{\langle k-\ell,kt-\ell s\rangle^{\sigma_1-1}}|\ell|^{1/2}\langle\ell,\ell s\rangle^{\sigma_4}|\widehat{\rho_Q}(s,\ell)|\,d\ell ds\right]^2 dkdt.$$

Consequently,

$$\mathcal{R}^{(\mathrm{B2}),1} \lesssim K_5\varepsilon^2\left\|\overline{R}\big[|k|^{1/2}\langle k,kt\rangle^{\sigma_4}|\widehat{\rho_Q}(t,k)|\big]\right\|^2_{L^2_t([0,T^*],L^2_k)} \tag{31}$$

with the integral operator $\overline{R}\colon L^2_t([0,T^*],L^2_k)\to L^2_t([0,T^*],L^2_k)$ defined by

$$\overline{R}[f](t,k) = \int_0^t\int_\ell \frac{\langle s\rangle|\widehat{w}(\ell)||k|^{1/2}|\ell|^{1/2}}{\langle k-\ell,kt-\ell s\rangle^{\sigma_1-1}}f(s,\ell)\,d\ell ds. \tag{32}$$

Through this kernel, density modes at earlier times $0\le s\le t$ can influence the k-th mode of the density at time t. Preventing a cascade of resonances is crucial: the ℓ-th mode of the density at time s may sequentially drive the k-th mode at time t, and so on, potentially causing growth in the nonlinear interaction at arbitrarily large times. Schur's test implies that

$$\begin{aligned}
&\left\|\overline{R}\right\|^2_{L^2_t([0,T^*],L^2_k)\to L^2_t([0,T^*],L^2_k)}\\
&\qquad\le\left(\sup_{t\in[0,T^*],k\in\mathbb{R}^d}\int_0^t\int_\ell\frac{\langle s\rangle|\widehat{w}(\ell)||k|^{1/2}|\ell|^{1/2}}{\langle k-\ell,kt-\ell s\rangle^{\sigma_1-1}}\,d\ell ds\right)\\
&\qquad\quad\times\left(\sup_{s\in[0,T^*],\ell\in\mathbb{R}^d}\int_s^{T^*}\int_k\frac{\langle s\rangle|\widehat{w}(\ell)||k|^{1/2}|\ell|^{1/2}}{\langle k-\ell,kt-\ell s\rangle^{\sigma_1-1}}\,dkdt\right).
\end{aligned}\tag{33}$$

The following lemma provides control over these integrals, analogous to [2, Lem. 3.1 & 3.2], with the proof extended to $d \geq 3$. The main challenge in managing the integrals in (33) arises from the $\langle s \rangle$-growth in the numerator, which becomes difficult to handle due to potential cancellations in $\langle k - \ell, kt - \ell s \rangle$ when k and ℓ are nearly collinear. To address this, a *resonant region* in $\mathbb{R}^d$ is identified, where k and ℓ are nearly collinear, and which diminishes over time. The integral over this resonant region converges as it shrinks at a sufficient rate to ensure convergence in the time variable. Outside the resonant region, where ℓ and k are sufficiently non-collinear, additional decay is extracted from the denominator through powers of $|kt - \ell s|$.

Lemma 2.2 (Control of resonances)**.** *Fix dimension $d \geq 3$ and let $T^* \geq 0$, $m \geq \max\{1, (d - 3/2)/2\}$ and $\beta \geq d + 6$. The following integral inequalities hold:*

$$\sup_{t \in [0,T^*], k \in \mathbb{R}^d} \int_0^t \int_{\ell \in \mathbb{R}^d} \frac{\langle s \rangle |k|^{1/2} |\ell|^{1/2}}{\langle \ell \rangle^m \langle k - \ell, kt - \ell s \rangle^\beta} d\ell ds \lesssim 1 \tag{34}$$

and

$$\sup_{s \in [0,T^*], \ell \in \mathbb{R}^d} \int_s^{T^*} \int_{k \in \mathbb{R}^d} \frac{\langle s \rangle |k|^{1/2} |\ell|^{1/2}}{\langle \ell \rangle^m \langle k - \ell, kt - \ell s \rangle^\beta} dk dt \lesssim 1. \tag{35}$$

Schur's test thus shows that the operator $\overline{R}$, defined in (32)*, satisfies*

$$\left\| \overline{R} \right\|_{L^2_t([0,T^*], L^2_k) \to L^2_t([0,T^*], L^2_k)} \lesssim 1.$$

证明. To establish inequality (34), fix $t \in [0, T^*]$ and $k \in \mathbb{R}^d$. If $k = 0$, the integral vanishes, so assume $k \neq 0$. The integral over early times is controlled as follows:

$$\int_0^1 \int_\ell \frac{\langle s \rangle |k|^{1/2} |\ell|^{1/2}}{\langle \ell \rangle^m \langle k - \ell, kt - \ell s \rangle^\beta} d\ell ds \lesssim \int_\ell \frac{|k|^{1/2} |\ell|^{1/2}}{\langle \ell \rangle^m \langle k - \ell \rangle^\beta} d\ell \lesssim \int_\ell \frac{|\ell|^{1/2}}{\langle \ell \rangle^{m - 1/2} \langle k - \ell \rangle^{\beta - 1/2}} d\ell \lesssim 1,$$

where $\langle k \rangle \lesssim \langle \ell \rangle \langle k - \ell \rangle$ was used.

Assume $T^* > 1$. Decompose $\ell \in \mathbb{R}^d$ into parallel and orthogonal components relative to k as $\ell_\parallel := (k \cdot \ell) k / |k|^2$ and $\ell_\perp := \ell - \ell_\parallel$. Define $I_R \subset \mathbb{R}^d$ as a cylinder around the line spanned by k with radius shrinking over time, and set $I_{NR} = I_R^c$, where

$$I_R = \left\{ \ell : |\ell_\perp| < |k|^b / \langle s \rangle^\zeta \right\} \quad \text{and} \quad I_{NR} = \left\{ \ell : |\ell_\perp| \geq |k|^b / \langle s \rangle^\zeta \right\},$$

with $b := (1 - \zeta)/2$, where ζ is chosen to satisfy

$$\frac{2}{d - 1/2} < \zeta < 1 \quad \text{and} \quad x := \frac{1}{1 - \zeta} < \frac{11}{2}.$$

Such a choice is possible since $(1-\zeta)^{-1} \to (2d-1)/(2d-5) \leq 5$ as $\zeta \downarrow 2/(d-1/2)$, for all $d \geq 3$.

The integral over the cylinder I_R satisfies, using $|\ell|^{1/2} \lesssim |\ell_\parallel|^{1/2} + |\ell_\perp|^{1/2}$,

$$\begin{aligned}\mathscr{I}_R &:= \int_1^t \int_{I_R} \frac{\langle s\rangle |k|^{1/2}|\ell|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt-\ell s\rangle^\beta}\, d\ell ds \\ &\lesssim \int_1^t \int_{I_R} \frac{\langle s\rangle |k|^{1/2}}{\langle \ell_\parallel\rangle^m \langle k-\ell_\parallel, kt-\ell_\parallel s\rangle^\beta} \big(|\ell_\parallel|^{1/2} + |\ell_\perp|^{1/2}\big)\, d\ell ds.\end{aligned}$$

Applying the bound for $|\ell_\perp|$ on I_R and the volume of its cross-section gives

$$\mathscr{I}_R \lesssim \int_1^t \int_{-\infty}^{\infty} \frac{\langle s\rangle |k|^{1/2}}{\langle \ell_\parallel\rangle^m \langle k-\ell_\parallel, kt-\ell_\parallel s\rangle^\beta} \frac{|k|^{(d-1)b}}{\langle s\rangle^{(d-1)\zeta}} \Big(|\ell_\parallel|^{1/2} + \frac{|k|^{b/2}}{\langle s\rangle^{\zeta/2}}\Big)\, d\ell_\parallel ds,$$

leading to two contributions $\mathscr{I}_1$ and $\mathscr{I}_2$. For $\mathscr{I}_1$, the change of variable $s \mapsto s/|\ell_\parallel|$ and $(d-1)\zeta \geq 1$ yield convergence under the conditions $\beta+m-2(d-1)b > 3/2$ and $\beta, m \geq 1/2+(d-1)b$. The second term $\mathscr{I}_2$ converges due to the choice of ζ, ensuring that $1/\langle s\rangle^{(d-1/2)\zeta-1}$ is integrable, for $\beta + m > 2 + 2(d-1/2)b$ and $m, \beta \geq 1/2 + (d-1/2)b$.

On I_{NR}, for $s \geq 1$, it holds that $|kt-\ell s| \gtrsim s|\ell_\perp| \gtrsim |k|^b \langle s\rangle^{1-\zeta}$, so

$$\begin{aligned}\mathscr{I}_{NR} &:= \int_1^t \int_{I_{NR}} \frac{\langle s\rangle |k|^{1/2}|\ell|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt-\ell s\rangle^\beta}\, d\ell ds \\ &\lesssim \int_1^t \int_{I_{NR}} \frac{\langle s\rangle |k|^{1/2}|\ell|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt-\ell s\rangle^{\beta - 1/(2b)} |k|^{1/2} \langle s\rangle^{(1-\zeta)/(2b)}}\, d\ell ds.\end{aligned}$$

Using $(1-\zeta)/(2b) - 1 = 0$, and applying the change of variable $s \mapsto s/|\ell|$, the integral $\mathscr{I}_{NR}$ converges for $\beta + m - 1/(2b) - 1/2 > d$.

To establish inequality (35), it is observed that:

$$\begin{aligned}\sup_{s\in[0,T^*],\ell\in\mathbb{R}^d} \int_s^{T^*} \int_{k\in\mathbb{R}^d} \frac{\langle s\rangle |k|^{1/2}|\ell|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt-\ell s\rangle^\beta}\, dkdt \\ \leq \sup_{s\in[0,T^*],\ell\in\mathbb{R}^d} \int_0^{T^*} \int_{k\in\mathbb{R}^d} \frac{\langle t\rangle |k|^{1/2}|\ell|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt-\ell s\rangle^\beta}\, dkdt.\end{aligned}$$

For fixed $s \in [0,T^*]$ and $\ell \in \mathbb{R}^d$, the integral over early times is estimated as

$$\int_0^1 \int_k \frac{\langle t\rangle |k|^{1/2}|\ell|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt-\ell s\rangle^\beta} dkdt \lesssim \int_k \frac{|k|^{1/2}|\ell|^{1/2}}{\langle \ell\rangle^m \langle k-\ell\rangle^\beta}\, dk \lesssim \int_k \frac{|k|^{1/2}}{\langle k\rangle^\beta}\, dk \lesssim 1.$$

Assume $T^* > 1$. If $\ell = 0$, the integral vanishes, so assume $\ell \neq 0$. Decompose $k \in \mathbb{R}^d$ into parallel and orthogonal components relative to ℓ, defining $k_\parallel := (\ell \cdot k)\ell/|\ell|^2$ and $k_\perp := k - k_\parallel$, and partition

$$J_R = \big\{k : |k_\perp| < |\ell|^b/\langle t\rangle^\zeta\big\} \quad \text{and} \quad J_{NR} = \big\{k : |k_\perp| \geq |\ell|^b/\langle t\rangle^\zeta\big\},$$

with ζ, b and x as in the proof of (34).

The integral over J_R satisfies

$$\begin{aligned}\mathscr{J}_R &:= \int_1^{T^*}\int_{J_R} \frac{\langle t\rangle |\ell|^{1/2}}{\langle \ell\rangle^m \langle k_\parallel - \ell, k_\parallel t - \ell s\rangle^\beta}\left(|k_\parallel|^{1/2} + |k_\perp|^{1/2}\right) dkdt \\ &\lesssim \int_1^{T^*}\int_{\mathbb{R}} \frac{|\ell|^{1/2}}{\langle \ell\rangle^m \langle k_\parallel - \ell, k_\parallel t - \ell s\rangle^\beta}\frac{|\ell|^{(d-1)b}}{\langle t\rangle^{(d-1)\zeta-1}}\left(|k_\parallel|^{1/2} + \frac{|\ell|^{b/2}}{\langle t\rangle^{\zeta/2}}\right) dk_\parallel dt,\end{aligned}$$

leading to two contributions, $\mathscr{J}_1$ and $\mathscr{J}_2$. Following the change of variable $t \mapsto t/|k_\parallel|$ in $\mathscr{J}_1$, the integrals converge under the conditions $m \geq (d-1/2)b + 1/2$ and $(d-1/2)\zeta - 1 > 1$.

On J_{NR}, for $t \geq 1$, it holds that $|kt - \ell s| \gtrsim t|k_\perp| \gtrsim |\ell|^b \langle t\rangle^{1-\zeta}$, so

$$\begin{aligned}\mathscr{J}_{NR} &:= \int_1^{T^*}\int_{J_{NR}} \frac{\langle t\rangle |\ell|^{1/2}|k|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt - \ell s\rangle^\beta}\, dkdt \\ &\lesssim \int_1^{T^*}\int_{J_{NR}} \frac{|\ell|^{1/2}|k|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt-\ell s\rangle^{\beta-x}|\ell|^{bx}\langle t\rangle^{x(1-\zeta)-1}}\, dkdt.\end{aligned}$$

Since $x(1-\zeta) = 1$ and $bx = 1/2$, the change of variable $t \mapsto t/|k|$ shows that the integral is uniformly bounded provided $\beta - x - 1/2 > d$. By the choice $x < 11/2$, this condition follows from $\beta \geq d+6$. The remaining conditions obtained above follow from $\beta \geq d+6$ and $m \geq \max\{1, (d-3/2)/2\}$. □

Applying Lemma 2.2 to (31) yields the bound

$$\mathcal{R}^{(\mathrm{B2}),1} \lesssim K_5\varepsilon^2 \left\| |k|^{1/2}\langle k, kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|^2_{L^2_t([0,T^*],L^2_k)}.$$

2.2.4 Conclusion of (28)

It has been shown that there exists a constant $K'_{\mathrm{LD}} > 0$, depending only on the fixed parameters in Proposition 1.12 and independent of $\hbar$, T^*, ε, Q_{in} and the bootstrap constants K_i, such that

$$\begin{aligned}&\left\| |k|^{1/2}\langle k, kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|^2_{L^2_t([0,T^*],L^2_k)} \\ &\qquad \leq K'_{\mathrm{LD}}\Big(\varepsilon^2 + K_1K_5\varepsilon^4 + K_5\varepsilon^2\left\| |k|^{1/2}\langle k, kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|^2_{L^2_t([0,T^*],L^2_k)}\Big).\end{aligned}$$

For ε sufficiently small so that $K'_{\mathrm{LD}}K_5\varepsilon^2 \leq 1/2$, the final term may be absorbed into the left-hand side, yielding

$$\left\| |k|^{1/2}\langle k, kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|^2_{L^2_t([0,T^*],L^2_k)} \leq 2K'_{\mathrm{LD}}\left(\varepsilon^2 + K_1K_5\varepsilon^4\right).$$

Choosing $K_1K_5\varepsilon^2 \leq 1$ and defining $K'_2 := 2K'_{\mathrm{LD}}$, the estimate (28) follows.

2.2.5 Estimate for (29)

Following the strategy of Section 2.2.1,

$$\big\||\hbar|k|^{3/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|^2_{L^2_t([0,T^*],L^2_k)} \\ \leq C_{\mathrm{LD}}\Big(\big\||\hbar|k|^{3/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_{Q_{\mathrm{FS}}}}(t,k)\big\|^2_{L^2_t([0,T^*],L^2_k)}+\mathcal{N}^{(\mathrm{B2}),2}\Big),$$

where

$$\mathcal{N}^{(\mathrm{B2}),2}=\int_{t=0}^{T^*}\int_k\bigg|\hbar|k|^{3/2}\langle k,kt\rangle^{\sigma_4}\int_0^t\int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(s,\ell)\hbar^{-1} \\ \times\sin\left(\tfrac{\hbar}{2}\ell\cdot k(t-s)\right)\widehat{W[P]}\left(s,k-\ell,kt-\ell s\right)\,d\ell ds\bigg|^2 dkdt.$$

The initial data term is controlled by Lemma 1.11 and the assumption on Q_{in}. For the nonlinear term, observe that the additional factor of $\hbar|k|$ in (29) cancels the $\hbar^{-1}$ denominator. The triangle inequality $\langle k,kt\rangle^{\sigma_4}\lesssim\langle k-\ell,kt-\ell s\rangle^{\sigma_4}+\langle \ell,\ell s\rangle^{\sigma_4}$ gives the decomposition $\mathcal{N}^{(\mathrm{B2}),2}\lesssim\mathcal{T}^{(\mathrm{B2}),2}+\mathcal{R}^{(\mathrm{B2}),2}$, where

$$\mathcal{T}^{(\mathrm{B2}),2}=\int_{t=0}^{T^*}\int_k\bigg[\int_0^t\int_\ell |k|^{3/2}\langle k-\ell,kt-\ell s\rangle^{\sigma_4}|\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)| \\ \times\left|\sin\left(\tfrac{\hbar}{2}\ell\cdot k(t-s)\right)\right|\left|\widehat{W[P]}\left(s,k-\ell,kt-\ell s\right)\right|d\ell ds\bigg]^2 dkdt,$$

$$\mathcal{R}^{(\mathrm{B2}),2}=\int_{t=0}^{T^*}\int_k\bigg[\int_0^t\int_\ell |k|^{3/2}\langle \ell,\ell s\rangle^{\sigma_4}|\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)| \\ \times\left|\sin\left(\tfrac{\hbar}{2}\ell\cdot k(t-s)\right)\right|\left|\widehat{W[P]}\left(s,k-\ell,kt-\ell s\right)\right|d\ell ds\bigg]^2 dkdt.$$

To control the transport part, the bound $\left|\sin\left(\frac{\hbar}{2}\ell\cdot k(t-s)\right)\right|\leq 1$ is used. The additional factor of $|k|$ in (29) is decomposed into $|k|\leq|\ell|+|k-\ell|$. This leads to two terms, with $\mathcal{T}^{(\mathrm{B2}),2}\lesssim\mathcal{T}_1^{(\mathrm{B2}),2}+\mathcal{T}_2^{(\mathrm{B2}),2}$ where

$$\mathcal{T}_1^{(\mathrm{B2}),2}=\int_{t=0}^{T^*}\int_k\bigg[\int_0^t\int_\ell |k|^{1/2}|\ell||\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)| \\ \times\left|\widehat{W[\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_4}P]}\left(s,k-\ell,kt-\ell s\right)\right|d\ell ds\bigg]^2 dkdt,$$

$$\mathcal{T}_2^{(\mathrm{B2}),2}=\int_{t=0}^{T^*}\int_k\bigg[\int_0^t\int_\ell |k|^{1/2}|\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)| \\ \times\left|\widehat{W[|\boldsymbol{\nabla}_x|\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_4}P]}\left(s,k-\ell,kt-\ell s\right)\right|d\ell ds\bigg]^2 dkdt.$$

The contribution $\mathcal{T}_1^{(\mathrm{B2}),2}$ is controlled by (30), and hence

$$\mathcal{T}_1^{(\mathrm{B2}),2}\lesssim K_1K_5\varepsilon^4.$$

For $\mathfrak{T}_2^{(\mathrm{B2}),2}$, observing that

$$\int_0^t \int_\ell \langle s\rangle^{d-3/2} |\widehat{\rho_Q}(s,\ell)|\, d\ell ds \lesssim \sqrt{K_5}\varepsilon \int_0^t \int_\ell \frac{\langle s\rangle^{d-3/2}}{\langle \ell, \ell s\rangle^{\sigma_1}}\, d\ell ds \lesssim \sqrt{K_5}\varepsilon$$

and proceeding as in Section 2.2.2 leads to

$$\mathfrak{T}_2^{(\mathrm{B2}),2} \lesssim K_5\varepsilon^2 \sup_{s\in[0,T^*]} \langle s\rangle^{-2d+3} \sum_{|\alpha|\le M} \left\| v^\alpha W \left[|\boldsymbol{\nabla}_x| \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_4} P\right](s,z,v)\right\|_{L^2_{z,v}}^2.$$

Using bootstrap assumption (B1),

$$\begin{aligned}
&\sup_{s\in[1,T^*]} \langle s\rangle^{-2d+3} \sum_{|\alpha|\le M} \left\| v^\alpha W \left[|\boldsymbol{\nabla}_x| \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_4} P\right](s,z,v)\right\|_{L^2_{z,v}}^2 \\
&\qquad \lesssim \sup_{s\in[1,T^*]} \langle s\rangle^{-2d+1} \sum_{|\alpha|\le M} \left\| v^\alpha W \left[|s\boldsymbol{\nabla}_x| \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_4} P\right](s,z,v)\right\|_{L^2_{z,v}}^2 \\
&\qquad \lesssim K_1\varepsilon^2,
\end{aligned}$$

while, for $s \in [0, \min\{1, T^*\}]$, the spatial-derivative estimate (39), which is derived from the bootstrap assumptions (B1), (B2), (B3) and (B5) alone, together with the choice of K_1 and the smallness condition $\varepsilon \le (\sqrt{K_2K_3} + \sqrt{K_1K_5})^{-1}$ made in Section 3.1, gives

$$\sup_{s\in[0,\min\{1,T^*\}]} \langle s\rangle^{-2d+3} \sum_{|\alpha|\le M} \left\| v^\alpha W \left[|\boldsymbol{\nabla}_x| \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_4} P\right](s,z,v)\right\|_{L^2_{z,v}}^2 \lesssim K_1\varepsilon^2.$$

Altogether, the transport term satisfies

$$\mathfrak{T}^{(\mathrm{B2}),2} \lesssim K_1K_5\varepsilon^4.$$

For the reaction term $\mathfrak{R}^{(\mathrm{B2}),2}$, bounding

$$\left|\sin\left(\tfrac{\hbar}{2}\ell\cdot k(t-s)\right)\right| \lesssim \hbar|\ell||k(t-s)|$$

and arguing as in Section 2.2.3 yields

$$\mathfrak{R}^{(\mathrm{B2}),2} \lesssim K_5\varepsilon^2 \int_0^{T^*}\int_k \left[\int_0^t \int_\ell \frac{\langle s\rangle |\widehat{w}(\ell)| \hbar|k|^{3/2} |\ell|^{1/2}}{\langle k-\ell, kt-\ell s\rangle^{\sigma_1-1}} |\ell|^{1/2} \langle \ell, \ell s\rangle^{\sigma_4} |\widehat{\rho_Q}(s,\ell)|\, d\ell ds\right]^2 dk dt.$$

Splitting the additional weight using the triangle inequality $\hbar|k| \le \hbar|\ell| + \hbar|k-\ell|$ yields two contributions: $\mathfrak{R}^{(\mathrm{B2}),2} \lesssim \mathfrak{R}_1^{(\mathrm{B2}),2} + \mathfrak{R}_2^{(\mathrm{B2}),2}$. The first contribution is exactly

$$\mathfrak{R}_1^{(\mathrm{B2}),2} = K_5\varepsilon^2 \left\| \overline{R}\left[\hbar|k|^{3/2} \langle k, kt\rangle^{\sigma_4} |\widehat{\rho_Q}(t,k)|\right]\right\|_{L^2_t([0,T^*],L^2_k)}^2,$$

where the integral operator $\overline{R}\colon L^2_t([0,T^*],L^2_k) \to L^2_t([0,T^*],L^2_k)$ is defined in (32). Lemma 2.2 implies $\left\|\overline{R}\right\|_{L^2_t([0,T^*],L^2_k)\to L^2_t([0,T^*],L^2_k)} \lesssim 1$, and hence

$$\mathfrak{R}_1^{(\mathrm{B2}),2} \lesssim K_5\varepsilon^2 \left\| \hbar|k|^{3/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k)\right\|_{L^2_t([0,T^*],L^2_k)}^2.$$

This term will be absorbed into the left-hand side. For the second contribution, use that $\hbar \in (0,1]$ and $|\widehat{w}(\ell)| \lesssim \langle \ell \rangle^{-m}$ with $m = M - 1/2$, by (16), and absorb $|k-\ell|$ into the denominator:

$$\begin{aligned}&\mathcal{R}_2^{(\mathrm{B2}),2}\\ &\lesssim K_5\varepsilon^2 \int_{t=0}^{T^*} \int_k \Big[\int_0^t \int_\ell \frac{\langle s\rangle |k|^{1/2} |\ell|^{1/2}}{\langle \ell\rangle^m \langle k-\ell, kt-\ell s\rangle^{\sigma_1 - 2}} |\ell|^{1/2} \langle \ell, \ell s\rangle^{\sigma_4} |\widehat{\rho_Q}(s,\ell)| \, d\ell ds \Big]^2 dk dt.\end{aligned}$$

By Lemma 2.2 and (28),

$$\mathcal{R}_2^{(\mathrm{B2}),2} \lesssim K_5\varepsilon^2 \big\| |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|^2_{L^2_t([0,T^*],L^2_k)} \lesssim K_2' K_5 \varepsilon^4,$$

provided that $\sigma_1 \geq d+8$.

2.2.6 Conclusion of (29)

There exists a constant $K''_{\mathrm{LD}} > 0$, depending only on the fixed parameters in Proposition 1.12 and independent of $\hbar$, T^*, ε, Q_{in} and the bootstrap constants K_i, such that

$$\begin{aligned}&\big\| \hbar |k|^{3/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|^2_{L^2_t([0,T^*],L^2_k)}\\ &\leq K''_{\mathrm{LD}} \Big(\varepsilon^2 + K_1 K_5 \varepsilon^4 + K_2' K_5 \varepsilon^4 + K_5 \varepsilon^2 \big\| \hbar |k|^{3/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|^2_{L^2_t([0,T^*],L^2_k)} \Big).\end{aligned}$$

Choosing ε sufficiently small such that $K''_{\mathrm{LD}} K_5 \varepsilon^2 \leq 1/2$ permits absorption into the left-hand side,

$$\big\| \hbar |k|^{3/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|^2_{L^2_t([0,T^*],L^2_k)} \leq 2K''_{\mathrm{LD}} \left(\varepsilon^2 + K_1 K_5 \varepsilon^4 + K_2' K_5 \varepsilon^4 \right).$$

Finally, for ε chosen such that $K_1 K_5 \varepsilon^2 \leq 1$ and $K_2' K_5 \varepsilon^2 \leq 1$, the improvement (29) follows upon setting $K_2'' := 3K''_{\mathrm{LD}}$. This completes the proof of the bootstrap improvement (I2).

2.3 Bootstrap improvement (I4)

This subsection establishes the improvement of bootstrap assumption (B4). The argument combines the linear damping result with a decomposition of the nonlinear term.

As in Section 2.2, the linear damping result (Proposition 2.1) is applied to the nonlinear equation, yielding

$$\big\| |k|^{1/2} \langle k, kt\rangle^{\sigma_2} \widehat{\rho_Q}(t,k) \big\|^2_{L^2_t([0,T^*])} \lesssim \big\| |k|^{1/2} \langle k, kt\rangle^{\sigma_2} \widehat{\rho_{Q_{\mathrm{FS}}}}(t,k) \big\|^2_{L^2_t([0,T^*])} + \mathcal{N}^{(\mathrm{B4})}(k),$$

where

$$\mathcal{N}^{(\mathrm{B4})}(k) = \int_{t=0}^{T^*} \Big| |k|^{1/2} \langle k, kt\rangle^{\sigma_2} \int_0^t \int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(s,\ell) \\ \times \hbar^{-1} \sin\left(\tfrac{\hbar}{2}\ell \cdot k(t-s)\right) \widehat{W[P]}\,(s, k-\ell, kt-\ell s)\; d\ell ds \Big|^2 dt.$$

The initial data term is controlled by Lemma 1.11, while the nonlinear contribution is decomposed into transport and reaction terms, with $\mathcal{N}^{(\mathrm{B4})} \lesssim \mathcal{T}^{(\mathrm{B4})}(k) + \mathcal{R}^{(\mathrm{B4})}(k)$, where

$$\mathcal{T}^{(\mathrm{B4})}(k) \lesssim \int_{t=0}^{T^*} \Big[\int_0^t \int_\ell |k|^{1/2} \langle k-\ell, kt-\ell s\rangle^{\sigma_2} |\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)| \\ \times |\ell||k(t-s)| \Big| \widehat{W[P]}\,(s, k-\ell, kt-\ell s) \Big|\, d\ell ds \Big]^2 dt$$

$$\mathcal{R}^{(\mathrm{B4})}(k) \lesssim \int_{t=0}^{T^*} \Big[\int_0^t \int_\ell |k|^{1/2} \langle \ell, \ell s\rangle^{\sigma_2} |\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)| \\ \times |\ell||k(t-s)| \Big| \widehat{W[P]}\,(s, k-\ell, kt-\ell s) \Big|\, d\ell ds \Big]^2 dt.$$

To control the transport term, apply the bounds $|\widehat{w}(\ell)| \lesssim 1$ and $|k(t-s)| \lesssim |k|t$. Then

$$\begin{aligned}
&\mathcal{T}^{(\mathrm{B4})}(k) \\
&\lesssim K_5 \varepsilon^2 |k|^3 \int_0^{T^*} t^2 \left[\int_0^t \int_\ell \frac{|\ell|}{\langle \ell, \ell s\rangle^{\sigma_1}} \langle k-\ell, kt-\ell s\rangle^{\sigma_2} \Big| \widehat{W[P]}\,(s, k-\ell, kt-\ell s) \Big|\, d\ell ds \right]^2 dt \\
&\lesssim K_5 \varepsilon^2 |k|^3 \int_0^{T^*} t^2 \Big[\int_0^t \left(\int_\ell \frac{|\ell|^2}{\langle \ell, \ell s\rangle^{2\sigma_1}} \frac{1}{|k-\ell|^{2\delta} \langle k-\ell, kt-\ell s\rangle^{2(\sigma_3-\sigma_2)}}\, d\ell \right)^{1/2} \\
&\qquad \times \left(\int_\ell |k-\ell|^{2\delta} \langle k-\ell, kt-\ell s\rangle^{2\sigma_3} \Big| \widehat{W[P]}\,(s, k-\ell, kt-\ell s) \Big|^2 d\ell \right)^{1/2} ds \Big]^2 dt,
\end{aligned}$$

where the second line uses Cauchy–Schwarz in ℓ. To estimate the first integral in ℓ, fix a parameter $j > 0$ such that $j \le \min\{\sigma_1, \sigma_3 - \sigma_2\}$. Using the inequality $\langle \ell, \ell s\rangle \langle k-\ell, kt-\ell s\rangle \gtrsim \langle k, kt\rangle$, it follows that

$$\frac{1}{\langle \ell, \ell s\rangle^{2\sigma_1} \langle k-\ell, kt-\ell s\rangle^{2(\sigma_3-\sigma_2)}} \lesssim \frac{1}{\langle \ell, \ell s\rangle^{2\sigma_1 - 2j} \langle k, kt\rangle^{2j}}.$$

To estimate the second integral in ℓ, the Sobolev embedding and Lemma 1.6 imply, for $d/2 < s \le \lceil (d+1)/2 \rceil$,

$$\begin{aligned}
&\int_\ell \sup_\eta \Big| \widehat{W[|\boldsymbol{\nabla}_x|^\delta \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_3} P]}(\ell, \eta) \Big|^2 d\ell \\
&\qquad \lesssim \int_\ell \int_\eta \Big| \langle \nabla_\eta \rangle^s \widehat{W[|\boldsymbol{\nabla}_x|^\delta \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_3} P]}(\ell, \eta) \Big|^2 d\eta d\ell \\
&\qquad \lesssim \sum_{|\alpha| \le M} \Big\| W\left[\boldsymbol{\xi}^\alpha |\boldsymbol{\nabla}_x|^\delta \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_3} P \right] \Big\|_{L^2_{z,v}}^2 \\
&\qquad \lesssim K_3 \varepsilon^2,
\end{aligned}$$

where the final step uses bootstrap assumption (B3). Substituting into the earlier bound yields

$$\begin{aligned}\mathcal{T}^{(\mathrm{B4})}(k) &\lesssim K_3K_5\varepsilon^4|k|^3\int_{t=0}^{T^*}\frac{t^2}{\langle k,kt\rangle^{2j}}\left[\int_{s=0}^{t}\left(\int_\ell \frac{|\ell|^2}{\langle \ell,\ell s\rangle^{2\sigma_1-2j}}\frac{1}{|k-\ell|^{2\delta}}\,d\ell\right)^{1/2}ds\right]^2 dt\\ &\lesssim K_3K_5\varepsilon^4\int_{t=0}^{|k|T^*}\frac{t^2}{\langle k,t\rangle^{2j}}dt\left[\int_{s=0}^{\infty}\frac{1}{\langle s\rangle^{d/2+1-\delta}}\,ds\right]^2\\ &\lesssim K_3K_5\varepsilon^4,\end{aligned}$$

provided that $\sigma_1 - j > d/2 + 1 - \delta$ and $j > 3/2$. These conditions are satisfied under the assumption $\sigma_3 - \sigma_2 > d/2$ and $\sigma_1 > d/2 + 5/2 - \delta$.

To control the reaction term, apply the bound $|\widehat{w}(\ell)| \lesssim 1$ and perform Cauchy–Schwarz in both ℓ and s:

$$\begin{aligned}\mathcal{R}^{(\mathrm{B4})}(k) &\lesssim \int_{t=0}^{T^*}|k|^3t^2\left(\int_{s=0}^{t}\int_\ell |\ell|\langle \ell,\ell s\rangle^{2\sigma_4}|\widehat{\rho_Q}(s,\ell)|^2\,d\ell ds\right)\\ &\qquad\times\left(\int_{s=0}^{t}\int_\ell |\ell|\langle \ell,\ell s\rangle^{-2(\sigma_4-\sigma_2)}\left|\widehat{W[P]}\,(s,k-\ell,kt-\ell s)\right|^2 d\ell ds\right)dt\\ &\lesssim K_2K_5\varepsilon^4\int_{t=0}^{T^*}|k|^3t^2\int_{s=0}^{t}\int_\ell \frac{|\ell|}{\langle \ell,\ell s\rangle^{2(\sigma_4-\sigma_2)}\langle k-\ell,kt-\ell s\rangle^{2\sigma_1}}\,d\ell ds dt.\end{aligned}$$

Fix a parameter $j > 0$ such that $j \le \min\{\sigma_1, \sigma_4 - \sigma_2\}$. Using the inequality

$$\langle \ell,\ell s\rangle\langle k-\ell,kt-\ell s\rangle \gtrsim \langle k,kt\rangle,$$

it follows that

$$\frac{1}{\langle \ell,\ell s\rangle^{2(\sigma_4-\sigma_2)}\langle k-\ell,kt-\ell s\rangle^{2\sigma_1}} \lesssim \frac{1}{\langle \ell,\ell s\rangle^{2(\sigma_4-\sigma_2-j)}\langle k,kt\rangle^{2j}},$$

and hence,

$$\begin{aligned}\mathcal{R}^{(\mathrm{B4})}(k) &\lesssim K_2K_5\varepsilon^4\int_{t=0}^{T^*}\frac{|k|^3t^2}{\langle k,kt\rangle^{2j}}\int_{s=0}^{t}\int_\ell\frac{|\ell|}{\langle \ell,\ell s\rangle^{2(\sigma_4-\sigma_2-j)}}\,d\ell ds dt\\ &\lesssim K_2K_5\varepsilon^4,\end{aligned}$$

where the ℓ-s integral converges for $j < \sigma_4 - \sigma_2 - (d+1)/2$, and the t integral converges for $j > 3/2$. For such a j to exist, the condition $\sigma_4 - \sigma_2 > d/2 + 2$ is required.

Conclusion of bootstrap improvement (I4)

Under the bootstrap assumptions (B2), (B3), and (B5), there exists a constant $\widetilde{K} > 0$, depending only on the fixed parameters in Proposition 1.12 and independent of $\hbar$, T^*, ε, Q_{in} and the bootstrap constants K_i, such that

$$\left\||k|^{1/2}\langle k,kt\rangle^{\sigma_2}\widehat{\rho_Q}(t,k)\right\|^2_{L^\infty_k L^2_t([0,T^*])} \le \widetilde{K}\left(\varepsilon^2 + (K_3K_5 + K_2K_5)\varepsilon^4\right).$$

Define $K_4 := \widetilde{K}$. Then for $\varepsilon^2 \le (K_3K_5 + K_2K_5)^{-1}$, the improvement (I4) in Proposition 1.12 follows.

3 Energy estimates on the density matrix

This section establishes the bootstrap improvements (I1), (I3) and (I5). The estimates are derived from energy bounds on P in weighted quantum Sobolev spaces, formulated via the Wigner transform. Throughout this section the Wigner transform of P is written $W[P](z,v)$, in the free-flow phase-space variables (z,v) introduced in Section 1.3.2.

The following lemma is used repeatedly in the analysis.

Lemma 3.1 ([2, Lem. 2.9])**.**

a. Let $f_1, f_2 \in L^2(\mathbb{R}^d_k \times \mathbb{R}^d_\eta)$ and $r \in L^1(\mathbb{R}^d)$. Then

$$\left| \int_\eta \int_k \int_\ell f_1(k,\eta) r(\ell) f_2(k-\ell, \eta - \ell t)\, d\ell dk d\eta \right| \lesssim \|f_1\|_{L^2_{k,\eta}} \|f_2\|_{L^2_{k,\eta}} \|r\|_{L^1_k}.$$

b. Let $f_1 \in L^2(\mathbb{R}^d_k \times \mathbb{R}^d_\eta)$ and $f_2 \in L^1(\mathbb{R}^d_k; L^2(\mathbb{R}^d_\eta))$ and $r \in L^2(\mathbb{R}^d)$. Then

$$\left| \int_\eta \int_k \int_\ell f_1(k,\eta) r(\ell) f_2(k-\ell, \eta - \ell t)\, d\ell dk d\eta \right| \lesssim \|f_1\|_{L^2_{k,\eta}} \|f_2\|_{L^1_k(L^2_\eta)} \|r\|_{L^2_k}.$$

证明. The proof of (a) relies on the Cauchy–Schwarz inequality, while the proof of (b) incorporates Young's convolution inequality. □

3.1 Bootstrap improvement (I1)

This subsection establishes the improvement of the bootstrap assumption (B1). The analysis is carried out on the Wigner transform, using the equivalence

$$\sum_{|\alpha|\le M} \|\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi \rangle^{\sigma_4} \langle t\boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi \rangle \, (\boldsymbol{\xi}^\alpha P(t))\|^2_{\mathcal{L}^2} = \sum_{|\alpha|\le M} \|\langle t\nabla_z, \nabla_v \rangle \, (v^\alpha W[P])\|^2_{H^{\sigma_4}}$$

as given by Lemma 1.6.

3.1.1 Estimate on the velocity derivative

Let α be a multi-index such that $|\alpha| \le M$. An energy estimate yields

$$(2\pi)^{2d} \frac{1}{2} \frac{d}{dt} \|\langle \nabla_v \rangle (v^\alpha W[P])\|^2_{H^{\sigma_4}} = \mathcal{L}^{(\mathrm{B1}),V} + \mathcal{N}^{(\mathrm{B1}),V},$$

where

$$\mathcal{L}^{(\mathrm{B1}),V} = \mathrm{Re}\int_{k,\eta} \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4}\overline{D_\eta^\alpha \widehat{W[P]}(t,k,\eta)}\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} D_\eta^\alpha \widehat{W[\mathrm{L}_P]}(t,k,\eta)\,d\eta dk,$$

$$\mathcal{N}^{(\mathrm{B1}),V} = \mathrm{Re}\int_{k,\eta} \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4}\overline{D_\eta^\alpha \widehat{W[P]}(t,k,\eta)}\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} D_\eta^\alpha \widehat{W[\mathrm{N}_P]}(t,k,\eta)\,d\eta dk.$$

The terms $\widehat{W[\mathrm{L}_P]}(t,k,\eta)$ and $\widehat{W[\mathrm{N}_P]}(t,k,\eta)$ are defined in (23).

To estimate $\mathcal{L}^{(\mathrm{B1}),V}$, Lemma 1.10 is applied to isolate the highest-order contribution, in which all D_η^α-derivatives fall on $\widehat{g}(\eta - kt)$, from the remainder terms. This gives the decomposition $\big|\mathcal{L}^{(\mathrm{B1}),V}\big| \lesssim \mathcal{L}_0^{(\mathrm{B1}),V} + \mathcal{L}_M^{(\mathrm{B1}),V}$, where

$$\mathcal{L}_0^{(\mathrm{B1}),V} = \int_{k,\eta} \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4}\big|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big|\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} \times |k||\widehat{w}(k)||\widehat{\rho_Q}(t,k)||\eta - kt|\big|D_\eta^\alpha \widehat{g}(\eta - kt)\big|\,d\eta dk,$$

$$\mathcal{L}_M^{(\mathrm{B1}),V} = \sum_{|\beta|\le|\alpha|-1}\int_{k,\eta} \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4}\big|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big|\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} \times \langle\hbar k\rangle^{|\alpha|-1}|k||\widehat{w}(k)||\widehat{\rho_Q}(t,k)|\big|D_\eta^\beta \widehat{g}(\eta - kt)\big|\,d\eta dk.$$

For $\mathcal{L}_0^{(\mathrm{B1}),V}$, observe that $\langle\eta\rangle \lesssim \langle kt\rangle\langle\eta - kt\rangle$ and $\langle k,\eta\rangle^{\sigma_4} \lesssim \langle k,kt\rangle^{\sigma_4}\langle\eta - kt\rangle^{\sigma_4}$. The decay assumption (16) on w implies $\langle kt\rangle|k||\widehat{w}(k)| \lesssim \langle t\rangle|k|^{1/2}$. Substituting these bounds gives

$$\begin{aligned}\big|\mathcal{L}_0^{(\mathrm{B1}),V}\big| &\lesssim \langle t\rangle\int_{k,\eta} \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4}\big|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big||k|^{1/2}\langle k,kt\rangle^{\sigma_4}|\widehat{\rho_Q}(t,k)| \\ &\qquad\times \langle\eta - kt\rangle^{\sigma_4+2}\big|D_\eta^\alpha \widehat{g}(\eta - kt)\big|\,d\eta dk \\ &\lesssim \langle t\rangle\big\|\langle\nabla_v\rangle W[P]\big\|_{H_M^{\sigma_4}}\big\||k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L_k^2}\|g\|_{H_M^{\sigma_4+2}}.\end{aligned}$$

To estimate $\mathcal{L}_M^{(\mathrm{B1}),V}$, the full decay assumption on the interaction kernel (16) is used to ensure $\langle k\rangle^{M-1/2}|\widehat{w}(k)| \lesssim 1$, which allows the bound

$$\langle kt\rangle\langle\hbar k\rangle^{|\alpha|-1}|k||\widehat{w}(k)| \lesssim \langle t\rangle\langle\hbar k\rangle|k|^{1/2}\langle k\rangle^{M-1/2}|\widehat{w}(k)| \lesssim \langle t\rangle\langle\hbar k\rangle|k|^{1/2},$$

valid uniformly for $\hbar \in (0,1]$. Applying Cauchy–Schwarz then gives

$$\mathcal{L}_M^{(\mathrm{B1}),V} \lesssim \langle t\rangle\|\langle\nabla_v\rangle W[P]\|_{H_M^{\sigma_4}}\big\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L_k^2}\|g\|_{H_M^{\sigma_4+1}}.$$

To control the nonlinear contribution $\mathcal{N}^{(\mathrm{B1}),V}$, Lemma 1.10 is used to isolate the highest-order term in which all D_η^α-derivatives fall on $\widehat{W[P]}$, yielding the decomposition $\mathcal{N}^{(\mathrm{B1}),V} = \mathcal{N}_0^{(\mathrm{B1}),V} + \mathcal{N}_M^{(\mathrm{B1}),V}$. The leading term will be treated explicitly below. The remainder is controlled using Lemma 1.10. Explicitly,

$$\begin{aligned}\mathcal{N}_0^{(\mathrm{B1}),V} = \mathrm{Im}\int_{k,\eta} &\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4}\overline{D_\eta^\alpha \widehat{W[P]}(t,k,\eta)}\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} \\ &\times\Bigg[\frac{\hbar^{-1}}{(2\pi)^d}\int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(t,\ell)\left(e^{-i(\hbar/2)\ell\cdot(\eta-kt)} - e^{i(\hbar/2)\ell\cdot(\eta-kt)}\right) \\ &\qquad\times D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t)\,d\ell\Bigg]dkd\eta\end{aligned}$$

and

$$\big|\mathcal{N}_M^{(\mathrm{B1}),V}\big| \lesssim \sum_{|\beta|\le|\alpha|-1} \int_{k,\eta,\ell} \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} \big|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big| \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} \\ \times |\widehat{w}(\ell)||\ell|\langle\hbar\ell\rangle^{|\alpha|-1} |\widehat{\rho_Q}(t,\ell)| \big|D_\eta^\beta \widehat{W[P]}(t,k-\ell,\eta-\ell t)\big| \, d\ell dk d\eta.$$

To estimate $\mathcal{N}_0^{(\mathrm{B1}),V}$, a cancellation is used by subtracting an integral that evaluates to zero:

$$\begin{aligned}
&\mathcal{N}_0^{(\mathrm{B1}),V} \\
&= \mathrm{Im} \int_{k,\eta,\ell} \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} \overline{D_\eta^\alpha \widehat{W[P]}(t,k,\eta)} \left(\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} - \langle\eta-\ell t\rangle\langle k-\ell,\eta-\ell t\rangle^{\sigma_4}\right) \\
&\qquad \times \left[\frac{\hbar^{-1}}{(2\pi)^d} \widehat{w}(\ell) \widehat{\rho_Q}(t,\ell) \left(e^{-i(\hbar/2)\ell\cdot(\eta-kt)} - e^{i(\hbar/2)\ell\cdot(\eta-kt)} \right) \right. \\
&\qquad\qquad \left. \times D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right] d\ell dk d\eta. \qquad (36)
\end{aligned}$$

The integral that has been subtracted is now shown to vanish. Observe that the Wigner transform of P satisfies the equation

$$\begin{cases}
\partial_t W[P](t,z,v) = W[\mathrm{L}_P](t,z,v) + W[\mathrm{N}_P](t,z,v) \\
W[\mathrm{L}_P](t,z,v) = -\dfrac{i\hbar^{-1}}{(2\pi)^d} \displaystyle\int_k e^{ik\cdot(z+vt)} \widehat{w}(k) \widehat{\rho_Q}(t,k) \big(g(v-\tfrac{\hbar}{2}k) - g(v+\tfrac{\hbar}{2}k)\big) dk, \\
W[\mathrm{N}_P](t,z,v) = -\dfrac{i\hbar^{-1}}{(2\pi)^d} \displaystyle\int_\ell e^{i\ell\cdot(z+vt)} \widehat{w}(\ell) \widehat{\rho_Q}(t,\ell) \\
\qquad\qquad \times \big(W[P](t,z+\tfrac{\hbar}{2}\ell t, v-\tfrac{\hbar}{2}\ell) - W[P](t,z-\tfrac{\hbar}{2}\ell t, v+\tfrac{\hbar}{2}\ell)\big)\, d\ell.
\end{cases} \qquad (37)$$

Using Plancherel's theorem, the subtracted integral corresponds to an expression on the physical side of the form:

$$\begin{aligned}
&\int_{k,\eta,\ell} \langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} \overline{D_\eta^\alpha \widehat{W[P]}(t,k,\eta)} \langle\eta-\ell t\rangle\langle k-\ell,\eta-\ell t\rangle^{\sigma_4} \\
&\times \left[\frac{\hbar^{-1}}{(2\pi)^d} \widehat{w}(\ell) \widehat{\rho_Q}(t,\ell) \left(e^{-i(\hbar/2)\ell\cdot(\eta-kt)} - e^{i(\hbar/2)\ell\cdot(\eta-kt)} \right) D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right] d\ell dk d\eta \\
&\quad = (2\pi)^d \int_{z,v} \overline{\langle\nabla_v\rangle\langle\nabla\rangle^{\sigma_4} \left((-iv)^\alpha W[P]\right)(t,z,v)} \hbar^{-1} \int_\ell e^{i\ell\cdot(z+vt)} \widehat{w}(\ell) \widehat{\rho_Q}(t,\ell) \\
&\qquad \times \Big[\left(\langle\nabla_v\rangle\langle\nabla\rangle^{\sigma_4} ((-iv)^\alpha W[P])\right)(t, z+\tfrac{\hbar}{2}\ell t, v-\tfrac{\hbar}{2}\ell) \\
&\qquad\qquad - \left(\langle\nabla_v\rangle\langle\nabla\rangle^{\sigma_4} ((-iv)^\alpha W[P])\right)(t, z-\tfrac{\hbar}{2}\ell t, v+\tfrac{\hbar}{2}\ell) \Big] \, d\ell dz dv,
\end{aligned}$$

where $\langle\nabla\rangle = \langle\nabla_z, \nabla_v\rangle$. To see that this integral has zero imaginary part, define

$$f(z,v) = \langle\nabla_v\rangle\langle\nabla_z,\nabla_v\rangle^{\sigma_4} \left((-iv)^\alpha W[P]\right)(t,z,v)$$

and write $f_\pm(z,v) := f(z \pm \frac{\hbar}{2}\ell t, v \mp \frac{\hbar}{2}\ell)$, so that the integral above equals $-(2\pi)^d I$, where

$$I := \hbar^{-1} \int_{z,v,\ell} \overline{f(z,v)} e^{i\ell\cdot(z+tv)} \widehat{\rho_Q}(t,\ell)\widehat{w}(\ell)\left[f_-(z,v) - f_+(z,v)\right] d\ell dz dv.$$

Apply the change of variables $z \mapsto z + \frac{\hbar}{2}\ell t$, $v \mapsto v - \frac{\hbar}{2}\ell$ in the first term and $z \mapsto z - \frac{\hbar}{2}\ell t$, $v \mapsto v + \frac{\hbar}{2}\ell$ in the second, followed by the substitution $\ell \mapsto -\ell$, which interchanges f_+ and f_-. Then

$$\begin{aligned} I &= \hbar^{-1} \int_{z,v,\ell} \overline{(f_+(z,v) - f_-(z,v))} e^{i\ell\cdot(z+tv)} \widehat{\rho_Q}(t,\ell)\widehat{w}(\ell) f(z,v)\, d\ell dv dz \\ &= \hbar^{-1} \int_{z,v,\ell} \overline{(f_-(z,v) - f_+(z,v))} e^{-i\ell\cdot(z+tv)} \widehat{\rho_Q}(t,-\ell)\widehat{w}(-\ell) f(z,v)\, d\ell dv dz \\ &= \overline{\hbar^{-1} \int_{z,v,\ell} \overline{f(z,v)} e^{i\ell\cdot(z+tv)} \widehat{\rho_Q}(t,\ell)\widehat{w}(\ell)\left[f_-(z,v) - f_+(z,v)\right] d\ell dz dv} \\ &= \overline{I}. \end{aligned}$$

The penultimate equality uses that w and $\rho_Q(t)$ are real-valued. Indeed, since $Q(t)$ is self-adjoint, $\rho_Q(t,x)$ is real-valued, and hence $\widehat{w}(-\ell) = \overline{\widehat{w}(\ell)}$ and $\widehat{\rho_Q}(t,-\ell) = \overline{\widehat{\rho_Q}(t,\ell)}$. Therefore, $I = \overline{I}$, so $\operatorname{Im} I = 0$.

Therefore, $\mathcal{N}_0^{(\mathrm{B1}),V}$ can be controlled using (36):

$$\begin{aligned} \left|\mathcal{N}_0^{(\mathrm{B1}),V}\right| \lesssim \int_{k,\eta,\ell} &\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} \left|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\right| \left|\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} - \langle\eta-\ell t\rangle\langle k-\ell,\eta-\ell t\rangle^{\sigma_4}\right| \\ &\times |\widehat{w}(\ell)||\ell||\widehat{\rho_Q}(t,\ell)||\eta - kt| \left|D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t)\right| d\ell dk d\eta. \end{aligned}$$

To estimate the difference of weights, write it as

$$\left(\langle\eta\rangle - \langle\eta-\ell t\rangle\right)\langle k,\eta\rangle^{\sigma_4} + \langle\eta-\ell t\rangle\left(\langle k,\eta\rangle^{\sigma_4} - \langle k-\ell,\eta-\ell t\rangle^{\sigma_4}\right).$$

The first bracket is bounded by $|\ell t| \le \langle \ell t\rangle$, by the reverse triangle inequality. By the Mean Value Theorem, the second is bounded by $\sigma_4 |(\ell,\ell t)| \langle k-\ell', \eta - \ell' t\rangle^{\sigma_4 - 1}$ for some $\ell' = s\ell$ with $s \in (0,1)$. Both $\langle k,\eta\rangle$ and $\langle k-\ell',\eta-\ell' t\rangle$ are at most $\langle \ell,\ell t\rangle + \langle k-\ell,\eta-\ell t\rangle$ by the triangle inequality, using $|\ell - \ell'| \le |\ell|$ for the latter. Hence

$$\begin{aligned} &|\langle\eta\rangle\langle k,\eta\rangle^{\sigma_4} - \langle\eta-\ell t\rangle\langle k-\ell,\eta-\ell t\rangle^{\sigma_4}| \\ &\qquad \lesssim \langle\ell t\rangle\langle\ell,\ell t\rangle^{\sigma_4} + \langle\eta-\ell t\rangle|(\ell,\ell t)|\langle\ell,\ell t\rangle^{\sigma_4-1} \\ &\qquad\quad + \langle\ell t\rangle\langle k-\ell,\eta-\ell t\rangle^{\sigma_4} + \langle\eta-\ell t\rangle|(\ell,\ell t)|\langle k-\ell,\eta-\ell t\rangle^{\sigma_4-1}. \end{aligned}$$

This leads to the decomposition, with $\mathcal{R}$ denoting the reaction contribution and $\mathcal{T}$ the transport contribution:

$$\left|\mathcal{N}_0^{(\mathrm{B1}),V}\right| \lesssim \mathcal{R}_{0,1}^{(\mathrm{B1}),V} + \mathcal{R}_{0,2}^{(\mathrm{B1}),V} + \mathcal{T}_{0,1}^{(\mathrm{B1}),V} + \mathcal{T}_{0,2}^{(\mathrm{B1}),V},$$

corresponding to the four terms above.

For $\mathcal{R}_{0,1}^{(\mathrm{B1}),V}$, note that $|\eta - kt| \lesssim \langle t\rangle\langle k-\ell, \eta-\ell t\rangle$ and $\langle \ell t\rangle|\ell||\widehat{w}(\ell)| \lesssim \langle t\rangle|\ell|^{1/2}$. These yield the bound

$$\mathcal{R}_{0,1}^{(\mathrm{B1}),V} \lesssim \langle t\rangle^2 \int_{k,\eta,\ell} \langle \eta\rangle\langle k,\eta\rangle^{\sigma_4} \big|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big| \langle \ell, \ell t\rangle^{\sigma_4}|\ell|^{1/2}|\widehat{\rho_Q}(t,\ell)| \\ \times \langle k-\ell, \eta-\ell t\rangle \big|D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t)\big|\, d\ell dk d\eta.$$

Lemma 3.1(b) then implies

$$\mathcal{R}_{0,1}^{(\mathrm{B1}),V} \lesssim \langle t\rangle^2 \|\langle \nabla_v\rangle W[P]\|_{H_M^{\sigma_4}} \big\||k|^{1/2}\langle k, kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L_k^2} \\ \times \int_k \big\|\langle k,\eta\rangle D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big\|_{L_\eta^2}\, dk.$$

The remaining integral is bounded using Cauchy–Schwarz in k:

$$\int_k \big\|\langle k,\eta\rangle D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big\|_{L_\eta^2}\, dk \lesssim \big\||k|^\delta\langle k\rangle^{d/2}\langle k,\eta\rangle D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big\|_{L_{k,\eta}^2} \lesssim \sqrt{K_3}\varepsilon.$$

The second reaction term $\mathcal{R}_{0,2}^{(\mathrm{B1}),V}$ is estimated similarly. It follows that

$$\mathcal{R}_{0,1}^{(\mathrm{B1}),V} + \mathcal{R}_{0,2}^{(\mathrm{B1}),V} \lesssim \sqrt{K_3}\varepsilon\langle t\rangle^2 \big\|\langle \nabla_v\rangle W[P]\big\|_{H_M^{\sigma_4}} \big\||k|^{1/2}\langle k, kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L_k^2}.$$

The transport term $\mathcal{T}_{0,1}^{(\mathrm{B1}),V}$ is estimated using the inequality

$$|\eta - kt| \lesssim \langle t(k-\ell), \eta - \ell t\rangle,$$

together with Lemma 3.1(a):

$$\begin{aligned}
\mathcal{T}_{0,1}^{(\mathrm{B1}),V} &\lesssim \int_{k,\eta,\ell} \langle \eta\rangle\langle k,\eta\rangle^{\sigma_4} \big|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big| \langle \ell t\rangle\langle k-\ell, \eta-\ell t\rangle^{\sigma_4} \\
&\qquad \times |\widehat{w}(\ell)||\ell||\widehat{\rho_Q}(t,\ell)||\eta - kt|\big|D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t)\big|\, d\ell dk d\eta \\
&\lesssim \|\langle \nabla_v\rangle W[P]\|_{H_M^{\sigma_4}} \|\langle t\nabla_z, \nabla_v\rangle W[P]\|_{H_M^{\sigma_4}} \int_\ell \langle \ell t\rangle|\ell||\widehat{\rho_Q}(t,\ell)||\widehat{w}(\ell)|\, d\ell \\
&\lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1}} \|\langle \nabla_v\rangle W[P]\|_{H_M^{\sigma_4}} \|\langle t\nabla_z, \nabla_v\rangle W[P]\|_{H_M^{\sigma_4}},
\end{aligned}$$

where the last step uses the decay assumption (16) and the bootstrap assumption (B5), provided that $\sigma_1 > d+2$. The second transport term $\mathcal{T}_{0,2}^{(\mathrm{B1}),V}$ is treated analogously.

For $\mathcal{N}_M^{(\mathrm{B1}),V}$, the weight $\langle \eta\rangle\langle k,\eta\rangle^{\sigma_4}$ is decomposed into transport and reaction components using

$$\langle \eta\rangle\langle k,\eta\rangle^{\sigma_4} \lesssim (\langle \ell t\rangle + \langle \eta - \ell t\rangle)\, \langle \ell, \ell t\rangle^{\sigma_4} + (\langle \ell t\rangle + \langle \eta - \ell t\rangle)\, \langle k-\ell, \eta - \ell t\rangle^{\sigma_4},$$

which leads to the bound

$$\left|\mathcal{N}_M^{(\mathrm{B1}),V}\right| \lesssim \mathcal{R}_{M,1}^{(\mathrm{B1}),V} + \mathcal{R}_{M,2}^{(\mathrm{B1}),V} + \mathcal{T}_{M,1}^{(\mathrm{B1}),V} + \mathcal{T}_{M,2}^{(\mathrm{B1}),V}.$$

To estimate the reaction terms, apply assumption (16) and proceed as in the estimate for $\mathcal{R}_{0,1}^{(\mathrm{B1}),V}$:

$$\mathcal{R}_{M,1}^{(\mathrm{B1}),V} + \mathcal{R}_{M,2}^{(\mathrm{B1}),V} \lesssim \sqrt{K_3}\varepsilon\langle t\rangle^2 \left\|\langle\nabla_v\rangle W[P]\right\|_{H_M^{\sigma_4}} \left\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|_{L_k^2}.$$

The transport terms $\mathcal{T}_{M,1}^{(\mathrm{B1}),V}$ and $\mathcal{T}_{M,2}^{(\mathrm{B1}),V}$ are treated analogously to $\mathcal{T}_{0,1}^{(\mathrm{B1}),V}$, yielding

$$\mathcal{T}_{M,1}^{(\mathrm{B1}),V} + \mathcal{T}_{M,2}^{(\mathrm{B1}),V} \lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1}} \|\langle\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}}^2.$$

3.1.2 Conclusion of the velocity derivative estimate

The preceding estimates imply

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\|\langle\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}}^2 \\
&\quad\lesssim \langle t\rangle \|\langle\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}} \left\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|_{L_k^2} \\
&\qquad + \sqrt{K_3}\varepsilon\langle t\rangle^2 \|\langle\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}} \left\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|_{L_k^2} \\
&\qquad + \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1}} \|\langle\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}} \|\langle t\nabla_z,\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}}.
\end{aligned}$$

Dividing through and estimating the derivative of the norm gives

$$\begin{aligned}
\frac{d}{dt}\|\langle\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}} &\lesssim \left(\langle t\rangle + \sqrt{K_3}\varepsilon\langle t\rangle^2\right) \left\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\right\|_{L_k^2} \\
&\quad + \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1}} \|\langle t\nabla_z,\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}}.
\end{aligned}$$

Integration in time, using the bootstrap assumptions (B1) and (B2), yields

$$\begin{aligned}
&\|\langle\nabla_v\rangle W[P](t)\|_{H_M^{\sigma_4}} - \|\langle\nabla_v\rangle W[Q_{\mathrm{in}}]\|_{H_M^{\sigma_4}} \\
&\quad\lesssim \int_0^t \left(\langle s\rangle + \sqrt{K_3}\varepsilon\langle s\rangle^2\right) \left\|\langle\hbar k\rangle|k|^{1/2}\langle k,ks\rangle^{\sigma_4}\widehat{\rho_Q}(s,k)\right\|_{L_k^2} ds \\
&\qquad + \int_0^t \frac{\sqrt{K_5}\varepsilon}{\langle s\rangle^{d+1}} \|\langle s\nabla_z,\nabla_v\rangle W[P](s)\|_{H_M^{\sigma_4}} ds \\
&\quad\lesssim \left(\int_0^t \left(\langle s\rangle + \sqrt{K_3}\varepsilon\langle s\rangle^2\right)^2 ds\right)^{1/2} \left\|\langle\hbar k\rangle|k|^{1/2}\langle k,ks\rangle^{\sigma_4}\widehat{\rho_Q}(s,k)\right\|_{L_t^2([0,T^*],L_k^2)} \\
&\qquad + \sqrt{K_1K_5}\varepsilon^2 \int_0^t \frac{\langle s\rangle^{d-1/2}}{\langle s\rangle^{d+1}} ds \\
&\quad\lesssim \sqrt{K_2}\varepsilon\left(1 + \sqrt{K_3}\varepsilon\langle t\rangle\right)\langle t\rangle^{3/2} + \sqrt{K_1K_5}\varepsilon^2.
\end{aligned}$$

Hence, there exists a constant $\widetilde{K}_V > 0$, depending only on the fixed parameters in Proposition 1.12 and independent of $\hbar$, T^*, ε, Q_{in} and the bootstrap constants K_i such that

$$\hbar^{-d} \sum_{|\alpha|\le M} \|\boldsymbol{\xi}^\alpha \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_4} \langle \boldsymbol{\nabla}_\xi\rangle P(t)\|_{L^2_{x,y}}^2 \\ \le \widetilde{K}_V \varepsilon^2 \left[\sqrt{K_2} + \varepsilon\big(\sqrt{K_2K_3} + \sqrt{K_1K_5}\big)\right]^2 \langle t\rangle^5. \quad (38)$$

3.1.3 Estimate on the spatial derivative

Let α be a multi-index with $|\alpha| \le M$. As in the velocity case, an energy estimate yields

$$(2\pi)^{2d}\frac{1}{2}\frac{d}{dt}\big\|\langle\nabla_z\rangle v^\alpha W[P]\big\|_{H^{\sigma_4}}^2 = \mathcal{L}^{(\mathrm{B1}),Z} + \mathcal{N}^{(\mathrm{B1}),Z},$$

where the linear and nonlinear terms are defined analogously. The linear contribution is controlled as in Section 3.1.1, giving

$$\mathcal{L}^{(\mathrm{B1}),Z} \lesssim \big\|\langle\nabla_z\rangle W[P]\big\|_{H^{\sigma_4}_M} \big\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L^2_k} \|g\|_{H^{\sigma_4+1}_M}.$$

For the nonlinear term, the decomposition $\mathcal{N}^{(\mathrm{B1}),Z} = \mathcal{N}_0^{(\mathrm{B1}),Z} + \mathcal{N}_M^{(\mathrm{B1}),Z}$ is used, following the same approach as in Section 3.1.1, with

$$\big|\mathcal{N}_0^{(\mathrm{B1}),Z}\big| \lesssim \int_{k,\eta,\ell} \langle k\rangle\langle k,\eta\rangle^{\sigma_4} \big|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big| |\langle k\rangle\langle k,\eta\rangle^{\sigma_4} - \langle k-\ell\rangle\langle k-\ell,\eta-\ell t\rangle^{\sigma_4}| \\ \times |\widehat{w}(\ell)||\ell||\widehat{\rho_Q}(t,\ell)||\eta-kt|\big|D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t)\big|\, d\ell dk d\eta,$$

$$\big|\mathcal{N}_M^{(\mathrm{B1}),Z}\big| \lesssim \sum_{|\beta|\le|\alpha|-1} \int_{k,\eta,\ell} \langle k\rangle\langle k,\eta\rangle^{\sigma_4} \big|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big| \langle k\rangle\langle k,\eta\rangle^{\sigma_4} \\ \times |\widehat{w}(\ell)||\ell|\langle\hbar\ell\rangle^{|\alpha|-1}|\widehat{\rho_Q}(t,\ell)|\big|D_\eta^\beta \widehat{W[P]}(t,k-\ell,\eta-\ell t)\big|\, d\ell dk d\eta.$$

To control $\mathcal{N}_0^{(\mathrm{B1}),Z}$, the Mean Value Theorem yields

$$|\langle k\rangle\langle k,\eta\rangle^{\sigma_4} - \langle k-\ell\rangle\langle k-\ell,\eta-\ell t\rangle^{\sigma_4}| \\ \lesssim \langle\ell\rangle\langle\ell,\ell t\rangle^{\sigma_4} + \langle k-\ell\rangle|(\ell,\ell t)|\langle\ell,\ell t\rangle^{\sigma_4-1} \\ + \langle\ell\rangle\langle k-\ell,\eta-\ell t\rangle^{\sigma_4} + \langle k-\ell\rangle|(\ell,\ell t)|\langle k-\ell,\eta-\ell t\rangle^{\sigma_4-1},$$

which splits the estimate into four contributions:

$$\big|\mathcal{N}_0^{(\mathrm{B1}),Z}\big| \lesssim \mathcal{R}_{0,1}^{(\mathrm{B1}),Z} + \mathcal{R}_{0,2}^{(\mathrm{B1}),Z} + \mathcal{T}_{0,1}^{(\mathrm{B1}),Z} + \mathcal{T}_{0,2}^{(\mathrm{B1}),Z}.$$

These terms are analogous to those in Section 3.1.1, and are treated using the same estimates, except that the leading-order reaction term $\mathcal{R}_{0,1}^{(\mathrm{B1}),Z}$ yields one

less power of $\langle t\rangle$. In conclusion, provided $\sigma_3 > d+2$, the estimate

$$\begin{aligned}\big|\mathcal{N}_0^{(\mathrm{B1}),Z}\big| \lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1}}\|\langle\nabla_z\rangle W[P]\|_{H_M^{\sigma_4}}\|\langle t\nabla_z,\nabla_v\rangle W[P]\|_{H_M^{\sigma_4}} \\ + \sqrt{K_3}\varepsilon\langle t\rangle\|\langle\nabla_z\rangle W[P]\|_{H_M^{\sigma_4}}\big\||k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L_k^2}\end{aligned}$$

follows. The same argument applies to $\mathcal{N}_M^{(\mathrm{B1}),Z}$.

3.1.4 Conclusion of the spatial derivative estimate

Proceeding as in Section 3.1.2, it follows that

$$\begin{aligned}&\frac{1}{2}\frac{d}{dt}\big\|\langle\nabla_z\rangle W[P](t)\big\|_{H_M^{\sigma_4}}^2 \\ &\quad\lesssim \big\|\langle\nabla_z\rangle W[P](t)\big\|_{H_M^{\sigma_4}}\big\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L_k^2} \\ &\qquad+\sqrt{K_3}\varepsilon\langle t\rangle\big\|\langle\nabla_z\rangle W[P](t)\big\|_{H_M^{\sigma_4}}\big\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L_k^2} \\ &\qquad+\frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1}}\big\|\langle\nabla_z\rangle W[P](t)\big\|_{H_M^{\sigma_4}}\big\|\langle t\nabla_z,\nabla_v\rangle W[P](t)\big\|_{H_M^{\sigma_4}}.\end{aligned}$$

This implies

$$\begin{aligned}\frac{d}{dt}\big\|\langle\nabla_z\rangle W[P](t)\big\|_{H_M^{\sigma_4}} &\lesssim \Big(1+\sqrt{K_3}\varepsilon\langle t\rangle\Big)\big\|\langle\hbar k\rangle|k|^{1/2}\langle k,kt\rangle^{\sigma_4}\widehat{\rho_Q}(t,k)\big\|_{L_k^2} \\ &\quad+\frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1}}\big\|\langle t\nabla_z,\nabla_v\rangle W[P](t)\big\|_{H_M^{\sigma_4}}.\end{aligned}$$

Integrating in time and applying the bootstrap assumptions (B1) and (B2) yields

$$\begin{aligned}&\big\|\langle\nabla_z\rangle W[P](t)\big\|_{H_M^{\sigma_4}} - \big\|\langle\nabla_z\rangle W[Q_{\mathrm{in}}]\big\|_{H_M^{\sigma_4}} \\ &\quad\lesssim \int_0^t\Big(1+\sqrt{K_3}\varepsilon\langle s\rangle\Big)\big\|\langle\hbar k\rangle|k|^{1/2}\langle k,ks\rangle^{\sigma_4}\widehat{\rho_Q}(s,k)\big\|_{L_k^2}\,ds \\ &\qquad+\int_0^t\frac{\sqrt{K_5}\varepsilon}{\langle s\rangle^{d+1}}\big\|\langle s\nabla_z,\nabla_v\rangle W[P](s)\big\|_{H_M^{\sigma_4}}\,ds \\ &\quad\lesssim \left(\int_0^t\Big(1+\sqrt{K_3}\varepsilon\langle s\rangle\Big)^2\,ds\right)^{1/2}\big\|\langle\hbar k\rangle|k|^{1/2}\langle k,ks\rangle^{\sigma_4}\widehat{\rho_Q}(s,k)\big\|_{L_t^2([0,T^*],L_k^2)} \\ &\qquad+\sqrt{K_1K_5}\varepsilon^2\int_0^t\frac{\langle s\rangle^{d-1/2}}{\langle s\rangle^{d+1}}\,ds \\ &\quad\lesssim \sqrt{K_2}\varepsilon\Big(1+\sqrt{K_3}\varepsilon\langle t\rangle\Big)\langle t\rangle^{1/2}+\sqrt{K_1K_5}\varepsilon^2.\end{aligned}$$

Hence, there exists a constant $\widetilde{K}_Z>0$, depending only on the fixed parameters in Proposition 1.12 and independent of $\hbar$, T^*, ε, Q_{in} and the bootstrap constants K_i, such that

$$\begin{aligned}\hbar^{-d}\sum_{|\alpha|\le M}\|\boldsymbol{\xi}^\alpha\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_4}\langle\boldsymbol{\nabla}_x\rangle P(t)\|_{L^2_{x,y}}^2 \\ \le \widetilde{K}_Z\varepsilon^2\Big[\sqrt{K_2}+\varepsilon\big(\sqrt{K_2K_3}+\sqrt{K_1K_5}\big)\Big]^2\langle t\rangle^3. \quad (39)\end{aligned}$$

3.1.5 Conclusion of bootstrap improvement (I1)

Choose

$$K_1 := 2\max\{\widetilde{K}_V, \widetilde{K}_Z\}\big(\sqrt{K_2}+1\big)^2$$

and impose $\varepsilon \le (\sqrt{K_2K_3}+\sqrt{K_1K_5})^{-1}$. Then, combining the bounds (38) and (39), it follows that

$$\hbar^{-d}\sum_{|\alpha|\le M}\|\boldsymbol{\xi}^\alpha\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_4}\langle\boldsymbol{\nabla}_\xi\rangle P(t)\|_{L^2_{x,y}}^2 \le \frac{1}{2}K_1\varepsilon^2\langle t\rangle^5$$

and

$$\hbar^{-d}\sum_{|\alpha|\le M}\|\boldsymbol{\xi}^\alpha\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_4}\langle\boldsymbol{\nabla}_x\rangle P(t)\|_{L^2_{x,y}}^2 \le \frac{1}{2}K_1\varepsilon^2\langle t\rangle^3.$$

Therefore, for $d\ge 3$, the bound

$$\hbar^{-d}\sum_{|\alpha|\le M}\|\boldsymbol{\xi}^\alpha\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_4}\langle t\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle P(t)\|_{L^2_{x,y}}^2 \le 2K_1\langle t\rangle^5\varepsilon^2 \le 2K_1\langle t\rangle^{2d-1}\varepsilon^2$$

holds, completing the proof of (I1).

3.2 Bootstrap improvement (I3)

This section establishes the bootstrap improvement (I3) by employing the equivalence

$$\sum_{|\alpha|\le M}\big\|\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_3}|\boldsymbol{\nabla}_x|^\delta\,(\boldsymbol{\xi}^\alpha P(t))\big\|_{\mathcal{L}^2}^2 = \sum_{|\alpha|\le M}\big\||\nabla_z|^\delta\langle\nabla_z,\nabla_v\rangle^{\sigma_3}\,(v^\alpha W[P])\big\|_{L^2}^2.$$

Fix a multi-index α with $|\alpha|\le M$. Differentiating one of the terms on the right-hand side yields

$$\begin{aligned}(2\pi)^{2d}\frac{1}{2}\frac{d}{dt}\big\||\nabla_z|^\delta\langle\nabla_z,\nabla_v\rangle^{\sigma_3}(v^\alpha W[P])\big\|_{L^2}^2 &\\ = \mathrm{Re}\int_{k,\eta}|k|^\delta\langle k,\eta\rangle^{\sigma_3}\overline{D_\eta^\alpha\widehat{W[P]}(t,k,\eta)}|k|^\delta\langle k,\eta\rangle^{\sigma_3}D_\eta^\alpha\widehat{W[\mathrm{L}_P]}(t,k,\eta)\,d\eta dk&\\ + \mathrm{Re}\int_{k,\eta}|k|^\delta\langle k,\eta\rangle^{\sigma_3}\overline{D_\eta^\alpha\widehat{W[P]}(t,k,\eta)}|k|^\delta\langle k,\eta\rangle^{\sigma_3}D_\eta^\alpha\widehat{W[\mathrm{N}_P]}(t,k,\eta)\,d\eta dk,&\end{aligned}$$

where $\widehat{W[\mathrm{L}_P]}(t,k,\eta)$ and $\widehat{W[\mathrm{N}_P]}(t,k,\eta)$ are defined in (23). Denote the linear and nonlinear contributions by $\mathcal{L}^{(\mathrm{B3})}$ and $\mathcal{N}^{(\mathrm{B3})}$, respectively.

To estimate $\mathcal{L}^{(\mathrm{B3})}$, Lemma 1.10 is applied to extract the leading-order contribution from the remainder terms. This yields $\big|\mathcal{L}^{(\mathrm{B3})}\big| \lesssim \mathcal{L}_0^{(\mathrm{B3})} + \mathcal{L}_M^{(\mathrm{B3})}$. For $\mathcal{L}_0^{(\mathrm{B3})}$, the inequality $\langle k,\eta\rangle^{\sigma_3}\lesssim\langle k,kt\rangle^{\sigma_3}\langle\eta-kt\rangle^{\sigma_3}$ is used, along with the estimate

$$\langle k,kt\rangle^{\sigma_3}|k|^{1+\delta} = |k|^{1/2}\langle k,kt\rangle^{\sigma_4}\frac{|k|^{1/2+\delta}}{\langle k,kt\rangle^{\sigma_4-\sigma_3}} \le \frac{|k|^{1/2}\langle k,kt\rangle^{\sigma_4}}{\langle t\rangle^{1/2+\delta}},$$

valid under the assumption $\sigma_4 - \sigma_3 \geq 1/2 + \delta$. Consequently,

$$\begin{aligned}\mathcal{L}_0^{(\mathrm{B3})} &\lesssim \frac{1}{\langle t\rangle^{1/2+\delta}} \int_{k,\eta} |k|^\delta \langle k,\eta\rangle^{\sigma_3} \big| D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big| |k|^{1/2} \langle k, kt\rangle^{\sigma_4} |\widehat{\rho_Q}(t,k)| \\ &\qquad\qquad \times \langle \eta - kt\rangle^{\sigma_3+1} \big| D_\eta^\alpha \widehat{g}(\eta - kt)\big|\, d\eta dk \\ &\lesssim \frac{1}{\langle t\rangle^{1/2+\delta}} \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P])\big\|_{L^2} \big\| |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k)\big\|_{L^2_k} \big\| g\big\|_{H^{\sigma_3+1}_M},\end{aligned}$$

where Cauchy–Schwarz is applied in η and then k. Similarly, under the assumption (16) on w,

$$\mathcal{L}_M^{(\mathrm{B3})} \lesssim \frac{1}{\langle t\rangle^{1/2+\delta}} \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P])\big\|_{L^2} \big\| \langle \hbar k\rangle |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k)\big\|_{L^2_k} \big\| g\big\|_{H^{\sigma_3}_M}.$$

To estimate the nonlinear term $\mathcal{N}^{(\mathrm{B3})}$, apply Lemma 1.10 to decompose it as $\mathcal{N}^{(\mathrm{B3})} = \mathcal{N}_0^{(\mathrm{B3})} + \mathcal{N}_M^{(\mathrm{B3})}$, where $\mathcal{N}_0^{(\mathrm{B3})}$ captures the contribution with the highest-order derivatives. As in Section 3.1.1, a cancellation is used to isolate a difference of weights. Specifically, subtracting an integral that evaluates to zero allows the replacement

$$|k|^\delta \langle k,\eta\rangle^{\sigma_3} \mapsto |k|^\delta \langle k,\eta\rangle^{\sigma_3} - |k-\ell|^\delta \langle k-\ell, \eta - \ell t\rangle^{\sigma_3},$$

which introduces a commutator structure and leads to additional decay. The resulting term satisfies

$$\begin{aligned}\big|\mathcal{N}_0^{(\mathrm{B3})}\big| \lesssim \int_{k,\eta,\ell} &|k|^\delta \langle k,\eta\rangle^{\sigma_3} \big| D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big| \big| |k|^\delta \langle k,\eta\rangle^{\sigma_3} - |k-\ell|^\delta \langle k-\ell,\eta-\ell t\rangle^{\sigma_3}\big| \\ &\times |\widehat{w}(\ell)||\ell| |\widehat{\rho_Q}(t,\ell)| |\eta - kt| \big| D_\eta^\alpha \widehat{W[P]}(t, k-\ell, \eta - \ell t)\big|\, d\ell dk d\eta,\end{aligned}$$

while the lower-order term satisfies

$$\begin{aligned}\big|\mathcal{N}_M^{(\mathrm{B3})}\big| \lesssim \sum_{|\beta|\leq|\alpha|-1} \int_{k,\eta,\ell} &|k|^\delta \langle k,\eta\rangle^{\sigma_3} \big| D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\big| |k|^\delta \langle k,\eta\rangle^{\sigma_3} \\ &\times |\widehat{w}(\ell)||\ell| \langle \hbar\ell\rangle^{|\alpha|-1} |\widehat{\rho_Q}(t,\ell)| \big| D_\eta^\beta \widehat{W[P]}(t, k-\ell, \eta-\ell t)\big|\, d\ell dk d\eta.\end{aligned}$$

To estimate $\big|\mathcal{N}_0^{(\mathrm{B3})}\big|$, apply the Mean Value Theorem to the difference of weights:

$$\begin{aligned}&\big| |k|^\delta \langle k,\eta\rangle^{\sigma_3} - |k-\ell|^\delta \langle k-\ell,\eta-\ell t\rangle^{\sigma_3}\big| \\ &\leq \big| |k|^\delta - |k-\ell|^\delta\big| \langle k,\eta\rangle^{\sigma_3} + |k-\ell|^\delta |\langle k,\eta\rangle^{\sigma_3} - \langle k-\ell,\eta-\ell t\rangle^{\sigma_3}| \\ &\lesssim |\ell|^\delta \langle k,\eta\rangle^{\sigma_3} + |k-\ell|^\delta |(\ell,\ell t)| \sup_{|\ell'|\leq|\ell|} \langle k-\ell', \eta - \ell' t\rangle^{\sigma_3 - 1} \\ &\lesssim (|\ell|^\delta + |k-\ell|^\delta) \langle \ell, \ell t\rangle^{\sigma_3} + |\ell|^\delta \langle k-\ell, \eta - \ell t\rangle^{\sigma_3} + |k-\ell|^\delta |(\ell,\ell t)| \langle k-\ell, \eta-\ell t\rangle^{\sigma_3 - 1}.\end{aligned}$$

This gives three contributions

$$\left|\mathcal{N}_0^{(\mathrm{B3})}\right| \lesssim \mathcal{R}_0^{(\mathrm{B3})} + \mathcal{T}_{0,1}^{(\mathrm{B3})} + \mathcal{T}_{0,2}^{(\mathrm{B3})}.$$

For the reaction term, note that $|\eta - kt| \le |\eta - \ell t| + t|k-\ell|$, which motivates decomposing $\mathcal{R}_0^{(\mathrm{B3})}$ into four parts according to whether the prefactor contains $|\ell|^\delta$ or $|k-\ell|^\delta$, and whether $|\eta - kt|$ is controlled by $|\eta - \ell t|$ or $t|k-\ell|$:

$$\begin{aligned}
\mathcal{R}_0^{(\mathrm{B3})} &\lesssim \int_{k,\eta,\ell} |k|^\delta \langle k,\eta\rangle^{\sigma_3} \left|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\right| |\ell|^\delta \langle \ell, \ell t\rangle^{\sigma_3} \\
&\qquad \times |\widehat{w}(\ell)||\ell| |\widehat{\rho_Q}(t,\ell)| \left(|\eta-\ell t| + t|k-\ell|\right) \left|D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t)\right| d\ell dk d\eta \\
&\quad + \int_{k,\eta,\ell} |k|^\delta \langle k,\eta\rangle^{\sigma_3} \left|D_\eta^\alpha \widehat{W[P]}(t,k,\eta)\right| |k-\ell|^\delta \langle \ell, \ell t\rangle^{\sigma_3} \\
&\qquad \times |\widehat{w}(\ell)||\ell| |\widehat{\rho_Q}(t,\ell)| \left(|\eta-\ell t| + t|k-\ell|\right) \left|D_\eta^\alpha \widehat{W[P]}(t,k-\ell,\eta-\ell t)\right| d\ell dk d\eta \\
&= \mathcal{R}_{1;V}^{(\mathrm{B3})} + \mathcal{R}_{1;Z}^{(\mathrm{B3})} + \mathcal{R}_{2;V}^{(\mathrm{B3})} + \mathcal{R}_{2;Z}^{(\mathrm{B3})},
\end{aligned}$$

where the subscripts V and Z indicate association with the velocity variable v (terms involving $|\eta - \ell t|$) and the spatial variable z (terms involving $t|k-\ell|$), respectively. These four terms are treated separately, beginning with $\mathcal{R}_{1;V}^{(\mathrm{B3})}$, which is the most delicate: it alone lacks a positive power of $|k-\ell|$ in the integrand, so it cannot be controlled directly through the (B3) norm, and (B2) is used in its place. The starting point is the algebraic bound

$$|\ell|^{1+\delta}\langle \ell, \ell t\rangle^{\sigma_3} = |\ell|^{\frac{1}{2}} \langle \ell, \ell t\rangle^{\sigma_4} \frac{|\ell|^{1/2+\delta}}{\langle \ell, \ell t\rangle^{\sigma_4 - \sigma_3}} \le |\ell|^{\frac{1}{2}} \langle \ell, \ell t\rangle^{\sigma_4} \frac{|\ell|^{1/2+\delta}}{|(\ell, \ell t)|^{1/2+\delta}} = \frac{|\ell|^{1/2}\langle \ell, \ell t\rangle^{\sigma_4}}{\langle t\rangle^{1/2+\delta}},$$

valid provided $\sigma_4 - \sigma_3 \ge 1/2 + \delta$, which replaces the density weight $|\ell|^{1+\delta}\langle \ell, \ell t\rangle^{\sigma_3}$ by the weight $|\ell|^{1/2}\langle \ell, \ell t\rangle^{\sigma_4}$ governed by (B2), and extracts the decay factor $\langle t\rangle^{-(1/2+\delta)}$. Applying Lemma 3.1(b) with the $L_k^1 L_\eta^2$ norm applied to $\eta D_\eta^\alpha \widehat{W}[P](t,k,\eta)$, and noting that the weight $|k|^{-2\delta}\langle k\rangle^{-2(\sigma_3 - 1)}$ is integrable when $d \ge 3$, then yields

$$\begin{aligned}
\mathcal{R}_{1;V}^{(\mathrm{B3})} &\lesssim \frac{1}{\langle t\rangle^{1/2+\delta}} \left\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \right\|_{L^2} \left\| |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L_k^2} \\
&\qquad\qquad \times \int_k \left\| \langle \eta\rangle D_\eta^\alpha \widehat{W}[P](t,k,\eta) \right\|_{L_\eta^2} dk \\
&\lesssim \frac{1}{\langle t\rangle^{1/2+\delta}} \left\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \right\|_{L^2}^2 \left\| |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L_k^2}.
\end{aligned}$$

Each of the remaining terms $\mathcal{R}_{1;Z}^{(\mathrm{B3})}$, $\mathcal{R}_{2;V}^{(\mathrm{B3})}$, $\mathcal{R}_{2;Z}^{(\mathrm{B3})}$ contains a positive power of $|k-\ell|$ in its integrand, allowing the use of Lemma 3.1(a) with the L^1 term applied to the density, incorporating the appropriate weights. For $\mathcal{R}_{1;Z}^{(\mathrm{B3})}$, observe

that $|k-\ell| \le |k-\ell|^\delta \langle k-\ell, \eta-\ell t\rangle^{\sigma_3}$, so

$$\begin{aligned}
\mathcal{R}_{1;Z}^{(\mathrm{B3})} &\lesssim t \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2}^2 \int_\ell |\ell|^{1+\delta} |\widehat{w}(\ell)| \langle \ell, \ell t\rangle^{\sigma_3} |\widehat{\rho_Q}(t,\ell)| \, d\ell \\
&\lesssim t \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2}^2 \big\| |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|_{L_k^2} \\
&\qquad\qquad \times \left(\int_\ell \frac{|\ell|^{1+2\delta}}{\langle \ell, \ell t\rangle^{2\sigma_4 - 2\sigma_3}} \, d\ell \right)^{1/2} \\
&\lesssim \frac{1}{\langle t\rangle^{d/2-1/2+\delta}} \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2}^2 \big\| |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|_{L_k^2}
\end{aligned}$$

provided that $\sigma_4 - \sigma_3 > d/2 + 1/2 + \delta$. The remaining terms $\mathcal{R}_{2;V}^{(\mathrm{B3})}$, $\mathcal{R}_{2;Z}^{(\mathrm{B3})}$ can be estimated in the same fashion, leading to

$$\mathcal{R}_{2;V}^{(\mathrm{B3})} + \mathcal{R}_{2;Z}^{(\mathrm{B3})} \lesssim \frac{1}{\langle t\rangle} \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2}^2 \big\| |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|_{L_k^2}.$$

Next, the transport terms $\mathcal{T}_{0,1}^{(\mathrm{B3})}$ and $\mathcal{T}_{0,2}^{(\mathrm{B3})}$ are addressed. For $\mathcal{T}_{0,1}^{(\mathrm{B3})}$, the additional factor of $|\eta - kt|$ necessitates the use of the bootstrap estimate (B1), which grows like $\langle t\rangle^{d-1/2}$. Since no time growth is allowed in (I3), decay must instead be extracted from $|\widehat{\rho_Q}(t,\ell)| \lesssim \sqrt{K_5}\varepsilon \langle \ell, \ell t\rangle^{-\sigma_1}$ to control this term. Using the bound $|\eta - kt| \lesssim \langle t(k-\ell), \eta - \ell t\rangle$, the term can be estimated as follows:

$$\begin{aligned}
\mathcal{T}_{0,1}^{(\mathrm{B3})} &\lesssim \int_{k,\eta,\ell} |k|^\delta \langle k, \eta\rangle^{\sigma_3} \Big| D_\eta^\alpha \widehat{W[P]}(t,k,\eta) \Big| |\ell|^{1+\delta} |\widehat{\rho_Q}(t,\ell)| \\
&\qquad \times \langle t(k-\ell), \eta - \ell t\rangle \langle k-\ell, \eta - \ell t\rangle^{\sigma_3} \big| D_\eta^\alpha \widehat{W[P]}(t, k-\ell, \eta - \ell t) \big| \, d\ell dk d\eta \\
&\lesssim \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2} \big\| \langle t\nabla_z, \nabla_v\rangle W[P] \big\|_{H_M^{\sigma_4}} \int_\ell |\ell|^{1+\delta} |\widehat{\rho_Q}(t,\ell)| \, d\ell \\
&\lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1+\delta}} \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2} \big\| \langle t\nabla_z, \nabla_v\rangle W[P] \big\|_{H_M^{\sigma_4}}.
\end{aligned}$$

For $\mathcal{T}_{0,2}^{(\mathrm{B3})}$, the additional flexibility allows for the expression $|\eta - kt| \lesssim \langle t\rangle \langle k - \ell, \eta - \ell t\rangle$ to be used, leading to control via the (B3) norm:

$$\begin{aligned}
\mathcal{T}_{0,2}^{(\mathrm{B3})} &\lesssim \langle t\rangle \int_{k,\eta,\ell} |k|^\delta \langle k, \eta\rangle^{\sigma_3} \big| D_\eta^\alpha \widehat{W[P]}(t,k,\eta) \big| |\ell| |(\ell, \ell t)| |\widehat{\rho_Q}(t,\ell)| \\
&\qquad \times \langle k-\ell, \eta - \ell t\rangle^{\sigma_3} |k-\ell|^\delta \big| D_\eta^\alpha \widehat{W[P]}(t, k-\ell, \eta - \ell t) \big| \, d\ell dk d\eta \\
&\lesssim \langle t\rangle \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2}^2 \int_\ell |\ell| |(\ell, \ell t)| |\widehat{\rho_Q}(t,\ell)| \, d\ell \\
&\lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^d} \big\| |\nabla_z|^\delta \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2}^2.
\end{aligned}$$

This estimate is valid provided that $\sigma_1 > d + 2$.

For the remainder term $\mathcal{N}_M^{(\mathrm{B3})}$, the triangle inequality yields $|k|^\delta \le |\ell|^\delta + |k-\ell|^\delta$ and $\langle k, \eta\rangle^{\sigma_3} \lesssim \langle \ell, \ell t\rangle^{\sigma_3} + \langle k-\ell, \eta - \ell t\rangle^{\sigma_3}$. This gives the bound

$$|k|^\delta \langle k, \eta\rangle^{\sigma_3} \lesssim (|\ell|^\delta + |k-\ell|^\delta) \langle \ell, \ell t\rangle^{\sigma_3} + |\ell|^\delta \langle k-\ell, \eta - \ell t\rangle^{\sigma_3} + |k-\ell|^\delta \langle k-\ell, \eta - \ell t\rangle^{\sigma_3}.$$

Consequently, three terms are obtained such that $\big|\mathcal{N}_M^{(\mathrm{B3})}\big| \lesssim \mathcal{R}_M^{(\mathrm{B3})} + \mathcal{T}_{M,1}^{(\mathrm{B3})} + \mathcal{T}_{M,2}^{(\mathrm{B3})}$. The reaction term is controlled in the same way as the leading-order contribution $\mathcal{R}_0^{(\mathrm{B3})}$ above:

$$\mathcal{R}_M^{(\mathrm{B3})} \lesssim \frac{1}{\langle t\rangle^{1/2+\delta}} \big\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2} \\ \times \big\| \langle \hbar k\rangle |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|_{L^2_k} \big\| |\nabla_z|^{\delta} W[P] \big\|_{H^{\sigma_3}_M}.$$

The transport term $\mathcal{T}_{M,1}^{(\mathrm{B3})}$ is treated similarly to $\mathcal{T}_{0,1}^{(\mathrm{B3})}$, leading to

$$\mathcal{T}_{M,1}^{(\mathrm{B3})} \lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1+\delta}} \big\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2} \big\| \langle t\nabla_z, \nabla_v\rangle W[P] \big\|_{H^{\sigma_4}_M}.$$

Finally, $\mathcal{T}_{M,2}^{(\mathrm{B3})}$ is handled as in $\mathcal{T}_{0,2}^{(\mathrm{B3})}$:

$$\mathcal{T}_{M,2}^{(\mathrm{B3})} \lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1}} \big\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P]) \big\|_{L^2} \big\| |\nabla_z|^{\delta} W[P] \big\|_{H^{\sigma_3}_M}.$$

Conclusion of bootstrap improvement (I3)

Combining the estimates derived above, it follows that for each multi-index α with $|\alpha| \le M$ and $d \ge 3$,

$$\frac{d}{dt} \big\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P](t)) \big\|_{L^2} \\ \lesssim \frac{1}{\langle t\rangle^{1/2+\delta}} \big\| \langle \hbar k\rangle |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \big\|_{L^2_k} \Big(1 + \big\| |\nabla_z|^{\delta} W[P](t) \big\|_{H^{\sigma_3}_M} \Big) \\ + \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d+1+\delta}} \big\| \langle t\nabla_z, \nabla_v\rangle W[P](t) \big\|_{H^{\sigma_4}_M} + \frac{\sqrt{K_5}\varepsilon}{\langle t\rangle^{d}} \big\| |\nabla_z|^{\delta} W[P](t) \big\|_{H^{\sigma_3}_M}.$$

Integrating in time and applying the bootstrap assumptions (B1), (B2), and (B3) gives

$$\big\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[P](t)) \big\|_{L^2} - \big\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v\rangle^{\sigma_3} (v^\alpha W[Q_{\mathrm{in}}]) \big\|_{L^2} \\ \lesssim (1 + \sqrt{K_3}\varepsilon) \int_0^t \frac{1}{\langle s\rangle^{1/2+\delta}} \Big\| \langle \hbar k\rangle |k|^{1/2} \langle k, ks\rangle^{\sigma_4} \widehat{\rho_Q}(s,k) \Big\|_{L^2_k} ds \\ + \sqrt{K_5}\varepsilon \int_0^t \frac{1}{\langle s\rangle^{3/2+\delta}} \sup_{0\le s\le T^*} \left(\frac{\| \langle s\nabla_z, \nabla_v\rangle W[P](s) \|_{H^{\sigma_4}_M}}{\langle s\rangle^{d-1/2}} \right) ds + \sqrt{K_3 K_5}\varepsilon^2 \int_0^t \frac{1}{\langle s\rangle^{d}}\, ds \\ \lesssim (1 + \sqrt{K_3}\varepsilon) \left(\int_0^t \frac{1}{\langle s\rangle^{1+2\delta}}\, ds \right)^{1/2} \Big\| \langle \hbar k\rangle |k|^{1/2} \langle k, kt\rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \Big\|_{L^2_t([0,T^*],L^2_k)} \\ + \sqrt{K_1 K_5}\varepsilon^2 + \sqrt{K_3 K_5}\varepsilon^2 \\ \lesssim \varepsilon \Big[\sqrt{K_2} + \big(\sqrt{K_2 K_3} + \sqrt{K_1 K_5} + \sqrt{K_3 K_5} \big)\varepsilon \Big].$$

Hence, there exists a positive constant $\widetilde{K} > 0$, depending only on the fixed parameters in Proposition 1.12 and independent of $\hbar$, T^*, ε, Q_{in} and the bootstrap

constants K_i, such that

$$\frac{1}{\hbar^d}\sum_{|\alpha|\le M}\big\|\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_3}|\boldsymbol{\nabla}_x|^\delta\boldsymbol{\xi}^\alpha P(t)\big\|_{L^2_{x,y}}^2 \le \widetilde{K}\varepsilon^2\Big[\sqrt{K_2}+\big(\sqrt{K_2K_3}+\sqrt{K_1K_5}+\sqrt{K_3K_5}\big)\varepsilon\Big]^2.$$

Choosing $K_3 := \widetilde{K}(\sqrt{K_2}+1)^2/2$ and $\varepsilon \le (\sqrt{K_2K_3}+\sqrt{K_1K_5}+\sqrt{K_3K_5})^{-1}$ yields the desired bootstrap improvement (I3).

3.3 Bootstrap improvement (I5)

This section establishes the improvement of the bootstrap assumption (B5): it suffices to show

$$\langle k,\eta\rangle^{\sigma_1}\big|\widehat{W[P]}(t,k,\eta)\big| \le \sqrt{2K_5}\,\varepsilon$$

for all $t\in[0,T^*]$, provided ε is sufficiently small.

Using the decomposition (23), the triangle inequality yields

$$\begin{aligned}&\langle k,\eta\rangle^{\sigma_1}\big|\widehat{W[P]}(t,k,\eta)\big|\\ &\quad\le \langle k,\eta\rangle^{\sigma_1}\bigg(\big|\widehat{W[Q_{\rm in}]}(k,\eta)\big|+\int_0^t\big|\widehat{W[\mathrm{L}_P]}(s,k,\eta)\big|\,ds+\int_0^t\big|\widehat{W[\mathrm{N}_P]}(s,k,\eta)\big|\,ds\bigg)\\ &\quad=: \mathcal{I}^{(\mathrm{B5})}(k,\eta)+\mathcal{L}^{(\mathrm{B5})}(k,\eta)+\mathcal{N}^{(\mathrm{B5})}(k,\eta).\end{aligned}$$

The initial term satisfies

$$\mathcal{I}^{(\mathrm{B5})}(k,\eta)=\langle k,\eta\rangle^{\sigma_1}\Big|\widehat{W[Q_{\rm in}]}(k,\eta)\Big| = \Big|\widehat{W[\langle\boldsymbol{\nabla}_x,\boldsymbol{\nabla}_\xi\rangle^{\sigma_1}Q_{\rm in}]}(k,\eta)\Big| \lesssim \varepsilon,$$

by Lemma 1.4 and the assumption on $Q_{\rm in}$ in Theorem 1.3.

To estimate the linear term, apply $\langle k,\eta\rangle^{\sigma_1}\lesssim\langle k,ks\rangle^{\sigma_1}\langle\eta-ks\rangle^{\sigma_1}$, the bound $|\widehat{w}(k)|\lesssim 1$, the bootstrap estimate (B4), and Lemma 1.2:

$$\begin{aligned}&\mathcal{L}^{(\mathrm{B5})}(k,\eta)\\ &\quad=\langle k,\eta\rangle^{\sigma_1}\int_0^t\big|2\hbar^{-1}\widehat{w}(k)\widehat{\rho_Q}(s,k)\sin\big(\tfrac{\hbar}{2}k\cdot(\eta-ks)\big)\,\widehat{g}(\eta-ks)\big|\,ds\\ &\quad\lesssim\int_0^t\langle k,ks\rangle^{\sigma_1}|\widehat{w}(k)||\widehat{\rho_Q}(s,k)||k|\langle\eta-ks\rangle^{\sigma_1}|\eta-ks||\widehat{g}(\eta-ks)|\,ds\\ &\quad\lesssim\big\||k|^{1/2}\langle k,kt\rangle^{\sigma_2}\widehat{\rho_Q}(t,k)\big\|_{L^2_t([0,T^*])}\bigg(\int_0^t|k|\langle\eta-ks\rangle^{2\sigma_1+2}|\widehat{g}(\eta-ks)|^2\,ds\bigg)^{1/2}\\ &\quad\lesssim\sqrt{K_4}\varepsilon\bigg(\int_0^\infty\langle\eta-\tfrac{k}{|k|}s\rangle^{2\sigma_1+2}\big|\widehat{g}(\eta-\tfrac{k}{|k|}s)\big|^2\,ds\bigg)^{1/2}\\ &\quad\lesssim\sqrt{K_4}\varepsilon\|g\|_{H^{\sigma_1+1}_M}.\end{aligned}$$

For the nonlinear term, the inequality $\langle k,\eta\rangle^{\sigma_1} \lesssim \langle \ell, \ell s\rangle^{\sigma_1} + \langle k-\ell, \eta-\ell s\rangle^{\sigma_1}$ leads to the decomposition $\mathcal{N}^{(\mathrm{B5})}(k,\eta) \lesssim \mathcal{R}^{(\mathrm{B5})}(k,\eta) + \mathcal{T}^{(\mathrm{B5})}(k,\eta)$, with

$$\mathcal{R}^{(\mathrm{B5})}(k,\eta) \lesssim \int_0^t \int_\ell \langle \ell, \ell s\rangle^{\sigma_1} |\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)||\ell||\eta - ks| \big|\widehat{W[P]}(s, k-\ell, \eta-\ell s)\big|\, d\ell ds,$$

$$\mathcal{T}^{(\mathrm{B5})}(k,\eta) \lesssim \int_0^t \int_\ell |\widehat{w}(\ell)||\widehat{\rho_Q}(s,\ell)||\ell||\eta - ks| \times \langle k-\ell, \eta-\ell s\rangle^{\sigma_1} \big|\widehat{W[P]}(s, k-\ell, \eta-\ell s)\big|\, d\ell ds.$$

For the reaction term, use $|\eta - ks| \lesssim \langle s\rangle \langle k-\ell, \eta-\ell s\rangle$ and Cauchy–Schwarz in time for fixed ℓ:

$$\begin{aligned}\mathcal{R}^{(\mathrm{B5})}(k,\eta) &\lesssim \sup_{\ell\in\mathbb{R}^d} \big\| |\ell|^{1/2} \langle \ell, \ell t\rangle^{\sigma_2} \widehat{\rho_Q}(t,\ell)\big\|_{L^2_t([0,T^*])} \\ &\times \int_\ell \left(\int_0^t \langle s\rangle^2 \langle \ell, \ell s\rangle^{-2(\sigma_2-\sigma_1)} |\ell| \langle k-\ell, \eta-\ell s\rangle^2 \Big|\widehat{W[P]}(s, k-\ell, \eta-\ell s)\Big|^2 ds \right)^{1/2} d\ell, \\ &\lesssim \sqrt{K_4 K_5}\varepsilon^2 \int_\ell \left(\int_0^t \langle s\rangle^2 \frac{|\ell|}{\langle \ell, \ell s\rangle^{2(\sigma_2-\sigma_1)} \langle k-\ell, \eta-\ell s\rangle^{2\sigma_1-2}}\, ds \right)^{1/2} d\ell.\end{aligned}$$

Setting $j_1 := \sigma_2 - \sigma_1$ and $j_2 := \sigma_1 - 1$, and noting $\langle \ell, \ell s\rangle^{2(\sigma_2-\sigma_1)} \geq \langle \ell s\rangle^{2j_1}$ and $\langle k-\ell, \eta-\ell s\rangle^{2\sigma_1-2} \geq \langle k-\ell\rangle^{2j_2}$, the bound becomes

$$\begin{aligned}\mathcal{R}^{(\mathrm{B5})}(k,\eta) &\lesssim \sqrt{K_4 K_5}\varepsilon^2 \int_\ell \frac{1}{\langle k-\ell\rangle^{j_2}} \left(\int_0^t |\ell| \frac{\langle s\rangle^2}{\langle \ell s\rangle^{2j_1}}\, ds \right)^{1/2} d\ell \\ &\lesssim \sqrt{K_4 K_5}\varepsilon^2 \int_\ell \frac{1}{\langle k-\ell\rangle^{j_2}} \left(\int_0^\infty \frac{\langle s/|\ell|\rangle^2}{\langle s\rangle^{2j_1}}\, ds \right)^{1/2} d\ell \\ &\lesssim \sqrt{K_4 K_5}\varepsilon^2 \int_\ell \frac{\max\{1, |\ell|^{-1}\}}{\langle \ell\rangle^{j_2}} d\ell,\end{aligned}$$

which converges for $j_1 > 3/2$ (i.e. $\sigma_2 - \sigma_1 > 3/2$) and $j_2 > d$ (i.e. $\sigma_1 > d+1$).

For the transport term, again use $|\eta - ks| \lesssim \langle s\rangle \langle k-\ell, \eta-\ell s\rangle$ and apply Cauchy–Schwarz in ℓ:

$$\begin{aligned}\mathcal{T}^{(\mathrm{B5})}(k,\eta) &\lesssim \int_0^t \langle s\rangle \int_\ell |\widehat{\rho_Q}(s,\ell)||\ell| \langle k-\ell, \eta-\ell s\rangle^{\sigma_1+1} \big|\widehat{W[P]}(s, k-\ell, \eta-\ell s)\big|\, d\ell ds \\ &\lesssim \sqrt{K_5}\varepsilon \int_0^t \langle s\rangle \left(\int_\ell \frac{|\ell|^2}{\langle \ell, \ell s\rangle^{2\sigma_1} |k-\ell|^{2\delta}}\, d\ell \right)^{1/2} \\ &\quad \times \left(\int_\ell |k-\ell|^{2\delta} \langle k-\ell, \eta-\ell s\rangle^{2\sigma_1+2} \big|\widehat{W[P]}(s, k-\ell, \eta-\ell s)\big|^2\, d\ell \right)^{1/2} ds.\end{aligned}$$

Treating the ℓ integral as in (24) gives

$$\sup_k \int_\ell \frac{|\ell|^2}{\langle \ell, \ell s\rangle^{2\sigma_1} |k-\ell|^{2\delta}}\, d\ell \lesssim \frac{1}{\langle s\rangle^{d+2-2\delta}},$$

provided that $\sigma_1 > d/2+1$. For the second ℓ integral, the change of variables $\ell \mapsto k-\ell$, the Sobolev embedding and bootstrap assumption (B3) yield

$$\begin{aligned}\big\||k-\ell|^{\delta}\langle k-\ell,\eta-\ell s\rangle^{\sigma_1+1}\widehat{W[P]}(s,k-\ell,\eta-\ell s)\big\|_{L^{\infty}_{\eta}L^{2}_{\ell}}&\\ \lesssim \big\||k|^{\delta}\langle k,\eta\rangle^{\sigma_3}\widehat{W[P]}(s,k,\eta)\big\|_{L^{2}_{k}L^{\infty}_{\eta}}&\\ \lesssim \sum_{|\alpha|\leq M}\big\|D^{\alpha}_{\eta}\big(|k|^{\delta}\langle k,\eta\rangle^{\sigma_3}\widehat{W[P]}(s,k,\eta)\big)\big\|_{L^{2}_{k,\eta}}&\\ \lesssim \sqrt{K_3}\varepsilon.&\end{aligned}$$

Therefore,

$$\mathcal{T}^{(\mathrm{B5})}(k,\eta)\lesssim \sqrt{K_3K_5}\varepsilon^2\int_0^t\frac{1}{\langle s\rangle^{d/2-\delta}}\,ds\lesssim\sqrt{K_3K_5}\varepsilon^2.$$

Conclusion of bootstrap improvement (I5)

Under the bootstrap assumptions (B3), (B4), and (B5), there exists a constant $\widetilde{K}>0$, depending only on the fixed parameters in Proposition 1.12 and independent of $\hbar$, T^*, ε, Q_{in} and the bootstrap constants K_i, such that

$$\langle k,\eta\rangle^{\sigma_1}\big|\widehat{W[P]}(t,k,\eta)\big|\leq\widetilde{K}\Big(\varepsilon+\sqrt{K_4}\varepsilon+\big(\sqrt{K_4K_5}+\sqrt{K_3K_5}\big)\varepsilon^2\Big).$$

Choose ε sufficiently small such that $\big(\sqrt{K_4K_5}+\sqrt{K_3K_5}\big)\,\varepsilon\leq 1$. Then

$$\langle k,\eta\rangle^{\sigma_1}\big|\widehat{W[P]}(t,k,\eta)\big|\leq\widetilde{K}\big(2+\sqrt{K_4}\big)\varepsilon.$$

Setting

$$\sqrt{K_5}:=\frac{\widetilde{K}(2+\sqrt{K_4})}{\sqrt{2}}$$

establishes the bootstrap improvement (I5):

$$\langle k,\eta\rangle^{\sigma_1}\big|\widehat{W[P]}(t,k,\eta)\big|\leq\sqrt{2}\sqrt{K_5}\,\varepsilon.$$

This completes the proof of Proposition 1.12.

参考文献

[1] J. Bedrossian - "Nonlinear echoes and Landau damping with insufficient regularity" , Tunis. J. Math. 3 (2021) no. 1, p. 121-205

[2] J. Bedrossian, N. Masmoudi & C. Mouhot - "Landau damping in finite regularity for unconfined systems with screened interactions" , Comm. Pure Appl. Math. 71 (2018) no. 3, p. 537-576

[3] M. Lewin & J. Sabin - "The Hartree equation for infinitely many particles, II: Dispersion and scattering in 2D" , Anal. PDE 7 (2014) no. 6, p. 1339-1363

[4] T. T. Nguyen & C. You - "Plasmons for the Hartree equations with Coulomb interaction" , Probab. Math. Phys. 6 (2025) no. 3, p. 913-960

[5] E. M. Stein & G. Weiss - "Interpolation of operators with change of measures" , Trans. Amer. Math. Soc. 87 (1958), p. 159-172