THE DERIVED TENSOR FUNCTOR WITH A COMPACT COMPLEX IS FULLY FAITHFUL

THE STACKS PROJECT AUTHORS

Editorial note. The counted statement is the author's own lemma environment labelled lemma-fully-faithful-in-compact-case (source lines 6338-6356); the proof environment begins on the second line after it and is its own proof, closing at source line 6374.

Editorial note. The statement and the proof invoke three author results of the same chapter by label; all three are retained as context spans and the generator expands the context transitively (budget 800 lines for this card), while reference targets in other chapters that carry canonical project tags are auto-declared as non-load-bearing source pointers.

Editorial note. The retained preamble is the chapter's own minimal preamble (sources/stacks-dga/preamble.tex, taken with its real line count), a verbatim excerpt of the native Xy-pic setup, the native theorem declarations with the mathematical macros, and the project's own no-op reference environments; no macro was invented or redefined.

Editorial scope. One original result of the Stacks Project's Differential Graded Algebra chapter, counted once: the derived tensor functor with a compact complex whose endomorphism algebra computes its Ext groups is fully faithful.

Allowed standard prerequisites (editorial).

- Ringed site and complexes of O-modules. A ringed site (C, O) and the derived category D(O) of complexes of O-modules, with its compact objects.. It is the setting and the hypothesis that K^bullet represents a compact object (source lines 6340-6344).
- Derived tensor product with a fixed complex. The left derived functor - tensor^L_E K^bullet : D(E,d) to D(O) of the earlier lemma, and its right adjoint R Hom(K^bullet, -) (the author's lemmas lemma-tensor-with-complex-derived and lemma-tensor-with-complex-hom-adjoint, retained as context).. The functor whose full faithfulness is asserted, and the adjunction used to reduce the claim (source lines 6350-6360).
- Endomorphism dg algebra computing Ext. E = Hom over the dg category of complexes from K^bullet to itself, assumed to compute the Ext groups of K^bullet in D(O).. It is the algebra the source category D(E,d) is taken over, and it is the second hypothesis of the statement (source lines 6345-6346, 6372-6373).

Date: stacks-project:dga.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

- Property (P) modules and their filtration sequences. Every dg E-module has a quasi-isomorphic representative with property (P), carrying an admissible filtration with graded pieces direct sums of shifts of E, and a split admissible short exact sequence displaying the filtration (the author's lemma lemma-property-P-sequence, retained as context).. The replacements and reductions of the proof run along this filtration (source lines 6366-6371).

Source reference. Original lemma 06XN: sources/stacks-chow/tags.txt, source lines 8857–8857. Cross-chapter pointer whose target carries the canonical tag 06XN in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 09IZ: sources/stacks-chow/tags.txt, source lines 12057–12057. Cross-chapter pointer whose target carries the canonical tag 09IZ in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 0149: sources/stacks-chow/tags.txt, source lines 1464–1464. Cross-chapter pointer whose target carries the canonical tag 0149 in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 09J0: sources/stacks-chow/tags.txt, source lines 12058–12058. Cross-chapter pointer whose target carries the canonical tag 09J0 in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 014C: sources/stacks-chow/tags.txt, source lines 1467–1467. Cross-chapter pointer whose target carries the canonical tag 014C in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 013N: sources/stacks-chow/tags.txt, source lines 1442–1442. Cross-chapter pointer whose target carries the canonical tag 013N in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Lemma 0.1. *The functor $K(Mod_{(E,d)}) \to K(\mathcal{O})$ of Lemma 0.6 has a left derived version defined on all of $D(E,d)$. We denote it $- \otimes^{\mathbf{L}}_E K^\bullet : D(E,d) \to D(\mathcal{O})$.*

Proof. We will use Derived Categories, Lemma 06XN to prove this. As our collection $\mathcal{P}$ of objects we will use the objects with property (P). Property (1) was shown in Lemma 0.5. Property (2) holds because if $s : P \to P'$ is a quasi-isomorphism of modules with property (P), then s is a homotopy equivalence by Lemma 0.4. □

Lemma 0.2. *Let $(\mathcal{C}, \mathcal{O})$ be a ringed site. Let $K^\bullet$ be a complex of $\mathcal{O}$-modules. Then the functor*

$$- \otimes^{\mathbf{L}}_E K^\bullet : D(E,d) \longrightarrow D(\mathcal{O})$$

of Lemma 0.1 is a left adjoint of the functor

$$R\mathrm{Hom}(K^\bullet, -) : D(\mathcal{O}) \longrightarrow D(E,d)$$

of Lemma 0.7.

Proof. The statement means that we have

$$\mathrm{Hom}_{D(E,\mathrm{d})}(M, R\,\mathrm{Hom}(K^\bullet, L^\bullet)) = \mathrm{Hom}_{D(\mathcal{O})}(M \otimes^{\mathbf{L}}_E K^\bullet, L^\bullet)$$

bifunctorially in M and $L^\bullet$. To see this we may replace M by a differential graded E-module P with property (P). We also may replace $L^\bullet$ by a K-injective complex of $\mathcal{O}$-modules $I^\bullet$. The computation of the derived functors given in the lemmas referenced in the statement combined with Lemma 0.4 translates the above into

$$\mathrm{Hom}_{K(\mathrm{Mod}_{(E,\mathrm{d})})}(P, \mathrm{Hom}_{\mathcal{B}}(K^\bullet, I^\bullet)) = \mathrm{Hom}_{K(\mathcal{O})}(P \otimes_E K^\bullet, I^\bullet)$$

where $\mathcal{B} = \mathrm{Comp}^{dg}(\mathcal{O})$. There is an evaluation map from right to left functorial in P and $I^\bullet$ (details omitted). Choose a filtration $F_\bullet$ on P as in the definition of property (P). By Lemma 0.3 and the fact that both sides of the equation are homological functors in P on $K(\mathrm{Mod}_{(E,\mathrm{d})})$ we reduce to the case where P is replaced by the differential graded E-module $\bigoplus F_iP$. Since both sides turn direct sums in the variable P into direct products we reduce to the case where P is one of the differential graded E-modules F_iP. Since each F_iP has a finite filtration (given by admissible monomorphisms) whose graded pieces are graded projective E-modules we reduce to the case where P is a graded projective E-module. In this case we clearly have

$$\mathrm{Hom}_{\mathrm{Mod}^{dg}_{(E,\mathrm{d})}}(P, \mathrm{Hom}_{\mathcal{B}}(K^\bullet, I^\bullet)) = \mathrm{Hom}_{\mathrm{Comp}^{dg}(\mathcal{O})}(P \otimes_E K^\bullet, I^\bullet)$$

as graded $\mathbf{Z}$-modules (because this statement reduces to the case $P = E[k]$ where it is obvious). As the isomorphism is compatible with differentials we conclude. □

Lemma 0.3. *Let (A, d) be a differential graded algebra. Let P be a differential graded A-module. If $F_\bullet$ is a filtration as in property (P), then we obtain an admissible short exact sequence*

$$0 \to \bigoplus F_iP \to \bigoplus F_iP \to P \to 0$$

of differential graded A-modules.

Proof. The second map is the direct sum of the inclusion maps. The first map on the summand F_iP of the source is the sum of the identity $F_iP \to F_iP$ and the negative of the inclusion map $F_iP \to F_{i+1}P$. Choose homomorphisms $s_i : F_{i+1}P \to F_iP$ of graded A-modules which are left inverse to the inclusion maps. Composing gives maps $s_{j,i} : F_jP \to F_iP$ for all $j > i$. Then a left inverse of the first arrow maps $x \in F_jP$ to $(s_{j,0}(x), s_{j,1}(x), \ldots, s_{j,j-1}(x), 0, \ldots)$ in $\bigoplus F_iP$. □

Lemma 0.4. *Let (A, d) be a differential graded algebra. Let M and N be differential graded A-modules.*

(1) *Let $P \to M$ be a P-resolution as in Lemma 0.5. Then*

$$\mathrm{Hom}_{D(A,d)}(M, N) = \mathrm{Hom}_{K(Mod_{(A,d)})}(P, N)$$

(2) *Let $N \to I$ be an I-resolution as in Lemma 0.10. Then*

$$\mathrm{Hom}_{D(A,d)}(M, N) = \mathrm{Hom}_{K(Mod_{(A,d)})}(M, I)$$

Proof. Let $P \to M$ be as in (1). Since $P \to M$ is a quasi-isomorphism we see that

$$\mathrm{Hom}_{D(A,\mathrm{d})}(P, N) = \mathrm{Hom}_{D(A,\mathrm{d})}(M, N)$$

by definition of the derived category. A morphism $f : P \to N$ in $D(A, \mathrm{d})$ is equal to $s^{-1}f'$ where $f' : P \to N'$ is a morphism and $s : N \to N'$ is a quasi-isomorphism. Choose a distinguished triangle

$$N \to N' \to Q \to N[1]$$

As s is a quasi-isomorphism, we see that Q is acyclic. Thus $\mathrm{Hom}_{K(\mathrm{Mod}_{(A,\mathrm{d})})}(P, Q[k]) = 0$ for all k by Lemma 0.9. Since $\mathrm{Hom}_{K(\mathrm{Mod}_{(A,\mathrm{d})})}(P, -)$ is cohomological, we conclude that we can lift $f' : P \to N'$ uniquely to a morphism $f : P \to N$. This finishes the proof.

The proof of (2) is dual to that of (1) using Lemma 0.8 in stead of Lemma 0.9. □

Lemma 0.5. *Let (A, d) be a differential graded algebra. Let M be a differential graded A-module. There exists a homomorphism $P \to M$ of differential graded A-modules such that*

(1) *$P \to M$ is a quasi-isomorphism, and*
(2) *P has property (P).*

Proof. Set $M = M_0$. We inductively choose short exact sequences

$$0 \to M_{i+1} \to P_i \to M_i \to 0$$

where the maps $P_i \to M_i$ are chosen as in Lemma 0.11. This gives a "resolution"

$$\ldots \to P_2 \xrightarrow{f_2} P_1 \xrightarrow{f_1} P_0 \to M \to 0$$

Then we set

$$P = \bigoplus\nolimits_{i \geq 0} P_i$$

as an A-module with grading given by $P^n = \bigoplus_{a+b=n} P^b_{-a}$ and differential (as in the construction of the total complex associated to a double complex) by

$$\mathrm{d}_P(x) = f_{-a}(x) + (-1)^a \mathrm{d}_{P_{-a}}(x)$$

for $x \in P^b_{-a}$. With these conventions P is indeed a differential graded A-module. Recalling that each P_i has a two step filtration $0 \to P'_i \to P_i \to P''_i \to 0$ we set

$$F_{2i}P = \bigoplus\nolimits_{i \geq j \geq 0} P_j \subset \bigoplus\nolimits_{i \geq 0} P_i = P$$

and we add P'_{i+1} to $F_{2i}P$ to get F_{2i+1}. These are differential graded submodules and the successive quotients are direct sums of shifts of A. By Lemma 0.12 we see that the inclusions $F_iP \to F_{i+1}P$ are admissible monomorphisms. Finally, we have to show that the map $P \to M$ (given by the augmentation $P_0 \to M$) is a quasi-isomorphism. This follows from Homology, Lemma 09IZ. □

Lemma 0.6. *The functor of Lemma 0.14 defines an exact functor of triangulated categories $K(Mod_{(E,d)}) \to K(\mathcal{O})$.*

Proof. The functor induces a functor between homotopy categories by Lemma 0.13. We have to show that $- \otimes_E K^\bullet$ transforms distinguished triangles into distinguished triangles. Suppose that $0 \to K \to L \to M \to 0$ is an admissible short exact sequence of differential graded E-modules. Let $s : M \to L$ be a graded E-module homomorphism which is left inverse to $L \to M$. Then s defines a map $M \otimes_E K^\bullet \to L \otimes_E K^\bullet$ of graded $\mathcal{O}$-modules (i.e., respecting $\mathcal{O}$-module structure and grading,

but not differentials) which is left inverse to $L \otimes_E K^\bullet \to M \otimes_E K^\bullet$. Thus we see that

$$0 \to K \otimes_E K^\bullet \to L \otimes_E K^\bullet \to M \otimes_E K^\bullet \to 0$$

is a termwise split short exact sequences of complexes, i.e., a defines a distinguished triangle in $K(\mathcal{O})$. □

Lemma 0.7. *In the situation above. If the right derived functor $R\operatorname{Hom}(K^\bullet, -)$ of $\operatorname{Hom}(K^\bullet, -) : K(\mathcal{A}) \to D(Ab)$ is everywhere defined on $D(\mathcal{A})$, then we obtain a canonical exact functor*

$$R\operatorname{Hom}(K^\bullet, -) : D(\mathcal{A}) \longrightarrow D(E, d)$$

of triangulated categories which reduces to the usual one on taking associated complexes of abelian groups.

Proof. Note that we have an associated functor $K(\mathcal{A}) \to K(\operatorname{Mod}_{(E,\mathrm{d})})$ by Lemma 0.15. We claim this functor is an exact functor of triangulated categories. Namely, let $f : A^\bullet \to B^\bullet$ be a map of complexes of $\mathcal{A}$. Then a computation shows that

$$\operatorname{Hom}_{\operatorname{Comp}^{dg}(\mathcal{A})}(K^\bullet, C(f)^\bullet) = C\left(\operatorname{Hom}_{\operatorname{Comp}^{dg}(\mathcal{A})}(K^\bullet, A^\bullet) \to \operatorname{Hom}_{\operatorname{Comp}^{dg}(\mathcal{A})}(K^\bullet, B^\bullet)\right)$$

where the right hand side is the cone in $\operatorname{Mod}_{(E,\mathrm{d})}$ defined earlier in this chapter. This shows that our functor is compatible with cones, hence with distinguished triangles. Let $X^\bullet$ be an object of $K(\mathcal{A})$. Consider the category of quasi-isomorphisms $s : X^\bullet \to Y^\bullet$. We are given that the functor $(s : X^\bullet \to Y^\bullet) \mapsto \operatorname{Hom}_{\mathcal{A}}(K^\bullet, Y^\bullet)$ is essentially constant when viewed in $D(Ab)$. But since the forgetful functor $D(E, \mathrm{d}) \to D(Ab)$ is compatible with taking cohomology, the same thing is true in $D(E, \mathrm{d})$. This proves the lemma. □

Lemma 0.8. *Let (A, d) be a differential graded algebra. Let I be a differential graded A-module with property (I). Then*

$$\operatorname{Hom}_{K(Mod_{(A,d)})}(N, I) = 0$$

for all acyclic differential graded A-modules N.

Proof. We will use that $K(\operatorname{Mod}_{(A,\mathrm{d})})$ is a triangulated category (Proposition 0.18). Let $F_\bullet$ be a filtration on I as in property (I). The short exact sequence of Lemma 0.17 produces a distinguished triangle. Hence by Derived Categories, Lemma 0149 it suffices to show that

$$\operatorname{Hom}_{K(\operatorname{Mod}_{(A,\mathrm{d})})}(N, I/F_iI) = 0$$

for all acyclic differential graded A-modules N and all i. Each of the differential graded modules I/F_iI has a finite filtration by admissible monomorphisms, whose graded pieces are products of $A^\vee[k]$. Thus it suffices to prove that

$$\operatorname{Hom}_{K(\operatorname{Mod}_{(A,\mathrm{d})})}(N, A^\vee[k]) = 0$$

for all acyclic differential graded A-modules N and all k. This follows from Lemma 0.16 and the fact that $(-)^\vee$ is an exact functor. □

Lemma 0.9. *Let (A, d) be a differential graded algebra. Let P be a differential graded A-module with property (P). Then*

$$\operatorname{Hom}_{K(Mod_{(A,d)})}(P, N) = 0$$

for all acyclic differential graded A-modules N.

Proof. We will use that $K(\mathrm{Mod}_{(A,\mathrm{d})})$ is a triangulated category (Proposition 0.18). Let $F_\bullet$ be a filtration on P as in property (P). The short exact sequence of Lemma 0.3 produces a distinguished triangle. Hence by Derived Categories, Lemma 0149 it suffices to show that

$$\mathrm{Hom}_{K(\mathrm{Mod}_{(A,\mathrm{d})})}(F_iP, N) = 0$$

for all acyclic differential graded A-modules N and all i. Each of the differential graded modules F_iP has a finite filtration by admissible monomorphisms, whose graded pieces are direct sums of shifts $A[k]$. Thus it suffices to prove that

$$\mathrm{Hom}_{K(\mathrm{Mod}_{(A,\mathrm{d})})}(A[k], N) = 0$$

for all acyclic differential graded A-modules N and all k. This follows from Lemma 0.19. □

Lemma 0.10. *Let (A, d) be a differential graded algebra. Let M be a differential graded A-module. There exists a homomorphism $M \to I$ of differential graded A-modules such that*

(1) *$M \to I$ is a quasi-isomorphism, and*
(2) *I has property (I).*

Proof. Set $M = M_0$. We inductively choose short exact sequences

$$0 \to M_i \to I_i \to M_{i+1} \to 0$$

where the maps $M_i \to I_i$ are chosen as in Lemma 0.20. This gives a "resolution"

$$0 \to M \to I_0 \xrightarrow{f_0} I_1 \xrightarrow{f_1} I_1 \to \ldots$$

Denote I the differential graded A-module with graded parts

$$I^n = \prod\nolimits_{i \geq 0} I_i^{n-i}$$

and differential defined by

$$\mathrm{d}_I(x) = f_i(x) + (-1)^i \mathrm{d}_{I_i}(x)$$

for $x \in I_i^{n-i}$. With these conventions I is indeed a differential graded A-module. Recalling that each I_i has a two step filtration $0 \to I_i' \to I_i \to I_i'' \to 0$ we set

$$F_{2i}I^n = \prod\nolimits_{j \geq i} I_j^{n-j} \subset \prod\nolimits_{i \geq 0} I_i^{n-i} = I^n$$

and we add a factor I_{i+1}' to $F_{2i}I$ to get $F_{2i+1}I$. These are differential graded submodules and the successive quotients are products of shifts of $A^\vee$. By Lemma 0.21 we see that the inclusions $F_{i+1}I \to F_iI$ are admissible monomorphisms. Finally, we have to show that the map $M \to I$ (given by the augmentation $M \to I_0$) is a quasi-isomorphism. This follows from Homology, Lemma 09J0. □

Lemma 0.11. *Let (A, d) be a differential graded algebra. Let M be a differential graded A-module. There exists a homomorphism $P \to M$ of differential graded A-modules with the following properties*

(1) *$P \to M$ is surjective,*
(2) *$\mathrm{Ker}(d_P) \to \mathrm{Ker}(d_M)$ is surjective, and*
(3) *P sits in an admissible short exact sequence $0 \to P' \to P \to P'' \to 0$ where P', P'' are direct sums of shifts of A.*

Proof. Let P_k be the free A-module with generators x, y in degrees k and $k+1$. Define the structure of a differential graded A-module on P_k by setting $\mathrm{d}(x) = y$ and $\mathrm{d}(y) = 0$. For every element $m \in M^k$ there is a homomorphism $P_k \to M$ sending x to m and y to $\mathrm{d}(m)$. Thus we see that there is a surjection from a direct sum of copies of P_k to M. This clearly produces $P \to M$ having properties (1) and (3). To obtain property (2) note that if $m \in \mathrm{Ker}(\mathrm{d}_M)$ has degree k, then there is a map $A[k] \to M$ mapping 1 to m. Hence we can achieve (2) by adding a direct sum of copies of shifts of A. □

Lemma 0.12. *Let (A, d) be a differential graded algebra. Let $M \to P$ be a surjective homomorphism of differential graded A-modules. If P is projective as a graded A-module, then $M \to P$ is an admissible epimorphism.*

Proof. This is immediate from the definitions. □

Lemma 0.13. *Let R be a ring. A functor $F : \mathcal{A} \to \mathcal{B}$ of differential graded categories over R induces functors $Comp(\mathcal{A}) \to Comp(\mathcal{B})$ and $K(\mathcal{A}) \to K(\mathcal{B})$.*

Proof. Omitted. □

Lemma 0.14. *In the situation above there is a functor*

$$- \otimes_E K^\bullet : Mod^{dg}_{(E,d)} \longrightarrow Comp^{dg}(\mathcal{O})$$

of differential graded categories. This functor sends E to $K^\bullet$ and commutes with direct sums.

Proof. Let M be a differential graded E-module. For every object U of $\mathcal{C}$ the complex $K^\bullet(U)$ is a left differential graded E-module as well as a right $\mathcal{O}(U)$-module. The actions commute, so we have a bimodule. Thus, by the constructions in Sections 2 and 1 we can form the tensor product

$$M \otimes_E K^\bullet(U)$$

which is a differential graded $\mathcal{O}(U)$-module, i.e., a complex of $\mathcal{O}(U)$-modules. This construction is functorial with respect to U, hence we can sheafify to get a complex of $\mathcal{O}$-modules which we denote

$$M \otimes_E K^\bullet$$

Moreover, for each U the construction determines a functor $\mathrm{Mod}^{dg}_{(E,\mathrm{d})} \to \mathrm{Comp}^{dg}(\mathcal{O}(U))$ of differential graded categories by Lemma 0.22. It is therefore clear that we obtain a functor as stated in the lemma. □

Lemma 0.15. *Let R be a ring. Let $\mathcal{A}$ be a differential graded category over R. Let x be an object of $\mathcal{A}$. Let*

$$(E, d) = \mathrm{Hom}_{\mathcal{A}}(x, x)$$

be the differential graded R-algebra of endomorphisms of x. We obtain a functor

$$\mathcal{A} \longrightarrow Mod^{dg}_{(E,d)}, \quad y \longmapsto \mathrm{Hom}_{\mathcal{A}}(x, y)$$

of differential graded categories by letting E act on $\mathrm{Hom}_{\mathcal{A}}(x, y)$ via composition in $\mathcal{A}$. This functor induces functors

$$Comp(\mathcal{A}) \to Mod_{(A,d)} \quad \text{and} \quad K(\mathcal{A}) \to K(Mod_{(A,d)})$$

by an application of Lemma 0.13.

Proof. This lemma proves itself. □

Lemma 0.16. *Let (A, d) be a differential graded algebra. Then we have*

$$\mathrm{Hom}_{Mod_{(A,d)}}(M, A^\vee[k]) = \mathrm{Ker}(d : (M^\vee)^k \to (M^\vee)^{k+1})$$

and

$$\mathrm{Hom}_{K(Mod_{(A,d)})}(M, A^\vee[k]) = H^k(M^\vee)$$

as functors in the differential graded A-module M.

Proof. This is clear from the discussion above. □

Lemma 0.17. *Let (A, d) be a differential graded algebra. Let I be a differential graded A-module. If $F_\bullet$ is a filtration as in property (I), then we obtain an admissible short exact sequence*

$$0 \to I \to \prod I/F_i I \to \prod I/F_i I \to 0$$

of differential graded A-modules.

Proof. Omitted. Hint: This is dual to Lemma 0.3. □

Proposition 0.18. *Let (A, d) be a differential graded algebra. The homotopy category $K(Mod_{(A,d)})$ of differential graded A-modules with its natural translation functors and distinguished triangles is a triangulated category.*

Proof. We know that $K(\mathrm{Mod}_{(A,\mathrm{d})})$ is a pre-triangulated category. Hence it suffices to prove TR4 and to prove it we can use Derived Categories, Lemma 014C. Let $K \to L$ and $L \to M$ be composable morphisms of $K(\mathrm{Mod}_{(A,\mathrm{d})})$. By Lemma 2.1 we may assume that $K \to L$ and $L \to M$ are admissible monomorphisms. In this case the result follows from Lemma 2.2. □

Lemma 0.19. *Let (A, d) be a differential graded algebra. Then we have*

$$\mathrm{Hom}_{Mod_{(A,d)}}(A[k], M) = \mathrm{Ker}(d : M^{-k} \to M^{-k+1})$$

and

$$\mathrm{Hom}_{K(Mod_{(A,d)})}(A[k], M) = H^{-k}(M)$$

for any differential graded A-module M.

Proof. Immediate from the definitions. □

Lemma 0.20. *Let (A, d) be a differential graded algebra. Let M be a differential graded A-module. There exists a homomorphism $M \to I$ of differential graded A-modules with the following properties*

1. *$M \to I$ is injective,*
2. *$\mathrm{Coker}(d_M) \to \mathrm{Coker}(d_I)$ is injective, and*
3. *I sits in an admissible short exact sequence $0 \to I' \to I \to I'' \to 0$ where I', I'' are products of shifts of $A^\vee$.*

Proof. We will use the functors $N \mapsto N^\vee$ (from left to right differential graded modules and from right to left differential graded modules) constructed in Section 3 and all of their properties. For every $k \in \mathbf{Z}$ let Q_k be the free left A-module with generators x, y in degrees k and $k+1$. Define the structure of a left differential graded A-module on Q_k by setting $\mathrm{d}(x) = y$ and $\mathrm{d}(y) = 0$. Arguing exactly as in the proof of Lemma 0.11 we find a surjection

$$\bigoplus\nolimits_{i \in I} Q_{k_i} \longrightarrow M^\vee$$

of left differential graded A-modules. Then we can consider the injection

$$M \to (M^\vee)^\vee \to (\bigoplus\nolimits_{i\in I} Q_{k_i})^\vee = \prod\nolimits_{i\in I} I_{k_i}$$

where $I_k = Q_{-k}^\vee$ is the "dual" right differential graded A-module. Further, the short exact sequence $0 \to A[-k-1] \to Q_k \to A[-k] \to 0$ produces a short exact sequence $0 \to A^\vee[k] \to I_k \to A^\vee[k+1] \to 0$.

The result of the previous paragraph produces $M \to I$ having properties (1) and (3). To obtain property (2), suppose $\overline{m} \in \mathrm{Coker}(\mathrm{d}_M)$ is a nonzero element of degree k. Pick a map $\lambda : M^k \to \mathbf{Q}/\mathbf{Z}$ which vanishes on $\mathrm{Im}(M^{k-1} \to M^k)$ but not on m. By Lemma 0.16 this corresponds to a homomorphism $M \to A^\vee[k]$ of differential graded A-modules which does not vanish on m. Hence we can achieve (2) by adding a product of copies of shifts of $A^\vee$. □

Lemma 0.21. *Let (A, d) be a differential graded algebra. Let $I \to M$ be an injective homomorphism of differential graded A-modules. If I is graded injective, then $I \to M$ is an admissible monomorphism.*

Proof. This is immediate from the definitions. □

Lemma 0.22. *Let R be a ring. Let (A, d) and (B, d) be differential graded algebras over R. Let N be a differential graded (A, B)-bimodule. Then $M \mapsto M \otimes_A N$ defines a functor*

$$- \otimes_A N : Mod^{dg}_{(A,d)} \longrightarrow Mod^{dg}_{(B,d)}$$

of differential graded categories. This functor induces functors

$$Mod_{(A,d)} \to Mod_{(B,d)} \quad \text{and} \quad K(Mod_{(A,d)}) \to K(Mod_{(B,d)})$$

by an application of Lemma 0.13.

Proof. Above we have seen how the construction defines a functor of underlying graded categories. Thus it suffices to show that the construction is compatible with differentials. Let M and M' be differential graded A-modules and let $f : M \to M'$ be an A-module homomorphism which is homogeneous of degree n. Then we have

$$\mathrm{d}(f) = \mathrm{d}_{M'} \circ f - (-1)^n f \circ \mathrm{d}_M$$

On the other hand, we have

$$\mathrm{d}(f \otimes \mathrm{id}_N) = \mathrm{d}_{M' \otimes_A N} \circ (f \otimes \mathrm{id}_N) - (-1)^n (f \otimes \mathrm{id}_N) \circ \mathrm{d}_{M \otimes_A N}$$

Applying this to an element $x \otimes y$ with $x \in M$ and $y \in N$ homogeneous we get

$$\begin{aligned} \mathrm{d}(f \otimes \mathrm{id}_N)(x \otimes y) =& \mathrm{d}_{M'}(f(x)) \otimes y + (-1)^{n+\deg(x)} f(x) \otimes \mathrm{d}_N(y) \\ & - (-1)^n f(\mathrm{d}_M(x)) \otimes y - (-1)^{n+\deg(x)} f(x) \otimes \mathrm{d}_N(y) \\ =& \mathrm{d}(f)(x \otimes y) \end{aligned}$$

Thus we see that $\mathrm{d}(f) \otimes \mathrm{id}_N = \mathrm{d}(f \otimes \mathrm{id}_N)$ and the proof is complete. □

1. Bimodules

2. Tensor product

Lemma 2.1. *Let (A, d) be a differential graded algebra. Let $L_1 \to L_2 \to \ldots \to L_n$ be a sequence of composable homomorphisms of differential graded A-modules. There exists a commutative diagram*

$$\begin{array}{ccccccc} L_1 & \longrightarrow & L_2 & \longrightarrow & \ldots & \longrightarrow & L_n \\ \uparrow & & \uparrow & & & & \uparrow \\ M_1 & \longrightarrow & M_2 & \longrightarrow & \ldots & \longrightarrow & M_n \end{array}$$

in $Mod_{(A,d)}$ such that each $M_i \to M_{i+1}$ is an admissible monomorphism and each $M_i \to L_i$ is a homotopy equivalence.

Proof. The case $n = 1$ is without content. Lemma 3.1 is the case $n = 2$. Suppose we have constructed the diagram except for M_n. Apply Lemma 3.1 to the composition $M_{n-1} \to L_{n-1} \to L_n$. The result is a factorization $M_{n-1} \to M_n \to L_n$ as desired. □

Lemma 2.2. *Let (A, d) be a differential graded algebra. Suppose that $\alpha : K \to L$ and $\beta : L \to M$ are admissible monomorphisms of differential graded A-modules. Then there exist distinguished triangles $(K, L, Q_1, \alpha, p_1, d_1)$, $(K, M, Q_2, \beta \circ \alpha, p_2, d_2)$ and $(L, M, Q_3, \beta, p_3, d_3)$ for which TR4 holds.*

Proof. Say $\pi_1 : L \to K$ and $\pi_3 : M \to L$ are homomorphisms of graded A-modules which are left inverse to α and β. Then also $K \to M$ is an admissible monomorphism with left inverse $\pi_2 = \pi_1 \circ \pi_3$. Let us write Q_1, Q_2 and Q_3 for the cokernels of $K \to L$, $K \to M$, and $L \to M$. Then we obtain identifications (as graded A-modules) $Q_1 = \mathrm{Ker}(\pi_1)$, $Q_3 = \mathrm{Ker}(\pi_3)$ and $Q_2 = \mathrm{Ker}(\pi_2)$. Then $L = K \oplus Q_1$ and $M = L \oplus Q_3$ as graded A-modules. This implies $M = K \oplus Q_1 \oplus Q_3$. Note that $\pi_2 = \pi_1 \circ \pi_3$ is zero on both Q_1 and Q_3. Hence $Q_2 = Q_1 \oplus Q_3$. Consider the commutative diagram

$$\begin{array}{ccccccccc} 0 & \to & K & \to & L & \to & Q_1 & \to & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \to & K & \to & M & \to & Q_2 & \to & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \to & L & \to & M & \to & Q_3 & \to & 0 \end{array}$$

The rows of this diagram are admissible short exact sequences, and hence determine distinguished triangles by definition. Moreover downward arrows in the diagram above are compatible with the chosen splittings and hence define morphisms of triangles

$$(K \to L \to Q_1 \to K[1]) \longrightarrow (K \to M \to Q_2 \to K[1])$$

and

$$(K \to M \to Q_2 \to K[1]) \longrightarrow (L \to M \to Q_3 \to L[1]).$$

Note that the splittings $Q_3 \to M$ of the bottom sequence in the diagram provides a splitting for the split sequence $0 \to Q_1 \to Q_2 \to Q_3 \to 0$ upon composing with $M \to Q_2$. It follows easily from this that the morphism $\delta : Q_3 \to Q_1[1]$ in the corresponding distinguished triangle

$$(Q_1 \to Q_2 \to Q_3 \to Q_1[1])$$

is equal to the composition $Q_3 \to L[1] \to Q_1[1]$. Hence we get a structure as in the conclusion of axiom TR4. □

3. Injective modules and differential graded algebras

Lemma 3.1. *Let (A, d) be a differential graded algebra. Let $\alpha : K \to L$ be a homomorphism of differential graded A-modules. There exists a factorization*

$$K \xrightarrow{\tilde{\alpha}} \tilde{L} \xrightarrow{\pi} L, \quad K \xrightarrow{\alpha} L$$

in $Mod_{(A,d)}$ such that

1. *$\tilde{\alpha}$ is an admissible monomorphism (see Definition 3.2),*
2. *there is a morphism $s : L \to \tilde{L}$ such that $\pi \circ s = id_L$ and such that $s \circ \pi$ is homotopic to $id_{\tilde{L}}$.*

Proof. The proof is identical to the proof of Derived Categories, Lemma 013N. Namely, we set $\tilde{L} = L \oplus C(1_K)$ and we use elementary properties of the cone construction. □

Definition 3.2. Let (A, d) be a differential graded algebra.

1. A homomorphism $K \to L$ of differential graded A-modules is an *admissible monomorphism* if there exists a graded A-module map $L \to K$ which is left inverse to $K \to L$.
2. A homomorphism $L \to M$ of differential graded A-modules is an *admissible epimorphism* if there exists a graded A-module map $M \to L$ which is right inverse to $L \to M$.
3. A short exact sequence $0 \to K \to L \to M \to 0$ of differential graded A-modules is an *admissible short exact sequence* if it is split as a sequence of graded A-modules.

Lemma 3.3. *Let $(\mathcal{C}, \mathcal{O})$ be a ringed site. Let $K^\bullet$ be a complex of $\mathcal{O}$-modules. Assume*

1. *$K^\bullet$ represents a compact object of $D(\mathcal{O})$, and*
2. *$E = \mathrm{Hom}_{Comp^{dg}(\mathcal{O})}(K^\bullet, K^\bullet)$ computes the ext groups of $K^\bullet$ in $D(\mathcal{O})$.*

Then the functor

$$- \otimes^{\mathbf{L}}_E K^\bullet : D(E, d) \longrightarrow D(\mathcal{O})$$

of Lemma 0.1 is fully faithful.

Proof. Because our functor has a left adjoint given by $R\mathrm{Hom}(K^\bullet, -)$ by Lemma 0.2 it suffices to show for a differential graded E-module M that the map

$$H^0(M) \longrightarrow \mathrm{Hom}_{D(\mathcal{O})}(K^\bullet, M \otimes^{\mathbf{L}}_E K^\bullet)$$

is an isomorphism. We may assume that $M = P$ is a differential graded E-module which has property (P). Since $K^\bullet$ defines a compact object, we reduce using Lemma 0.3 to the case where P has a finite filtration whose graded pieces are direct sums of $E[k]$. Again using compactness we reduce to the case $P = E[k]$. The assumption on $K^\bullet$ is that the result holds for these. □