# A RESOLVENT CRITERION FOR CONVERGENCE OF DYNAMICAL STATIONARY FLUCTUATIONS

LABBÉ, CYRIL, LASLIER, BENOÎT, TONINELLI, FABIO, ZAMBOTTI, LORENZO

*Editorial note.* Only selected theorem counted; complete mathematical body retained including the source figure. The figure is a preserved vector crop of publisher PDF page 41; extraction coordinates retained. Unavailable STIX wedge/vee glyph font uses standard semantic wedge/vee symbols. Original assumption counter uses the amsart theorem counter.

*Allowed standard prerequisites (editorial).*

- Standard functional analysis, measure theory and the author-cited classical theorems. Standard prior results stay cited; complete source-local proofs and definitions retained.

Contents

1. Introduction 1
2. A black box theorem 3
3. The zero-range process 9
4. Pair of reflected interfaces 19
5. $\nabla\phi$ interfaces with convex potential in one dimension 28
References 29

## 1. Introduction

This work deals with scaling limits of stationary fluctuations of Markov processes. We typically think of statistical physics models such as particle systems (exclusion process, zero-range process), evolving interfaces ($\nabla\phi$ interfaces, height functions) or dimer models, and the limiting process is usually given by a stochastic PDE. There exists by now a large literature on this topic. However, establishing rigorously such a convergence result often requires a delicate estimate, the so-called Boltzmann–Gibbs principle, that allows to replace space-time averages of discrete observables by a function of the local density up to some negligible error. We refer to [17, Section 11.1] for more details on the Boltzmann–Gibbs principle in a reversible setting, [6] in a non-reversible setting and also [14] for an extension called a second order Boltzmann–Gibbs principle; let us mention also that the Boltzmann–Gibbs principle, together with the relative entropy method, plays an important role for the proof of convergence of *non-equilibrium fluctuations* in some interacting particle systems [16]. Moreover, there are some interesting models from statistical physics for which the scaling limit of the stationary fluctuations is still out of reach of existing methods.

In this article, we present a general theorem, dubbed "black box theorem" in the sequel, that ensures convergence of a sequence of stationary Markov processes towards some limiting stationary Markov process, provided a set of five assumptions are fulfilled. Having in mind the models of statistical physics alluded to above, this theorem is designed to circumvent the use of the Boltzmann–Gibbs principle. As we will see, this theorem not only provides simpler proof for existing convergence results (see the example of the zero-range process below) but it also allows to prove new convergence results (see the pair of ordered interfaces below).

Our black box theorem relies on a change of perspective on how one characterizes the limiting process. In classical methods, the characterization of the limit goes through a martingale problem and the Boltzmann–Gibbs principle is used for computing the quadratic variation of the scaling limits of the discrete martingales. In our approach, we instead require a characterization of the limit through an identity satisfied by its resolvents, see Assumption 2.1 below.

The remaining four assumptions concern the sequence of stationary Markov processes. We naturally require the sequence of invariant measures to converge to the invariant measure of the limit, Assumption 2.2, and we ask the sequence of Markov processes to be tight, Assumption 2.3. The last two assumptions deal specifically with the resolvents of the sequence of Markov processes. In Assumption 2.4, we require the

*Date*: doi:10.5802/ahl.277; CC BY 4.0.

sequence of resolvents to be equicontinuous in order to extract converging subsequences. Finally, Assumption 2.5 imposes that the sequence of resolvents are characterized by an identity that is asymptotically the same as the identity satisfied by the resolvents of the limit.

Our black box theorem finds its origin in an article of the fourth author [31] on the convergence of the stationary fluctuations of a $\nabla\phi$ interface model with reflection towards the solution of a reflected SPDE introduced by Nualart and Pardoux [26]. This convergence had been established by Funaki and Olla [11] using the Boltzmann–Gibbs principle: the proof in [31] allowed to circumvent this delicate step. Later this technique was turned into a "black box" result in [32], however this approach was restricted to continuous-in-time approximations, while a paper of the first author [9] proved that the convergence result of [31] was true also for discrete approximations. The present work generalises both techniques to a much more general setting that encompasses both discrete and continuous state-spaces.

In the present paper, we apply our general result to three models of statistical physics. We first consider in Section 3 the one-dimensional zero-range process on a segment and show how our approach circumvents the use of the delicate Boltzmann–Gibbs principle, and simply requires some standard estimates under the invariant measure of the discrete dynamics. We emphasize that the result for the zero-range process is not new (and that it holds under weaker conditions on the transition rates than those we assume below) and that we present it here for pedagogical reasons, that is, to first illustrate our method in a simple context.

The second and main application of our general result concerns an evolving pair of ordered lattice paths $\mathtt{v} \geq \mathtt{w}$ on $[0, 2N]$, that are bound to the height 0 at 0 and $2N$ and evolve according to a corner-flip dynamics: the corners of $\mathtt{v}$ and $\mathtt{w}$ flip at some prescribed rates, except if the flip breaks the ordering of the two interfaces. This last constraint gives rise to a pair of *reflection measures* in the evolution equations: one that pushes up $\mathtt{v}$ and one that pushes down $\mathtt{w}$. Note that this process is a particular instance of a model introduced previously in [19, 28] in the context of dynamics of planar structures and tilings. At a higher level, our motivation to look at this process is the aim to understand the scaling limit of the evolution of *two-dimensional* random surfaces. The Markov chain on a pair of ordered lattice paths is then a toy model where the paths represent the level lines of the height function, and only two level lines are present. See the end of this section, as well as Remark 4.8, for a natural generalization with an arbitrary number $k$ of interacting level lines.





FIGURE 1. Example of a pair of reflected interfaces: $k = 3$ and $k = 9$ are *contact points*. At contact points, a reflection term prevents the jumps that would break the ordering to occur.

In Section 4, we will show that the stationary fluctuations of this model of reflected interfaces, under a diffusive rescaling, converge to the stationary solution of the following system of stochastic PDEs driven by independent space-time white noises $\xi^v$, $\xi^w$

$$\begin{cases} \partial_t v = \frac{1}{2}\partial_x^2 v + \xi^v + \frac{1}{\sqrt{2}}\eta, \quad x \in [0,1], \quad t \geq 0, \\ \partial_t w = \frac{1}{2}\partial_x^2 w + \xi^w - \frac{1}{\sqrt{2}}\eta, \quad x \in [0,1], \quad t \geq 0, \\ v \geq w, \quad \eta(dt,dx) \geq 0, \quad \int_{\mathbb{R}_+\times(0,1)} (v-w)(t,x)\eta(dt,dx) = 0. \end{cases} \tag{1}$$

In these equations, $\eta$ is a random measure on $\mathbb{R}_+ \times (0,1)$, carried by the space-time points where $v$ and $w$ meet, and whose role is to preserve the ordering $v \geq w$. This system of stochastic PDEs is intimately related to the reflected stochastic heat equation introduced by Nualart and Pardoux [26], see Section 4.1. The main difficulty in the proof of the convergence of the discrete dynamics consists in showing that the two discrete reflection measures converge to the *same* reflection measure $\eta$ in the limit: this question, for a very similar model, was left open in [9]. Implementing the Boltzmann–Gibbs principle to prove such a statement seems quite delicate: instead, checking the assumptions necessary for our black box theorem to apply mainly boils down to obtaining some estimates under the invariant measure of the process.

Finally we state a convergence result for discrete-time approximations of $\nabla\phi$ interfaces (with or without a "hard wall constraint" that imposes positivity of $\phi$) with convex potential in one dimension in Section 5, although we do not work out all details. These models have relevance in statistical physics, as describing

interface pinning/wetting phenomena in two-dimensions, see for instance the classic reference [10] as well as the monograph [12].

Although the three examples presented above are reversible w.r.t. their invariant measures, we emphasize that our black box result requires neither reversibility, nor the validity of the so-called "gradient condition" [17].

There is a number of directions that could be investigated in the future. First, our proof of the convergence of an evolving pair of ordered interfaces could be extended to cover systems of $k$ ordered interfaces for some arbitrary but fixed value $k \geq 2$, see Remark 4.8 for more details. Second, a very challenging question would be to apply the black box theorem (or some generalization thereof) to prove the convergence of the fluctuations of the height function of the dimer model on the honeycomb graph (or rhombus tilings of the plane), under the "tower dynamic" introduced in [19, 28], to a two-dimensional stochastic heat equation with additive noise. Informally speaking, this tiling dynamics corresponds to a system of $k$ ordered interfaces with $k$ itself of order $N$. The exact form of the limit stochastic heat equation can be guessed from the hydrodynamic limit equation obtained by the second and third authors in [20, 21].

We emphasize that the rhombus-tiling tower-dynamic of [28] satisfies a certain "gradient condition", that comes together with a $L^1$ contraction property [20, Section 3]. This feature allows for instance to prove a hydrodynamic limit [20, 21], that is, the convergence of the height profile, under diffusive scaling, to the solution of a non-linear parabolic PDE. This is in contrast with the usual rhombus tilings "single-flip" Glauber dynamic, for which no "gradient condition" or exact contraction property holds, so that even proving that the relaxation time scales diffusively is highly non-trivial (see [7, 22] for progress in this direction).

Third, it would be interesting to apply our black-box result to non-reversible systems, in particular, weakly asymmetric particle systems whose fluctuations converge to the KPZ equation.

**Organization of the paper.** Section 2 contains our "black box" theorem that, under certain general assumptions, allows to obtain the scaling limit of the fluctuation process of a sequence of Markov chains. The rest of the work applies the theorem to concrete cases: the zero-range process on a segment (Section 3), the pair of reflected interfaces (Section 4) and 1-dimensional $\nabla\phi$ interfaces with convex potentials (Section 5). In the latter case, we do not work out the details.

## 2. A black box theorem

This section presents a general result that ensures convergence in law of a sequence of stationary Markov processes (typically on discrete state-spaces) towards some stationary Markov process (typically on a continuous state-space). It relies on five assumptions: the first assumption concerns the limit process and the four others concern the sequence of discrete processes.

**Assumption on the limit.** We consider a collection of probability measures $(\mathbb{P}_x)_{x\in E}$ on the Skorohod space $\mathbb{D}([0,\infty), E)$ of càdlàg functions from $\mathbb{R}_+$ to a separable[1] metric space $E$ satisfying:

(1) $(x, A) \to \mathbb{P}_x(A)$ is a kernel, i.e., for every $x \in E$, $A \mapsto \mathbb{P}_x(A)$ is a probability measure on $E$ and for every $A \in \mathcal{B}(E)$, $x \mapsto \mathbb{P}_x(A)$ is measurable,

(2) the canonical process $(u(t))_{t\geq 0}$ under $\mathbb{P}_x$ satisfies the Markov property: for all $x \in E$, $\mathbb{P}_x$-a.s.

$$\mathbb{E}_x[F((u(t+s), s \geq 0)) \mid \mathcal{F}_t] = \mathbb{E}_{u(t)}[F(u(s), s \geq 0)], \quad \forall\, t \geq 0,$$

for any bounded measurable map $F : \mathbb{D}([0,\infty), E) \to \mathbb{R}$, and where $(\mathcal{F}_t, t \geq 0)$ is the canonical filtration.

We assume that the hence defined Markov process admits a stationary (*not necessarily reversible*) probability measure $m$, which has full topological support on $E$.

For any bounded measurable function $f : E \to \mathbb{R}$ and any $t \geq 0$ we set

$$P_t f(x) := \mathbb{E}_x[f(u(t))], \quad x \in E.$$

By a density argument, $P_t$ can be extended into a bounded operator on $L^2(E, dm)$. We then assume that $(P_t, t \geq 0)$ is a strongly continuous semigroup of bounded operators on $L^2(E, dm)$. The resolvent operators $\mathcal{R}_\lambda$, $\lambda > 0$ are defined as

$$\mathcal{R}_\lambda f := \int_0^\infty e^{-\lambda t} P_t f dt, \quad f \in L^2(E, dm).$$

It is well-known that the law of the process $u$ is completely characterized by the knowledge of the action of the operators $\mathcal{R}_\lambda$, $\lambda > 0$ on a large enough set of functions, see Section 2.1 for more details.

[1] We do not require $E$ to be a Polish space since completeness is not necessary for our black box theorem to hold. In concrete examples, we will actually consider spaces $E$ that are not complete.

We can now spell out our assumption, we refer the reader to the discussion below the statement for some explanations.

**Assumption 2.1** (Characterization of the limit)**.** *There exists a collection of finite complex measures $\Sigma^\psi$ on $E$, indexed by $\psi \in \mathcal{C}$ where $\mathcal{C}$ is some set of bounded and continuous functions from $E$ to $\mathbb{C}$, such that the following holds.*

*For any $\lambda > 0$ and any bounded Lipschitz function $f : E \to \mathbb{R}$, we assume that $\mathcal{R}_\lambda f$ admits a unique bounded Lipschitz representative in $L^2(E, m)$; moreover, this representative is the unique bounded Lipschitz function $F : E \to \mathbb{R}$ that satisfies the identity*

$$-\int_E F(h)\Sigma^\psi(dh) + \lambda \int_E F(h)\psi(h)m(dh) = \int_E f(h)\psi(h)m(dh), \quad \forall\, \psi \in \mathcal{C}. \tag{2}$$

Let us give some explanations on this assumption. An alternative definition of the resolvent operator can be given in terms of the generator $\mathcal{L}$ of the process: $\mathcal{R}_\lambda = (\lambda - \mathcal{L})^{-1}$. Provided the class $\mathcal{C}$ is large enough, this leads to the weak characterization of $\mathcal{R}_\lambda f$ as the unique element such that

$$\langle (\lambda - \mathcal{L})\mathcal{R}_\lambda f, \psi\rangle = \langle f, \psi\rangle$$

for all $\psi \in \mathcal{C}$. Here $\langle \cdot, \cdot \rangle$ denotes the inner product of the Hilbert space $L^2(E, m)$. By duality, we obtain

$$\forall\, \psi \in \mathcal{C}\ \langle \mathcal{R}_\lambda f, (\lambda - \mathcal{L}^*)\psi\rangle = \langle f, \psi\rangle\,.$$

Comparing with equation (2), this is consistent if $\mathcal{L}^*\psi(h)m(dh) = \Sigma^\psi(dh)$. However we insist that in Assumption 2.1 the function $\psi$ does not need to be in the domain of $\mathcal{L}^*$ (or $\mathcal{L}$ in the reversible case) so that the identity might only be formal. For example, for the pair of reflected interfaces, one can think of $\mathcal{L}$ as having a Dirac weight on configurations where the interfaces meet at a point (the only situation where the dynamics sees the reflection). At a rigorous level, this translates as the fact that $\Sigma^\psi$ gives positive mass to these configurations despite them having $m$-mass 0.

In the reversible case, an important tool to establish a characterization such as Assumption 2.1 is the general theory of self-adjoint operator and Dirichlet forms. In particular [24, Theorem I.2.8] says that for any $f \in L^2(E, dm)$ and for any $\lambda > 0$, $\mathcal{R}_\lambda f$ is the unique element of the domain $\mathcal{D}(\mathcal{E})$ of the Dirichlet form such that

$$\mathcal{E}(\mathcal{R}_\lambda f, \psi) + \lambda \int_E \mathcal{R}_\lambda f(h)\psi(h)m(dh) = \int_E f(h)\psi(h)m(dh),$$

for all $\psi$ that lie in (a dense subset of) $\mathcal{D}(\mathcal{E})$. When it is possible to determine $\mathcal{D}(\mathcal{E})$ and to justify an integration by parts in the Dirichlet form, this provides a sufficient condition on the set $\mathcal{C}$, see the proof of Proposition 4.2 for an example of this method.

Let us illustrate the assumption on a concrete example. Take $u$ to be the solution of the stochastic heat equation in dimension 1

$$\partial_t u = \frac{1}{2}\partial_x^2 u + \xi, \quad x \in [0, 1], \quad t \geq 0,$$

endowed with Dirichlet b.c. where $\xi$ is a space-time white noise. In this example, $E$ can be taken to be $L^1([0, 1], dx)$ and $m$ the law of the Brownian bridge on $[0, 1]$. The above assumption is then satisfied with $\mathcal{C}$, the set of all maps $\psi_\varphi : E \to \mathbb{C}$ defined by $\psi_\varphi(h) = \exp(i\langle h, \varphi\rangle)$ where $i = \sqrt{-1}$, $\langle \cdot, \cdot \rangle$ is the usual inner product in $L^2([0, 1], dx)$ and $\varphi : [0, 1] \to \mathbb{R}$ is any $C^2$-function that vanishes at 0 and 1. Then

$$d\Sigma^{\psi_\varphi}(h) = \psi_\varphi(h)\Big(\frac{i}{2}\langle h, \varphi''\rangle - \frac{1}{2}\|\varphi\|^2_{L^2([0,1],dx)}\Big)m(dh).$$

Let us mention that the choice $E = L^1([0, 1], dx)$ is relatively arbitrary at this stage. However, for establishing the convergence of approximating processes, this choice will play a role.

**Assumptions on the sequence of approximating processes.** We are now given a sequence of Markov processes $u_N$ on some spaces $E_N$. Each process $u_N$ admits an invariant probability measure $m_N$ on $E_N$ which has full topological support. We let $\mathcal{R}^N_\lambda$ denote its resolvent operator.

**Assumption 2.2** (State-space)**.** *Each $E_N$ is a Borel subset of $E$, endowed with the trace metric and Borel sigma-algebra. Furthermore the distance function to $E_N$ is achieved: namely, there exists a map $\pi_N : E \to E_N$ such that for all $x \in E$*

$$d_E(x, \pi_N(x)) = \inf_{y \in E_N} d_E(x, y).$$

*After extending the measure $m_N$ from $E_N$ to $E$ by setting $m_N(E \backslash E_N) = 0$, $m_N$ converges weakly towards $m$ as $N \to \infty$.*

The map $\pi_N$ will be called a projection. Let us emphasize that we do *not* assume that this projection is unique nor that it is continuous. Important examples satisfying this assumption are:

(1) $E_N$ is a finite subset of $E$: in this case existence of $\pi_N$ holds but uniqueness typically fails,
(2) $E$ is a Hilbert space and $E_N$ is a closed convex subset (for instance, a finite-dimensional subspace): existence and uniqueness hold by the Projection Theorem,
(3) $E$ is a subset of a Banach space $B$ (endowed with the same distance/norm), and $E_N \subset E$ is a closed subset of a finite dimensional subspace of $B$: existence holds but uniqueness may fail. For instance, take $E$ to be the subset of all functions of the Banach space $L^1([0,1], dx)$ that are non-negative a.e., and let $E_N$ be the subset of all non-negative functions that are constant on $[k/N, (k+1)/N)$ for every $k \in \{0, \dots, N-1\}$.

Below, we identify three conditions on the sequence $u_N$ that ensure that it converges in law towards $u$.

**Assumption 2.3** (Tightness)**.** *There exists a Polish space $E^-$ in which $E$ is continuously embedded, and such that, if each $u_N$ starts from its invariant measure $m_N$, then the sequence of processes $(u_N)_N$ is tight in $\mathbb{D}([0,\infty), E^-)$.*

This assumption spares us from proving tightness in $\mathbb{D}([0,\infty), E)$. In the applications presented later on, $E^-$ will be a Sobolev space with a negative index and the proof of the tightness will follow from the Lyons–Zheng decomposition [23] for reversible Markov processes.

We turn to a regularity assumption on the sequence of resolvents that will allow us to extract converging subsequences. We denote by $\mathrm{Lip}(E_N)$ the set of all Lipschitz functions $f : E_N \to \mathbb{R}$, that is, the set of all maps $f$ such that

$$[f]_{\mathrm{Lip}(E_N)} := \sup_{x,y \in E_N} \frac{|f(x) - f(y)|}{d_E(x,y)} < \infty.$$

**Assumption 2.4 (Equicontinuity).** For any given $\lambda > 0$ and any given Lipschitz function $f : E_N \to \mathbb{R}$, we assume that $\mathcal{R}_\lambda^N f$ admits a unique bounded Lipschitz representative in $L^2(E_N, m_N)$, that we still denote $\mathcal{R}_\lambda^N f$ for convenience. Furthermore there exists a constant $C > 0$ such that for all $\lambda > 0$, all $N \geq 1$ and all $f \in \mathrm{Lip}(E_N)$

$$[\mathcal{R}_\lambda^N f]_{\mathrm{Lip}(E_N)} \leq C [f]_{\mathrm{Lip}(E_N)}.$$

This is the main place where specific properties of the approximating process are used. Indeed in the applications that we will present, this estimate will be derived by probabilistic means, through finding a coupling under which the distance between two copies of the process is a sub-martingale. The choice of the underlying space $E$ plays an important role here: for instance, for some particle systems, the difference of the height functions of two evolving configurations decays stochastically in $L^1$ but not in $L^2$.

Finally, we spell out an assumption that can be viewed as an approximation of the characterization of $u$.

**Assumption 2.5** (Discrete characterization and convergence)**.** *For any $\psi \in \mathcal{C}$, there exists a finite complex measure $\Sigma_N^\psi$ on $E_N$ such that the following holds. For any $\lambda > 0$ and any $f \in L^2(E_N, dm_N)$, $F := \mathcal{R}_\lambda^N f$ satisfies the identity*

$$-\int_{E_N} F(h) \Sigma_N^\psi(dh) + \lambda \int_{E_N} F(h)\psi(h) m_N(dh) = \int_{E_N} f(h)\psi(h) m_N(dh). \tag{3}$$

*In addition, after extending $\Sigma_N^\psi$ from $E_N$ to $E$ by setting $\Sigma_N^\psi(E \backslash E_N) = 0$, the finite complex measure $\Sigma_N^\psi$ converges weakly to $\Sigma^\psi$ as $N \to \infty$.*

In the particular case where $E_N$ is a finite set and provided $\mathcal{C}$ is dense in $L^2(E_N, dm_N)$, equation (3) is always verified with $\Sigma_N^\psi(dh) := \mathcal{L}_N^* \psi(h) m_N(dh)$, where $\mathcal{L}_N^*$ is the adjoint w.r.t. $m_N$ of the generator $\mathcal{L}_N$ of the process. Indeed in that case the discussion after Assumption 2.1 is always rigorous since it only involves finite dimensional vector spaces. The core of the above assumption is therefore the convergence of the measure $\Sigma_N^\psi$ to $\Sigma^\psi$, or equivalently of the typically well defined $\mathcal{L}_N^* \psi(h)$ to the maybe only formal $\mathcal{L}^* \psi(h)$.

As one can guess, we will pass to the limit on identity (3) along some converging subsequence of resolvents (that can be extracted by Assumption 2.4) and we will identify limit points using Assumption 2.1.

We now have all the ingredients at hand to state our main result.

**Theorem 2.6.** *Let $u_N$ start from its invariant measure $m_N$ for every $N \geq 1$. Given Assumptions 2.1, 2.2, 2.3, 2.4 and 2.5, the sequence $(u_N)_N$ converges in law in $\mathbb{D}([0,\infty), E^-)$ to the process $u$ starting from its invariant measure $m$.*

Let us mention that in practice the convergence can be strengthened: if the tightness (Assumption 2.3) is established using moment bounds, and if one can establish uniform in $N$ moment bounds under the invariant measure $m_N$, then interpolation inequalities allow to lift the tightness in $\mathbb{D}([0,\infty), F)$ where $F$ is a space that interpolates between $E^-$ and $E$. We refer to [8, Lemma 5.2] for an application of this idea.

The rest of this section is devoted to the proof of this result.

2.1. **Reduction to an identity on the resolvents.** Let $\mathbb{P}^N_{m_N}$ be the law of $u_N$ starting from $m_N$ and let $\mathbb{P}$ be the law of $u$ starting from $m$. From the tightness Assumption 2.3, all we need to prove is that any limit of a converging subsequence of $(\mathbb{P}^N_{m_N})_N$ matches with $\mathbb{P}$.

First of all, observe that the restriction to $E$ of any function $f \in \mathrm{Lip}(E^-)$ lies in $\mathrm{Lip}(E)$. Indeed, we assumed that $E$ is continuously embedded in $E^-$, and therefore the bounded linear embedding map from $E$ to $E^-$ is Lipschitz continuous.

Below, all functions $f \in \mathrm{Lip}(E^-)$ will be evaluated on $E$: to alleviate the notation we will write $f$ for $f_{|E}$, and similarly with $E_N$.

Having this in mind, assume that for any $n \geq 1$, any bounded functions $f_0, f_1, \ldots, f_n \in \mathrm{Lip}(E^-)$ and any $\lambda_1, \ldots, \lambda_n > 0$, the following convergence holds

$$\int_{E_N} f_0(u_0)\mathcal{R}^N_{\lambda_1}\Big(f_1\mathcal{R}^N_{\lambda_2}\Big(\ldots f_{n-1}\mathcal{R}^N_{\lambda_n} f_n\Big)\ldots\Big)(u_0)m_N(du_0) \xrightarrow[N\to\infty]{} \int_E f_0(u_0)\mathcal{R}_{\lambda_1}\Big(f_1\mathcal{R}_{\lambda_2}\Big(\ldots f_{n-1}\mathcal{R}_{\lambda_n} f_n\Big)\ldots\Big)(u_0)m(du_0). \tag{4}$$

With this assumption at hand, let us prove that any limit of a converging subsequence of $(\mathbb{P}^N_{m_N})_N$ matches with $\mathbb{P}$. Let $P$ be the limit of a converging subsequence $(\mathbb{P}_{N_i})_i$. A monotone class argument yields $P = \mathbb{P}$ provided $P(u(t_1) \in A_1; \ldots; u(t_1 + \cdots + t_n) \in A_n) = \mathbb{P}(u(t_1) \in A_1; \ldots; u(t_1 + \cdots + t_n) \in A_n)$ for any collection $A_1, \ldots, A_n$ of closed sets of $E^-$, any collection $t_1, \ldots, t_n \geq 0$ and any integer $n \geq 1$. Since the indicator of any closed set of $E^-$ is the bounded pointwise limit of a sequence of bounded Lipschitz functions on $E^-$, it suffices to show that

$$\mathbb{E}_P\Big[f_0(u_0)\prod_{j=1}^n f_j(u_{t_1+\cdots+t_j})\Big] = \mathbb{E}_{\mathbb{P}}\Big[f_0(u_0)\prod_{j=1}^n f_j(u_{t_1+\cdots+t_j})\Big], \tag{5}$$

holds for all $n$ and bounded $f_0, f_1, \ldots, f_n \in \mathrm{Lip}(E^-)$. By the uniqueness of the Laplace transform of finite measures on $\mathbb{R}^n_+$, it boils down to proving[2]

$$\int_{(0,\infty)^n} e^{-\sum_{j=1}^n \lambda_j t_j}\mathbb{E}_P\Big[f_0(u_0)\prod_{j=1}^n f_j(u_{t_1+\cdots+t_j})\Big]dt_1\ldots dt_n = \int_{(0,\infty)^n} e^{-\sum_{j=1}^n \lambda_j t_j}\mathbb{E}_{\mathbb{P}}\Big[f_0(u_0)\prod_{j=1}^n f_j(u_{t_1+\cdots+t_j})\Big]dt_1\ldots dt_n, \tag{6}$$

for any $n \geq 1$, any $\lambda_1, \ldots, \lambda_n > 0$ and any bounded $f_0, f_1, \ldots, f_n \in \mathrm{Lip}(E^-)$.

The Markov property applied to the process $u_N$ shows that

$$\int_{E_N} f_0(u_0)\mathcal{R}^N_{\lambda_1}\Big(f_1\mathcal{R}^N_{\lambda_2}\Big(\ldots f_{n-1}\mathcal{R}^N_{\lambda_n} f_n\Big)\ldots\Big)(u_0)m_N(du_0) = \int_{(0,\infty)^n} e^{-\sum_j \lambda_j t_j}\mathbb{E}_{\mathbb{P}^N_{m_N}}\Big[f_0(u^N_0)\prod_{j=1}^n f_j(u^N_{t_1+\cdots+t_j})\Big]dt_1\ldots dt_n. \tag{7}$$

The weak convergence of $\mathbb{P}_{N_i}$ towards $P$ ensures that the r.h.s. converges, along the subsequence $N_i$, to

$$\int_{(0,\infty)^n} e^{-\sum_j \lambda_j t_j}\mathbb{E}_P\Big[f_0(u_0)\prod_{j=1}^n f_j(u_{t_1+\cdots+t_j})\Big]dt_1\ldots dt_n. \tag{8}$$

On the other hand, the identity (4) ensures that the l.h.s. of (7) converges to

$$\int_E f_0(u_0)\mathcal{R}_{\lambda_1}\Big(f_1\mathcal{R}_{\lambda_2}\Big(\ldots f_{n-1}\mathcal{R}_{\lambda_n} f_n\Big)\ldots\Big)m(du_0) = \int_{(0,\infty)^n} e^{-\sum_j \lambda_j t_j}\mathbb{E}_{\mathbb{P}}\Big[f_0(u_0)\prod_{j=1}^n f_j(u_{t_1+\cdots+t_j})\Big]dt_1\ldots dt_n,$$

as $N \to \infty$. Therefore, we have established (6), and the equality $\mathbb{P} = P$ follows. We have thus reduced the proof of Theorem 2.6 to the proof of (4).

[2]Of course $\mathbb{E}_P\Big[f_0(u_0)\prod_{j=1}^n f_j(u_{t_1+\cdots+t_j})\Big]dt_1\ldots dt_n$ is not a finite measure, but

$$e^{-\sum_{j=1}^n t_j\varepsilon}\mathbb{E}_P\Big[f_0(u_0)\prod_{j=1}^n f_j(u_{t_1+\cdots+t_j})\Big]dt_1\ldots dt_n$$

is a finite measure on $\mathbb{R}^n$ for any given $\varepsilon > 0$.

2.2. **Some technical tools.** In this paragraph, we collect some technical tools that will eventually allow us to extract converging subsequences from the sequence of discrete resolvents, and to pass to the limit on integrals involving these discrete resolvents against the discrete invariant measures. It turns out that our setting requires some care since continuous maps on $E_N$ cannot be canonically lifted into continuous maps on $E$. This is because in Assumption 2.2 we assumed existence of a projection $\pi_N$ from $E$ to $E_N$ without requiring its *continuity*. (It is worth keeping in mind that $\pi_N$ cannot be continuous when $E_N$ is a discrete subset of a continuous set $E$).

Interestingly, our assumptions imply that the amplitude of the possible jumps of the projection $\pi_N$ "vanish" as $N \to \infty$. More precisely, for every set $G \subset E$, we denote by $G^\varepsilon$ the open set of all the points in $E$ lying at a distance smaller than $\varepsilon$ from $G$. For all $\varepsilon > 0$, the weak convergence of $m_N$ to $m$ implies that

$$E = \varliminf_{N\to\infty} E_N^\varepsilon. \tag{9}$$

Indeed, for all $x \in E$ and all $\varepsilon > 0$, if we denote by $B(x,\varepsilon)$ the open ball in $E$ centered at $x$ and of radius $\varepsilon$, then we have

$$m\big(B(x,\varepsilon)\big) \le \varliminf_{N\to\infty} m_N\big(B(x,\varepsilon)\big).$$

Since $m$ has full topological support on $E$, we deduce that the l.h.s. is strictly positive so that, for all $N$ large enough $B(x,\varepsilon) \cap E_N \neq \emptyset$, thus ensuring that $x \in E_N^\varepsilon$ for all $N$ large enough. This implies that for all $x, y \in E$, there exists $N_0(x,y) \ge 1$ such that for all $N \ge N_0$, $d(\pi_N(x), \pi_N(y)) \le d(x,y) + 2\varepsilon$ where we write $d$ instead of $d_E$ to lighten notations.

We can now state a modification of Arzelà–Ascoli's theorem. Recall from Assumption 2.2 that $\pi_N(x)$ is an element of $E_N$ that minimizes the distance from $x$ to $E_N$.

**Theorem 2.7** (Modified Arzelà–Ascoli's theorem)**.** *Let $F_N : E_N \to \mathbb{R}, N \ge 1$ be a collection of uniformly bounded and equicontinuous maps, i.e. $\sup_{N\ge1} \|F_N\|_\infty < \infty$ and*

$$\omega(\delta) := \sup_{N\ge1} \big\{ |F_N(x) - F_N(y)| : x, y \in E_N,\ d(x,y) \le \delta \big\}, \tag{10}$$

*goes to* 0 *as* $\delta \downarrow 0$. *Then, there exists a bounded and uniformly continuous map* $F : E \to \mathbb{R}$, *as well as a subsequence* $(N_j)_{j\geq 1}$ *such that:*

(1) *For any compact set* $J \subset E$, $\sup_{x\in J\cap E_{N_j}} |F(x) - F_{N_j}(x)| \to 0$ *as* $j \to \infty$,

(2) $F_{N_j}(\pi_{N_j}x)$ *converges to* $F(x)$ *for all* $x \in E$.

**Remark 2.8.** The proof shows that the modulus of continuity of $F$ is bounded by $\omega(3\,\cdot\,)$, and can be adjusted to show that it is actually bounded by $\omega((1+\epsilon)\cdot\,)$ for any $\epsilon > 0$. In particular, if the $F_N$ are uniformly Lipschitz, so is $F$.

**Remark 2.9.** In the case where $E_N$ is a finite dimensional subspace of a Hilbert space $E$, the projection $\pi_N$ is 1-Lipschitz and then the classical Arzelà–Ascoli theorem can be applied to the sequence $F_N(\pi_N\,\cdot\,)$. In our more general setting, this is unfortunately not the case.

*Proof.* Let $(x_i)_{i\geq 1} \subset E$ be a dense sequence. By a diagonal argument, there exists a subsequence $N_j \to \infty$ such that, for every $i \geq 1$, $F_{N_j}(\pi_{N_j}x_i)$ converges to a limit, that we call $F(x_i)$. Necessarily, $\sup_i |F(x_i)| \leq \sup_N \|F_N\|_\infty < \infty$. Fix $i \neq \ell \geq 1$. From (9), we deduce that both $d(x_i, \pi_{N_j}x_i)$ and $d(x_\ell, \pi_{N_j}x_\ell)$ are smaller than $d(x_i, x_\ell)/2$ as soon as $j$ is large enough. Then, we have

$$|F(x_i) - F(x_\ell)| = \varlimsup_{j\to\infty} \big|F_{N_j}(\pi_{N_j}x_i) - F_{N_j}(\pi_{N_j}x_\ell)\big| \leq \omega\big(2\,d(x_i, x_\ell)\big). \tag{11}$$

Let $x \in E$. For each $k \geq 1$, let $i_k \geq 1$ be such that $x \in B(x_{i_k}, k^{-1})$. Using (11), it is then elementary to check that $F(x_{i_k}), k \geq 1$ is a Cauchy sequence. Let $F(x)$ be its limit: using (11), it is straightforward to check that this definition does not depend on the choice of the sequence $i_k$, and that it matches with the map $F$ already defined on $\{x_i, i \geq 1\}$. Using (11) once again, it is simple to obtain the bound $|F(x) - F(y)| \leq \omega(3\,d(x,y))$ for all $x, y \in E$. The function $F$ is therefore uniformly continuous on $E$.

Let $J$ be a compact set of $E$. For any $k \geq 1$, there exists $p_k \geq 1$ such that $\bigcup_{i\leq p_k} B(x_i, k^{-1})$ covers $J$. By (9), taking $j = j(k)$ large enough, for all $i \in \{1, \dots, p_k\}$ we have $x_i \in E_{N_j}^{1/k}$ together with

$$|F_{N_j}(\pi_{N_j}x_i) - F(x_i)| \leq \frac{1}{k}. \tag{12}$$

Let $x \in J$. There exists $i_k \leq p_k$ such that $d(x, x_{i_k}) \leq k^{-1}$. Then, for all $j$ large enough we get

$$\begin{aligned}\big|F(x) - F_{N_j}(\pi_{N_j}x)\big| &\leq |F(x) - F(x_{i_k})| + \big|F(x_{i_k}) - F_{N_j}(\pi_{N_j}x_{i_k})\big| \\ &\quad + \big|F_{N_j}(\pi_{N_j}x_{i_k}) - F_{N_j}(\pi_{N_j}x)\big| \\ &\leq \omega\big(3k^{-1}\big) + k^{-1} + \omega\big(2k^{-1} + d(x, \pi_{N_j}x)\big).\end{aligned}$$

If $x \in J \cap E_{N_j}$, then $x = \pi_{N_j}x$ and the first assertion of the theorem follows. Regarding the second assertion, fix $x \in E$ and observe that we can apply the previous arguments to $J = \{x\}$. By (9), we know that for all $j$ large enough $d(x, \pi_{N_j}x) \leq k^{-1}$ and therefore the previous bound ensures that $F_{N_j}(\pi_{N_j}x)$ converges to $F(x)$. This ends the proof of Theorem 2.7. □

We conclude this section by a criteria for "diagonal" convergence where both the measure and the integrand depend on $N$. The proof is very close to [29, Lemma 1].

**Lemma 2.10.** *Let* $\psi : E \to \mathbb{C}$ *be a bounded and continuous function. Let* $F_N : E_N \to \mathbb{R}$, $N \geq 1$ *be a sequence of uniformly bounded and equicontinuous maps in the sense of Proposition 2.7, and assume that* $F_N(\pi_N\cdot\,)$ *converges* $m$*-almost everywhere to some bounded continuous map* $F : E \to \mathbb{R}$. *Suppose that* $\rho_N$ *is a sequence of complex finite measures on each* $E_N$, *that we extend to* $E$ *by setting* $\rho_N(E\backslash E_N) = 0$, *and that converges weakly to some complex finite measure* $\rho$ *on* $E$. *Then* $\int_E F_N \circ \pi_N \psi \, d\rho_N \to \int F\psi \, d\rho$.

*Proof.* Without loss of generality, we can assume that $0 \leq F_N \leq 1$, $0 \leq \psi \leq 1$ and that the measures $\rho_N$ are real-valued and non-negative. Let $N_j, j \geq 1$ be a sequence going to $\infty$. By Theorem 2.7, we can extract a subsubsequence $F_{N_{i_j}}$ such that the two assertions of the theorem are satisfied: since the limit is continuous on $E$, it necessarily coincides with the function $F$. The tightness of the sequence $\rho_{N_{i_j}}$ implies that for every $n \geq 1$, there exists a compact set $J_n$ such that $\varlimsup_j \rho_{N_{i_j}}(J_n^c) \leq n^{-1}$. Using the fact that $\rho_{N_{i_j}}$ is supported by $E_{N_{i_j}}$, we write

$$\begin{aligned}&\int_E F_{N_{i_j}} \circ \pi_{N_{i_j}} \psi d\rho_{N_{i_j}} \\ &\qquad\qquad \leq \rho_{N_{i_j}}(J_n^c) + \int_{J_n} \Big(F_{N_{i_j}}(\pi_{N_{i_j}}\,\cdot\,) - F(\,\cdot\,)\Big)\psi d\rho_{N_{i_j}} + \int_{J_n} F\psi d\rho_{N_{i_j}}.\end{aligned} \tag{13}$$

The first assertion of the theorem and the boundedness of $\psi$ imply that

$$\overline{\lim_j} \int_{J_n} \Big| F_{N_{i_j}}(\pi_{N_{i_j}} \cdot) - F(\cdot) \Big| \psi d\rho_{N_{i_j}} = 0.$$

The weak convergence of $F\psi\rho_N$ towards $F\psi\rho$ ensures that

$$\overline{\lim_j} \int_{J_n} F\psi d\rho_{N_{i_j}} \le \int_{J_n} F\psi d\rho.$$

Therefore, we get

$$\overline{\lim_j} \int_E F_{N_{i_j}} \circ \pi_{N_{i_j}} \psi d\rho_{N_{i_j}} \le \frac{1}{n} + \int_E F\psi d\rho. \tag{14}$$

Passing to the limit on $n$, we obtain $\overline{\lim}_j \int F_{N_{i_j}} \circ \pi_{N_{i_j}} \psi d\rho_{N_{i_j}} \le \int F\psi d\rho$. Upon replacing $F_N$ by $1 - F_N$, one obtains $\underline{\lim}_j \int F_{N_{i_j}} \circ \pi_{N_{i_j}} \psi d\rho_{N_{i_j}} \ge \int F\psi d\rho$. This ends the proof of Lemma 2.10. □

2.3. **Convergence of the resolvents.** We come back to our original goal: the proof of (4). It boils down to the following lemma.

**Lemma 2.11.** *Let $g_N : E_N \to \mathbb{R}$, $N \ge 1$ be a sequence of uniformly bounded and uniformly Lipschitz maps, such that $g_N \circ \pi_N$ converges m-a.e. to some bounded Lipschitz function $g : E \to \mathbb{R}$. Then, for all $\lambda > 0$, the map $\mathcal{R}^N_\lambda g_N(\pi_N \cdot)$ converges pointwise to $\mathcal{R}_\lambda g(\cdot)$ as $N \to \infty$.*

*Proof.* Using Assumption 2.4, we know that the sequence $F_N(\cdot) = \mathcal{R}^N_\lambda g_N(\cdot)$, $N \ge 1$, is uniformly bounded and equicontinuous in the sense of Theorem 2.7. Furthermore Assumption 2.5 states that for all $\psi \in \mathcal{C}$

$$-\int_{E_N} F_N(h)\Sigma^\psi_N(dh) + \lambda \int_{E_N} F_N(h)\psi m_N(dh) = \int_{E_N} g_N(h)\psi(h)\, m_N(dh).$$

Let $F_{N_i}, N_i \ge 1$ be an arbitrary subsequence. By Theorem 2.7, there exists a bounded Lipschitz map $F$ on $E$ and a subsubsequence $F_{N_{i_j}}$ converging to $F$ in the sense of the statement. The weak convergences of $\Sigma^\psi_N$ and $m_N$ to $\Sigma^\psi$ and $m$ (guaranteed by Assumptions 2.5 and 2.2), together with the regularity properties of the integrands are sufficient to apply Lemma 2.10 along the subsubsequence $N_{i_j}$ and deduce that

$$-\int_E F(h)\Sigma^\psi(dh) + \lambda \int_E F(h)\psi(h)m(dh) = \int_E g(h)\psi(h)\, m(dh).$$

Since this holds for all $\psi \in \mathcal{C}$, Assumption 2.1 ensures that $F = \mathcal{R}_\lambda g$. Since the subsequence $N_i$ was arbitrary, we can conclude. □

A recursion based on Lemmas 2.10 and 2.11 then yields (4).

## 3. The zero-range process

For pedagogical reasons, we show first how our method works when applied to the one-dimensional zero-range process. In this case, the convergence of the fluctuation process is already known (see [17, Chapter 11] and references therein), which allows the reader to compare our new approach to the classical one. Note that we bypass entirely the need to prove the so-called "Boltzmann–Gibbs principle" ([17, Section 11.1]). The equicontinuity of the resolvents will follow from a model dependent but very soft argument, while the "replacement" part of the proof, Assumption 2.5, uses technical but routine local CLT results.

3.1. **Model and notations.** A configuration $n$ is a map from the lattice $\{1, \dots, N\}$ to $\mathbb{N}$ and we interpret $n(i)$ as a number of particles at site $i$. The jump rate of the particles is given by a function $\tau : \mathbb{N} \to \mathbb{R}_+$ on which we assume: (1) $\tau(0) = 0$, (2) $\tau$ is a non-decreasing function, (3) $\tau$ grows at most polynomially at infinity. (The second hypothesis implies attractiveness).

The dynamics is as follows: at rate $N^2\tau(n(i))$ a Poisson clock rings at site $i$ and when this happens a particle jumps right or left by one step, each with probability $1/2$, except for $i = 1$ (resp. $i = N$) in which case the jump to the left (resp. to the right) is not allowed. In other words, at rate $N^2\tau(n(i))/2$, $(n(i), n(i+1)) \to$

$(n(i) - 1, n(i+1)+1)$ whenever $i \in \{1, \dots, N-1\}$ and at the same rate $(n(i), n(i-1)) \to (n(i)-1, n(i-1)+1)$ whenever $i \in \{2, \dots, N\}$.

Define the probability measure

$$\nu_a(k) := \frac{1}{Z_a} \frac{a^k}{\prod_{i=1}^k \tau(i)}, \quad k \in \mathbb{N}, \quad Z_a := \sum_{j \geq 0} \frac{a^j}{\prod_{i=1}^j \tau(i)}.$$

Let $a^* := \sup\{a > 0 : Z_a < \infty\}$. Depending on the growth rate of the function $\tau$, this quantity is either finite or infinite. We introduce $R(a) := \sum_k k\nu_a(k)$, the first moment of $\nu_a$, and $R^* := \lim_{a \uparrow a^*} R(a)$. We then let $\Phi : (0, R^*) \to (0, a^*)$ be the reciprocal of $R$: for any parameter $\overline{n} \in (0, R^*)$, the measure $\nu_{\Phi(\overline{n})}$ has mean $\overline{n}$. Given some $\overline{n} \in (0, R^*)$, the above dynamics, restricted to the set $\Omega_{N,\overline{n}}$ of configurations with $\lfloor N\overline{n} \rfloor$ particles, admits a unique invariant (and reversible) probability measure

$$\pi_{N,\overline{n}}(n) = \frac{C(\overline{n}, N)}{\prod_{i=1}^N \prod_{j=1}^{n(i)} \tau(j)}, \quad \forall\, n = (n(1), \dots, n(N)) \in \Omega_{N,\overline{n}}.$$

This measure is nothing but the product measure of the $\nu_a$ conditioned to $\Omega_{N,\overline{n}}$, for any arbitrary parameter $a \in (0, a^*)$. Let us point out however that a "natural" choice of parameter $a$ would be $\Phi(\overline{n})$.

From now on, $\overline{n} \in (0, R^*)$ is fixed but arbitrary.

Given an evolving configuration $(n_N(t,i), t \geq 0, i \in \{1, \dots, N\})$, we introduce the evolving height function as follows

$$u_N(t,x) := \frac{1}{\sqrt{N}} \sum_{k=1}^{\lfloor xN \rfloor} \big(n_N(t,k) - \overline{n}\big), \quad x \in [0,1]. \tag{15}$$

It turns out that it is convenient to work on the Banach space $E = L^1([0,1], dx)$.

Since the mapping $n_N \to u_N$ is bijective, $(u_N(t), t \geq 0)$ is a Markov process taking values in $E$. We let $\mathcal{L}_N$ be its generator:

$$\begin{aligned} \mathcal{L}_N f(u_N) = \sum_{k=1}^{N-1} \frac{N^2 \tau(n_N(k))}{2} \Big( f\Big(u_N^{k,k+1}\Big) - f(u) \Big) \\ + \sum_{k=2}^{N} \frac{N^2 \tau(n_N(k))}{2} \Big( f\Big(u_N^{k,k-1}\Big) - f(u) \Big), \end{aligned} \tag{16}$$

where $u_N^{k,k\pm1}$ is obtained from $u_N$ by removing a particle at site $k$ and adding one at site $k \pm 1$.

We also let $m_N$ be its invariant probability measure, and $\mathcal{E}_N$ the associated Dirichlet form under the invariant measure $m_N$. Note that $u_N$ actually takes values in a finite set $E_N \subset E$.

For later purposes, we derive some scaling limits under the invariant measure. These limits depend on a couple of parameters defined as follows. Set $a := \Phi(\overline{n})$ and

$$\begin{aligned} \overline{\tau} = \mathrm{E}_{\nu_a}[\tau(n)] &= \sum_k \tau(k)\nu_a(k), \qquad \rho = \mathrm{Cov}_{\nu_a}(n, \tau(n)) &= \sum_k (k - \overline{n})(\tau(k) - \overline{\tau})\nu_a(k), \\ \alpha = \mathrm{Var}_{\nu_a}(n) &= \sum_k (k - \overline{n})^2 \nu_a(k), \; \gamma = \mathrm{Var}_{\nu_a}(\tau(n)) &= \sum_k (\tau(k) - \overline{\tau})^2 \nu_a(k). \end{aligned}$$

Elementary computations show that

$$\overline{\tau} = \rho = a.$$

Given a configuration $n$ we introduce the height function associated to the field of jump rates:

$$q_N(x) := \frac{1}{\sqrt{N}} \sum_{k=1}^{\lfloor xN \rfloor} \big(\tau(n(k)) - \overline{\tau}\big), \quad x \in [0,1],$$

as well as the static counterpart of (15)

$$h_N(x) := \frac{1}{\sqrt{N}} \sum_{k=1}^{\lfloor xN \rfloor} (n(k) - \overline{n}), \quad x \in [0,1]. \tag{17}$$

**Lemma 3.1.** *The pair $(h_N, q_N)$, under the probability measure $m_N$, converges in law to a centered Gaussian process $(\beta(x), \zeta(x))_{0\le x\le 1}$ whose covariance is characterized by*

$$\begin{aligned}\mathrm{Cov}(\beta(s),\beta(t)) &= \alpha\min(s,t)(1-\max(s,t)),\\ \mathrm{Cov}(\zeta(s),\zeta(t)) &= \gamma\min(s,t) - \frac{\rho^2}{\alpha}st,\\ \mathrm{Cov}(\beta(s),\zeta(t)) &= \rho\min(s,t)(1-\max(s,t)).\end{aligned}$$

*In addition the following bounds hold: for all $p \ge 1$*

$$\begin{aligned}\sup_N m_N\left[\sup_{x\in[0,1]}|h_N(x)|^p\right] &< \infty,\\ \sup_N m_N\left[\sup_{x\in[0,1]}|q_N(x)|^p\right] &< \infty,\\ \sup_N m_N[\tau(n(1))^p] &< \infty.\end{aligned}$$

This result implies that $\frac{1}{\sqrt{\alpha}}\beta$ is a standard Brownian bridge on $[0,1]$, and that $\zeta - \frac{\rho}{\alpha}\beta$ is a centered Gaussian process independent of $\beta$.

*Proof.* To prove the convergence of $(h_N, q_N)$ under $m_N$, one first considers the product law $\mu_N := \otimes_{k=1}^N \nu_a$ with $a = \Phi(\overline{n})$. It is standard to prove that under[3] $\mu_N$, the pair $(h_N, q_N)$ converges towards a centered Gaussian process $(B, Z)$ with covariance

$$\begin{aligned}\mathrm{Cov}(B(s),B(t)) &= \alpha\min(s,t),\\ \mathrm{Cov}(Z(s),Z(t)) &= \gamma\min(s,t),\\ \mathrm{Cov}(B(s),Z(t)) &= \rho\min(s,t).\end{aligned}$$

($B$ and $Z$ are scaled and correlated Brownian motions). At this point, we observe that $Z = \frac{\rho}{\alpha}B + Z'$ where $Z'$ is a centered Gaussian process independent of $B$. Consequently if we let $(\beta,\zeta)$ be the centered Gaussian process obtained by conditioning $(B,Z)$ to $B(1)=0$, then $\beta$ is a scaled Brownian Bridge and $\zeta = \frac{\rho}{\alpha}\beta + Z'$, with $\beta$ independent of $Z'$. The covariance structure of the statement follows.

We now concentrate on the proof of the convergence of $h_N$ under $m_N$ towards $\beta$: the extension to the convergence of the pair $(h_N, q_N)$ under $m_N$ is relatively straightforward. By [27, Theorem 1, Chapter VII] a local limit theorem holds: for all $x \in (0,1]$, if we let $k_N := \lfloor xN \rfloor$ then

$$\sup_{y\in\frac{\mathbb{N}-k_N\overline{n}}{\sqrt{N}}}\left|\sqrt{N}\mu_N(h_N(k_N/N)=y) - g_{\alpha x}(y)\right| \longrightarrow 0, \quad N\longrightarrow\infty, \tag{18}$$

where $g_{\sigma^2}$ denotes the density of the centered Gaussian of variance $\sigma^2$. The independence and stationarity of the increments allows to extend this LLT to finite dimensional marginals.

Following an observation made earlier in this subsection and given the definition of $h_N$, we have

$$m_N = \mu_N(\,\cdot\mid h_N(1)=\delta_N), \quad \delta_N := \frac{1}{\sqrt{N}}(\lfloor N\overline{n}\rfloor - N\overline{n}).$$

This allows to deduce the following uniform in $N$ absolute continuity result. There exists a constant $C>0$ such that for all $N$ large enough and all non-negative and measurable map $f:\mathbb{R}^{k_N}\to\mathbb{R}_+$

$$m_N\Big[f(h_N(1/N),\dots,h_N(k_N/N))\Big] \le C\mu_N\Big[f(h_N(1/N),\dots,h_N(k_N/N))\Big] \tag{19}$$

where $k_N := \lceil N/2\rceil$. Indeed, $C$ can simply be taken to be the supremum over all $N$ large enough of

$$\sup_{y\in\frac{\mathbb{N}-k_N\overline{n}}{\sqrt{N}}}\frac{\mu_N(h_N(1)-h_N(k_N/N)=-y)}{\mu_N(h_N(1)=0)} = \sup_{y\in\frac{\mathbb{N}-k_N\overline{n}}{\sqrt{N}}}\frac{\mu_N(h_N(1-k_N/N)=-y)}{\mu_N(h_N(1)=0)},$$

and this quantity is finite thanks to (18).

Let us now establish that the sequence $(h_N)_N$ is tight. Actually, our proof will also establish the moment bounds of the statement on the expectation of the suprema of $h_N$. To that end, we establish an estimate on the Hölder semi-norm of $u_N$: for any $\beta\in(0,1/2)$ and $p\ge 1$

$$\limsup_{N\ge1}\sup_{k\neq\ell\in\{1,\dots,N\}} m_N\left[\left(\frac{|h_N(k/N)-h_N(\ell/N)|}{|\frac{k}{N}-\frac{\ell}{N}|^\beta}\right)^p\right] < \infty. \tag{20}$$

[3]With minor abuse of notation, we let $\mu_N$ denote also the law of $(h_N, q_N)$ when $n$ has law $\mu_N$.

By symmetry and using the triangle inequality, we can assume that $k, \ell \leq N/2$. From the uniform absolute continuity (19), it suffices to establish (20) under $\mu_N$ rather than $m_N$. This estimate follows from standard techniques, using the fact that under $\nu_a$, $n - \bar{n}$ admits moments of all orders.

Let us now deduce the convergence of the finite dimensional marginals of $h_N$ under $m_N$ towards those of $\beta$. For simplicity, let us concentrate on one-dimensional marginals. Fix $a < b$ and $x \in (0,1)$, and set $k_N := \lfloor xN \rfloor$. We compute

$$\begin{aligned} m_N\big(a < h_N(k_N/N) < b\big) & \\ &= \sum_{y \in \frac{\mathbb{N} - k_N \bar{n}}{\sqrt{N}} : a<y<b} m_N(h_N(k_N/N) = y) \\ &= \sum_{y \in \frac{\mathbb{N} - k_N \bar{n}}{\sqrt{N}} : a<y<b} \frac{\mu_N(h_N(k_N/N) = y)\mu_N(h_N(1 - k_N/N) = -y + \delta_N)}{\mu_N(h_N(1) = \delta_N)}. \end{aligned}$$

Now observe that

$$\inf_{y \in (a,b)} g_{\alpha x}(y) > 0, \quad \inf_{y \in (a,b)} g_{\alpha(1-x)}(-y + \delta_N) > 0, \quad g_\alpha(\delta_N) \longrightarrow g_\alpha(0) > 0.$$

Consequently the LLT (18) allows to write (with $o(1)$ being uniform over all $y$ in the sum)

$$\begin{aligned} m_N\big(a < h_N(k_N/N) < b\big) &= \sum_{y \in \frac{\mathbb{N} - k_N \bar{n}}{\sqrt{N}} : a<y<b} \frac{1}{\sqrt{N}} \frac{g_{\alpha x}(y) g_{\alpha(1-x)}(-y + \delta_N)}{g_\alpha(\delta_N)} (1 + o(1)) \\ &\longrightarrow \int_{a<y<b} \frac{g_{\alpha x}(y) g_{\alpha(1-x)}(-y)}{g_\alpha(0)} dy = \mathbb{P}(a < \beta(x) < b). \end{aligned}$$

Combined with the bound (20) already established, this is sufficient to deduce that the law of $h_N(x)$ under $m_N$ converges to that of $\beta(x)$.

Let us finally establish the bound on the moments of $\tau(n(1))$. From the absolute continuity (19), it is sufficient to prove

$$\limsup_N \nu_a[\tau(n(1))^p] < \infty,$$

and this follows from the fact that $\tau$ grows at most polynomially at infinity. □

Note that Lemma 3.1 implies that Assumption 2.2 holds (recall item (3) in the discussion after the formulation of that assumption).

3.2. **The SPDE.** Consider the solution $u$ of

$$\begin{cases} \partial_t u = c\partial_x^2 u + \sigma\xi, \quad x \in [0,1], \quad t \geq 0, \\ u(t = 0, \cdot) = u_0(\cdot), \end{cases} \tag{21}$$

endowed with Dirichlet b.c., where $c, \sigma > 0$ are some constants, $\xi$ is a gaussian space-time white noise and $u_0$ is some initial condition in $E = L^1([0,1], dx)$. (We refer to Section 4.1 for the precise meaning given to such an equation).

The process $u$ admits a unique invariant and reversible measure $m_{c,\sigma}$ which is the law of a scaled Brownian bridge (pinned at 0 at $x \in \{0,1\}$). More precisely, let $Y(x), x \in [0,1]$ be a standard Brownian bridge then $m_{c,\sigma}$ is the law of $\sqrt{\frac{\sigma^2}{2c}} Y$.

Since $u$ is a Feller process, it is associated to a strongly continuous semigroup $P_t$ in $L^2(E,m)$, the latter being the space of all maps $F : E \to \mathbb{R}$ that are square integrable against $m$. The general theory ensures that $P_t$ gives rise to a generator $\mathcal{L}$, a Dirichlet form $\mathcal{E}$ and resolvents $\mathcal{R}_\lambda$, $\lambda > 0$.

Our next result shows that Assumption 2.1 is satisfied for the process $u(t, \cdot)$. The class $\mathcal{C}$ is taken to be the set of all functions $\psi_\varphi(u) := \exp(i\langle u, \varphi \rangle)$, where $\varphi : [0,1] \to \mathbb{R}$ is a $C^2$ function compactly supported in $(0,1)$, and where $\langle \cdot, \cdot \rangle$ is the inner product in $L^2([0,1], dx)$.

**Proposition 3.2.** *Let $f \in L^2(E, dm)$ be a Lipschitz function from $E$ to $\mathbb{R}$. For any $\lambda > 0$, $\mathcal{R}_\lambda f$ is the unique element of $L^2(E, dm)$ such that for any $C^2$ function $\varphi : [0,1] \to \mathbb{R}$ with compact support in $(0,1)$, the following holds:*

$$\int \mathcal{R}_\lambda f(h) \psi_\varphi(h) \left( \lambda + \frac{\sigma^2}{2} \|\varphi\|_{L^2}^2 - ic\langle h, \varphi'' \rangle \right) m(dh) = \int f(h) \psi_\varphi(h) m(dh). \tag{22}$$

*As a consequence Assumption 2.1 is satisfied with the class $\mathcal{C}$ and the measures*

$$\Sigma^{\psi_\varphi}(dh) = \psi_\varphi(h)\Big(ic\langle h, \varphi''\rangle - \frac{\sigma^2}{2}\|\varphi\|_{L^2}^2\Big)dm(h), \quad \psi_\varphi \in \mathcal{C}. \tag{23}$$

Such a result is certainly not new, but for the sake of completeness we provide a proof. Let us emphasize that the uniqueness part relies on elementary facts on finite-dimensional Ornstein–Uhlenbeck processes.

*Proof. Step 1: Extension to more general $\varphi$.* Assume that some map $F \in L^2(E, dm)$ satisfies for all $C^2$ function $\varphi$ with compact support in $(0,1)$

$$\int F(h)\psi_\varphi(h)\Big(\lambda + \frac{\sigma^2}{2}\|\varphi\|_{L^2}^2 - ic\langle h, \varphi''\rangle\Big)m(dh) = \int f(h)\psi_\varphi(h)m(dh). \tag{24}$$

Let us show that it remains true for all $C^2$ function $\varphi$ that is not necessarily compactly supported in $(0,1)$ but only vanishes at the boundaries. First of all, it is not hard to prove the following moment bound on the Brownian bridge: for any given $p \geq 1$

$$m\left[\sup_{x\in[0,2/n]\cup[1-2/n,1]} |h(x)|^p\right] \lesssim (1/n)^{p/2}. \tag{25}$$

uniformly over all $n \geq 1$. Now let $\chi_n : [0,1] \to [0,1]$ be a smooth function that equals 1 on $[2/n, 1-2/n]$ and 0 outside $[1/n, 1-1/n]$ and such that there exists $C > 0$ such that for all $n \geq 1$

$$\sup_x |\chi_n'(x)| \leq Cn, \quad \sup_x |\chi_n''(x)| \leq Cn^2.$$

Set $\varphi_n(x) := \varphi(x)\chi_n(x)$. Then (24) evaluated at $\varphi_n$ converges to the same equation with $\varphi$. The only delicate term is $\langle u, \varphi''\rangle$, let us provide the details. Since $\varphi - \varphi_n$ vanishes outside $I_n := [0, 2/n] \cup [1-2/n, 1]$ we find

$$\begin{aligned}\int |F(h)\psi_\varphi(h)\langle h, (\varphi'' - \varphi_n'')\rangle| m(dh) \\ \leq \left(\int F(h)^2 m(dh)\right)^{1/2} m\left[\sup_{x\in I_n}|h(x)|^2\right]^{1/2} \\ \int_{[0,2/n]\cup[1-2/n,1]} |\varphi''(1-\chi_n) + 2\varphi'\chi_n' + \varphi\chi_n''|.\end{aligned}$$

By (25), the product of the first two terms goes to 0 as $n \to \infty$. Since $\varphi$ vanishes at 0 and 1, it is of order $1/n$ on $I_n$, and the integral can be bounded by a term of order 1 uniformly over all $n \geq 1$.

*Step 2: Existence.* The general theory [24, Theorem I.2.8] ensures that $\mathcal{R}_\lambda f$ is the only $F \in \mathcal{D}(\mathcal{E})$ satisfying

$$\mathcal{E}(F, g) + \lambda \int F(h)g(h)m(dh) = \int f(h)g(h)m(dh), \tag{26}$$

for all $g \in \mathcal{D}(\mathcal{E})$, the latter being the domain of the Dirichlet form. The general theory again ensures that for any $g \in \mathcal{D}(\mathcal{L})$, the domain of the generator, it holds

$$\mathcal{E}(\mathcal{R}_\lambda f, g) = -\int \mathcal{R}_\lambda f(h)\mathcal{L}g(h)m(dh).$$

Assume for now that $\psi_\varphi$ belongs to $\mathcal{D}(\mathcal{L})$, and that

$$\mathcal{L}\psi_\varphi(h) = \psi_\varphi(h)\Big(ic\langle h, \varphi''\rangle - \frac{\sigma^2}{2}\|\varphi\|_{L^2}^2\Big).$$

Combining the last three identities, and using the general fact that $\mathcal{D}(\mathcal{L}) \subset \mathcal{D}(\mathcal{E})$, we deduce that $\mathcal{R}_\lambda f$ satisfies (22).

*Step 3: Computation of $\mathcal{L}\psi_\varphi$.* Let us show that $\psi_\varphi$ belongs to $\mathcal{D}(\mathcal{L})$. Instead of identifying explicitly the domain of the generator, we prove that the following convergence holds in $L^2(E, dm)$

$$\frac{P_t\psi_\varphi - \psi_\varphi}{t}(h) \longrightarrow \psi_\varphi(h)\Big(ic\langle h, \varphi''\rangle - \frac{\sigma^2}{2}\|\varphi\|_{L^2}^2\Big), \quad t \downarrow 0.$$

Recall that the Markov process $u(t, \cdot)$ can be built as the solution of (21). From this evolution equation, we deduce that

$$d\langle u(t,\cdot), \varphi\rangle = c\langle u(t,\cdot), \varphi''\rangle dt + \sigma\|\varphi\|_{L^2} dB(t),$$

where $B$ is a Brownian motion. Since $P_t\psi_\varphi(u_0) = \mathbb{E}[e^{i\langle u(t,\cdot),\varphi\rangle} \mid u(t,0) = u_0]$, an explicit computation yields the desired result.

*Step 4: Uniqueness.* Let us show that this is the unique $L^2(E, dm)$ map that satisfies this identity. Given the first step above, it suffices to prove that if $F \in L^2(E, dm)$ satisfies

$$\int F(h)\psi_\varphi(h)\Big(\lambda + \frac{\sigma^2}{2}\|\varphi\|_{L^2}^2 - ic\langle h, \varphi''\rangle\Big)m(dh) = 0,$$

for all $C^2$ function $\varphi$ that vanishes at the boundaries, then $F = 0$.

Let $F$ be such a function. Let $e_\mu$ be a normalized eigenfunction of the Dirichlet Laplacian on $[0, 1]$ associated with some eigenvalue $-\mu \leq 0$. For $t \in \mathbb{R}$, set $\varphi := te_\mu$. Then

$$\psi_\varphi(h)\Big(\lambda + \frac{\sigma^2}{2}\|\varphi\|_{L^2}^2 - ic\langle h, \varphi''\rangle\Big) = e^{it\langle h, e_\mu\rangle}\Big(\lambda + t^2\frac{\sigma^2}{2} + i\mu ct\langle h, e_\mu\rangle\Big).$$

Writing $x := \langle h, e_\mu\rangle$ and $g(x) := e^{itx}$, we observe that the last expression coincides with $(\lambda - \mathcal{L}^{OU})g(x)$, where $\mathcal{L}^{OU}$ is the generator of the 1-d Ornstein–Uhlenbeck process

$$dX(t) = -\mu cX(t)dt + \sigma dB(t).$$

Its invariant probability measure $m^{OU}$ is the pushforward of $m$ through $h \mapsto \langle h, e_\mu\rangle$. Let $\mathcal{F}_\mu$ be the sigma-field on $E$ generated by this last map and let $F_\mu : \mathbb{R} \to \mathbb{R}$ be a measurable map s.t. $m[F \mid \mathcal{F}_\mu](h) = F_\mu(\langle h, e_\mu\rangle)$. We have shown that

$$\begin{aligned}\int_{\mathbb{R}} F_\mu(x)(\lambda - \mathcal{L}^{OU})g(x)dm^{OU}(x) &= \int_E F(h)\psi_\varphi(h)\Big(\lambda + \frac{\sigma^2}{2}\|\varphi\|_{L^2}^2 - ic\langle h, \varphi''\rangle\Big)m(dh)\\ &= 0.\end{aligned}$$

As a consequence

$$\int F_\mu(x)(\lambda - \mathcal{L}^{OU})g(x)m^{OU}(dx) = 0,$$

for all $g$ in the vector space spanned by $\{x \mapsto e^{itx} : t \in \mathbb{R}\}$. By approximation arguments, for any $f \in L^2(\mathbb{R}, dm^{OU})$ there exists a sequence $g_n$ in the latter vector space s.t. $\|(\lambda - \mathcal{L}^{OU})g_n - f\|_{L^2(\mathbb{R}, dm^{OU})}$ goes to 0 as $n \to \infty$ and this suffices to deduce that

$$\int F_\mu(x)f(x)m^{OU}(dx) = 0.$$

Consequently $F_\mu = 0$ $m^{OU}$-a.e., and in turn $m[F \mid \mathcal{F}_\mu] = 0$ $m$-a.e.

The very same arguments applies if we consider finitely many $e_{\mu_i}$ simultaneously. More precisely, if we enumerate the eigenvalues of the Dirichlet Laplacian $0 > -\mu_1 > -\mu_2 > \dots$ and if we let $\mathcal{F}_n$ be the sigma-field on $E$ generated by the maps $h \mapsto \langle h, e_{\mu_1}\rangle$, $\dots$, $h \mapsto \langle h, e_{\mu_n}\rangle$, then $m[F \mid \mathcal{F}_n] = 0$ $m$-a.e. for any $n \geq 1$. As a consequence $m[F \mid \mathcal{F}_\infty] = 0$ $m$-a.e., where $\mathcal{F}_\infty$ is the smallest $\sigma$-field generated by all the $\mathcal{F}_n$, $n \geq 1$ augmented with all $m$-null sets. Now we observe that $L^2 \subset L^1 = E$ and that $m(L^2) = 1$. Since the $(e_{\mu_n})_{n\geq 1}$ is a Hilbert basis of $L^2$, we deduce that for $m$-a.e. $h$

$$F(h) = F\left(\sum_{n\geq 1}\langle h, e_{\mu_n}\rangle e_{\mu_n}\right).$$

We thus deduce that $F$ coincides $m$-a.e. with a $\mathcal{F}_\infty$-measurable map. Consequently $F = m[F \mid \mathcal{F}_\infty] = 0$ $m$-a.e. □

3.3. **Equicontinuity.** Assume that for any $v, w \in \Omega_{N,\bar{n}}$ there exists a coupling of the processes $u_N^v$ and $u_N^w$ starting respectively from $v$ and $w$ such that:

$$\mathbb{E}[\|u_N^v(t, \cdot) - u_N^w(t, \cdot)\|_{L^1}] \leq \|v - w\|_{L^1}. \tag{27}$$

Then we deduce that for any $f \in \mathrm{Lip}(E_N)$

$$\begin{aligned}\big|\mathcal{R}_\lambda^N f(v) - \mathcal{R}_\lambda^N f(w)\big| &\leq \int_0^\infty e^{-\lambda t}\mathbb{E}[|f(u_N^v(t, \cdot)) - f(u_N^w(t, \cdot))|]dt\\ &\leq [f]_{\mathrm{Lip}(E_N)}\int_0^\infty e^{-\lambda t}\mathbb{E}[\|u_N^v(t, \cdot) - u_N^w(t, \cdot)\|_{L^1}]dt\\ &\leq \lambda^{-1}[f]_{\mathrm{Lip}(E_N)}\|v - w\|_{L^1},\end{aligned}$$

and Assumption 2.4 is proved.

**Remark 3.3.** Note that, if the r.h.s. of (27) were replaced by $\|v - w\|_{L^1}$ times a function that grows less than exponentially in $t$, Assumption 2.4 would still follow.

We are thus left with constructing the above coupling. Actually we will construct a coupling of all processes $u_N^v$, $v \in \Omega_{N,\bar n}$ such that

(1) it preserves the ordering of the height functions: if $v \geq w$ then $u_N^v(t,\cdot) \geq u_N^w(t,\cdot)$ at all times $t \geq 0$,
(2) for any $v, w$, at all times $t \geq 0$ $\mathbb{E}[\|u_N^v(t,\cdot) - u_N^w(t,\cdot)\|_{L^1}] \leq \|v - w\|_{L^1}$.

To do so, consider a collection of independent Poisson clocks $P_{i,i+1}^k$, $i \in \{1,\dots,n-1\}$, $k \geq 1$ and $P_{i,i-1}^k$, $i \in \{2,\dots,n\}$, $k \geq 1$ of rates $\frac{1}{2}(\tau(k) - \tau(k-1))$. With these Poisson clocks at hand, we can construct the processes $n_N^v$ in the following way. If $P_{i,i\pm1}^k$ rings at time $t$ then for all $v$ such that $n_N^v(t_-, i) \geq k$ we apply the following transition: $n_N^v(t,i) = n_N^v(t_-,i) - 1$ and $n_N^v(t, i \pm 1) = n_N^v(t_-, i \pm 1) + 1$.

It is then elementary to check that this coupling preserves the ordering of the height functions. Regarding the second property, we first consider the particular case where $v \geq w$. Then at all times $t \geq 0$

$$\big\|u_N^w(t,\cdot) - u_N^v(t,\cdot)\big\|_{L^1} = \sum_{i=1}^N \frac{1}{N}\big(u_N^v(t,i) - u_N^w(t,i)\big).$$

Consequently

$$\partial_t \mathbb{E}\big[\big\|u_N^v(t,\cdot) - u_N^w(t,\cdot)\big\|_{L^1}\big] = \sum_{i=1}^{N-1} \frac{1}{N}\mathbb{E}[\mathcal{L}_N(u_N^v(t,i) - u_N^w(t,i))].$$

Since

$$\mathbb{E}\big[\mathcal{L}_N u_N^v(t,i)\big] = \frac{N^{3/2}}{2}\mathbb{E}\big[\tau\big(n_N^v(t,i+1)\big) - \tau\big(n_N^v(t,i)\big)\big],$$

and similarly with $w$, we obtain

$$\begin{aligned}(28)\quad &\partial_t \mathbb{E}[\|u_N^v(t,\cdot) - u_N^w(t,\cdot)\|_{L^1}] \\ &\qquad = \frac{\sqrt{N}}{2}\mathbb{E}[\tau(n_N^v(t,N)) - \tau(n_N^w(t,N))] - \frac{\sqrt{N}}{2}\mathbb{E}[\tau(n_N^v(t,1)) - \tau(n_N^w(t,1))].\end{aligned}$$

Since $u_N^v(t,\cdot) \geq u_N^w(t,\cdot)$ and since the total number of particles is the same in both configurations, we deduce that

$$n_N^v(t,N) \leq n_N^w(t,N), \quad n_N^v(t,1) \geq n_N^w(t,1).$$

Since $k \mapsto \tau(k)$ is non-decreasing, we deduce that $\partial_t \mathbb{E}[\|u_N^v(t,\cdot) - u_N^w(t,\cdot)\|_{L^1}] \leq 0$ so that $\mathbb{E}[\|u_N^v(t,\cdot) - u_N^w(t,\cdot)\|_{L^1}] \leq \|v - w\|_{L^1}$.

Let us now consider the general case where $v, w$ are not necessarily ordered. We introduce the configurations

$$\max(i) := v(i) \vee w(i), \quad \min(i) := v(i) \wedge w(i).$$

Our coupling ensures that

$$u_N^{\min}(t,\cdot) \leq u_N^v(t,\cdot), u_N^w(t,\cdot) \leq u_N^{\max}(t,\cdot).$$

In addition

$$\|v - w\|_{L^1} = \|\max - \min\|_{L^1}.$$

From the first part of the argument, we thus deduce

$$\begin{aligned}\mathbb{E}[\|u_N^v(t,\cdot) - u_N^w(t,\cdot)\|_{L^1}] &\leq \mathbb{E}\big[\big\|u_N^{\max}(t,\cdot) - u_N^{\min}(t,\cdot)\big\|_{L^1}\big] \\ &\leq \|\max - \min\|_{L^1} = \|v - w\|_{L^1}.\end{aligned}$$

3.4. **Discrete characterization and convergence.** The goal of this subsection is to check that Assumption 2.5 is fulfilled. Recall the parameters introduced above Lemma 3.1. We set

$$c := \frac{\rho}{2\alpha}, \quad \sigma := \sqrt{\overline{\tau}},$$

and we let $m = m_{c,\sigma}$ be the law on $E$ of a scaled Brownian bridge, as introduced in Section 3.2.

Recall that $\mathcal{C}$ is the set of all maps $\psi_\varphi(h) = \exp(i\langle h, \varphi\rangle)$, $v \in E$, where $\varphi$ is a $C^2$ function from $[0,1]$ into $\mathbb{R}$ compactly supported in $(0,1)$. Recall the definition of the measure $\Sigma^{\psi_\varphi}$ from Section 3.2

$$\Sigma^{\psi_\varphi}(dh) = \psi_\varphi(h)\left(ic\langle h, \varphi''\rangle - \frac{\sigma^2}{2}\|\varphi\|_{L^2}^2\right)m(dh).$$

We set

$$\Sigma_N^{\psi_\varphi}(dh) := \mathcal{L}_N\psi_\varphi(h) \cdot m_N(dh).$$

**Proposition 3.4.** *The measure $\Sigma_N^{\psi_\varphi}$ converges weakly towards $\Sigma^{\psi_\varphi}$ in the set of finite complex measures on $E$.*

*Proof.* We need to show that for any bounded and continuous function $F : E \to \mathbb{R}$

$$\int_E F(h)\Sigma_N^{\psi_\varphi}(dh) \longrightarrow \int_E F(h)\Sigma^{\psi_\varphi}(dh), \quad N \longrightarrow \infty.$$

From now on, $F$ will implicitly be such a function. Recall that $\varphi$ is a $C^2$ function compactly supported in $(0,1)$, consequently there exist $\epsilon > 0$ such that $\varphi(x) = 0$ for all $x \in [0,\epsilon] \cup [1-\epsilon, 1]$. From now on, we always assume that $N \geq 1/\epsilon$ so that $\varphi(1/N) = \varphi((N-1)/N) = 0$.

Using the expression of the generator in (16), we have

$$\mathcal{L}_N\psi_\varphi(h) = \sum_{k=1}^{N-1} N^2\tau(n_k)\psi_\varphi(h)\left(\frac{e^{\frac{i}{\sqrt{N}}\int_{\frac{k-1}{N}}^{\frac{k}{N}}\varphi(x)dx} + e^{-\frac{i}{\sqrt{N}}\int_{\frac{k}{N}}^{\frac{k+1}{N}}\varphi(x)dx}}{2} - 1\right).$$

A Taylor expansion yields

$$\begin{aligned} e^{\frac{i}{\sqrt{N}}\int_{\frac{k-1}{N}}^{\frac{k}{N}}\varphi(x)dx} &= 1 + \frac{i}{\sqrt{N}}\int_{\frac{k-1}{N}}^{\frac{k}{N}}\varphi(x)dx - \frac{1}{2N^3}\varphi^2(k/N) + O(N^{-4}) \\ &= 1 + \frac{i}{N^{3/2}}\varphi((k-1)/N) - \frac{1}{2N^3}\varphi^2(k/N) \\ &\quad + \frac{i}{\sqrt{N}}\int_{\frac{k-1}{N}}^{\frac{k}{N}}\big(\varphi(x) - \varphi((k-1)/N)\big)dx + O(N^{-4}), \end{aligned}$$

and similarly

$$\begin{aligned} e^{-\frac{i}{\sqrt{N}}\int_{\frac{k}{N}}^{\frac{k+1}{N}}\varphi(x)dx} &= 1 - \frac{i}{N^{3/2}}\varphi(k/N) - \frac{1}{2N^3}\varphi^2(k/N) \\ &\quad + \frac{i}{\sqrt{N}}\int_{\frac{k}{N}}^{\frac{k+1}{N}}\big(\varphi(x) - \varphi(k/N)\big)dx + O(N^{-4}), \end{aligned}$$

This leads to

$$\mathcal{L}_N\psi_\varphi(h) = \sum_{k=1}^{N-1} N^2\tau(n_k)\psi_\varphi(h)(I_1(k) + I_2(k) + I_3(k)),$$

where

$$I_1(k) = -\frac{i}{2N^{3/2}}\left(\varphi\left(\frac{k}{N}\right) - \varphi\left(\frac{k-1}{N}\right)\right), \quad I_2(k) = -\frac{1}{2N^3}\varphi^2\left(\frac{k}{N}\right),$$

and $I_3$ is a remainder that satisfies $|I_3(k)| \leq C_\varphi N^{-7/2}$ for all $k$, where $C_\varphi > 0$ is a constant that only depends on $\varphi$. Let us control the term provided by $I_2$, that is

$$\begin{aligned} &-\int_E \frac{1}{2N}\sum_{k=1}^{N-1}\tau(n_k)\psi_\varphi(h)\varphi^2(k/N)F(h)m_N(dh) \\ &\qquad = -\frac{\overline{\tau}}{2}\int_E \frac{1}{N}\sum_{k=1}^{N-1}\psi_\varphi(h)\varphi^2(k/N)F(h)m_N(dh) \\ &\qquad\qquad - \frac{1}{2N}\int_E\sum_{k=1}^{N-1}(\tau(n_k) - \overline{\tau})\psi_\varphi(h)\varphi^2(k/N)F(h)m_N(dh). \end{aligned}$$

The first term on the r.h.s. converges towards

$$-\frac{\overline{\tau}}{2}\int_E F(h)\|\varphi\|^2 m(dh).$$

The second term equals

$$-\frac{1}{2\sqrt{N}}\int_E \psi_\varphi(h)F(h)\left(\sum_{k=1}^{N-1} q_N\left(\frac{k}{N}\right)\left[\varphi^2\left(\frac{k}{N}\right) - \varphi^2\left(\frac{k+1}{N}\right)\right]\right)m_N(dh).$$

The last bound stated in Lemma 3.1 ensures that this quantity converges to 0 as $N \to \infty$. Regarding $I_1$, we first compute

$$
\begin{aligned}
-\sum_{k=1}^{N-1} N^2 \tau(n_k)\psi_\varphi(h)\frac{i}{2N^{3/2}}\big(\varphi(k/N)-\varphi((k-1)/N)\big) & \\
&= -\frac{i}{2N}\sum_{k=1}^{N-1} N^{1/2}(\tau(n_k)-\overline{\tau})\psi_\varphi(h)N\big(\varphi(k/N)-\varphi((k-1)/N)\big) \\
&= \frac{i}{2N}\sum_{k=1}^{N-1} q_N(k/N)N^2\Delta_N\varphi(k/N)\psi_\varphi(h),
\end{aligned}
$$

where $\Delta_N\varphi(k/N) = \varphi((k+1)/N) - 2\varphi(k/N) + \varphi((k-1)/N)$. By Lemma 3.1, this expression, multiplied by $F$ and integrated against $m_N$, converges to

$$\frac{i}{2}\mathbb{E}\big[F(\beta)\langle\zeta,\varphi''\rangle\psi_\varphi(\beta)\big].$$

Since $\zeta - \frac{\rho}{\alpha}\beta$ is a centered Gaussian process independent of $\beta$, we deduce that this last expression coincides with

$$\frac{i}{2}\mathbb{E}\big[F(\beta)\langle\frac{\rho}{\alpha}\beta,\varphi''\rangle\psi_\varphi(\beta)\big] = \frac{i}{2}\int F(h)\frac{\rho}{\alpha}\langle h,\varphi''\rangle\psi_\varphi(h)m(dh).$$

Finally, the term $I_3$ has a negligible contribution. □

3.5. **Tightness.** Let $H^{-\beta}$ be the Sobolev space of all distributions $f$ on $(0,1)$ such that

$$\|f\|^2_{H^{-\beta}} := \sum_{n\geq 1} n^{-2\beta}\langle f, e_n\rangle^2 < \infty,$$

where $(e_n)_{n\geq 1}$ is the usual eigenbasis of the Dirichlet Laplacian on $[0,1]$.

Note that $L^1$ is continuously embedded in $H^{-\beta}$ as soon as $\beta > 1/2$. The next proposition shows that Assumption 2.3 is satisfied with $E^- = H^{-\beta}$ for any $\beta > 1$.

**Proposition 3.5.** *Take $\beta > 1$. The sequence $u_N$, $N \geq 1$ is tight in $\mathbb{D}([0,\infty), H^{-\beta})$ and all limiting points are in $C([0,\infty), H^{-\beta})$.*

*Proof.* To obtain tightness in $\mathbb{D}([0,\infty), H^{-\beta})$ with limiting points in $\mathbb{C}([0,\infty), H^{-\beta})$ it suffices to show that the sequence of initial laws $u_N(0,\cdot)$ is tight and that for all $T,\delta > 0$ we have

$$\lim_{h\downarrow 0}\overline{\lim_{N\to\infty}}\frac{1}{h}\sup_{0\leq s\leq T}\mathbb{P}\bigg(\sup_{s\leq t\leq s+h}\|u_N(t)-u_N(s)\|_{H^{-\beta}} > \delta\bigg) = 0, \tag{29}$$

see for instance [2, Theorem 13.2]. The former is immediate since we start from the stationary measure $m_N$ that weakly converges to $m$. To check the latter condition, we apply the so-called Lyons–Zheng decomposition [23] that exploits the reversibility of the process w.r.t. its invariant measure. Fix some arbitrary $T > 0$. Let $\mathcal{F}_t, t \geq 0$ be the filtration associated with $u_N(t), t \geq 0$, and let $\widetilde{\mathcal{F}}_t, t \in [0,T]$ be the filtration associated with the reversed process $u_N(T-t), t \in [0,T]$. For any $n \geq 1$, the process $\widehat{u}_N(t,n) := \langle u_N(t), e_n\rangle$ satisfies

$$\widehat{u}_N(t,n) - \widehat{u}_N(s,n) = \int_s^t \mathcal{L}_N\widehat{u}(r,n)dr + M_N(t,n) - M_N(s,n), \tag{30}$$

where $M_N(t,n)$ is an $\mathcal{F}_t$-martingale. Using the reversibility of the process, it is simple to get the identity

$$
\begin{aligned}
&\widehat{u}_N(T-(T-s),n) - \widehat{u}_N(T-(T-t),n) \\
&\qquad = \int_{T-t}^{T-s}\mathcal{L}_N\widehat{u}_N(T-r,n)dr + \widetilde{M}_N(T-s,n) - \widetilde{M}_N(T-t,n),
\end{aligned} \tag{31}
$$

where $\widetilde{M}_N(t,n)$ is an $\widetilde{\mathcal{F}}_t$ martingale. Therefore,

$$\widehat{u}_N(t,n) - \widehat{u}_N(s,n) = \frac{1}{2}\Big(M_N(t,n) - M_N(s,n) + \widetilde{M}_N(T-t,n) - \widetilde{M}_N(T-s,n)\Big). \tag{32}$$

Using the Burkholder–Davis–Gundy inequality (see for instance [18]) to the martingales we deduce that for all $p \geq 2$

$$\mathbb{E}^N\left[\sup_{t\in[s,s+h]}\left|M_N(t,n)-M_N(s,n)\right|^p\right]^{\frac{1}{p}} \lesssim \mathbb{E}^N\left[\sup_{t\in[s,s+h]}\left\langle M_N(\cdot,n)-M_N(s,n)\right\rangle_t^{\frac{p}{2}}\right]^{\frac{1}{p}} + \mathbb{E}^N\left[\sup_{r\in(s,s+h]}\left|M_N(r,n)-M_N(r-,n)\right|^p\right]^{\frac{1}{p}},$$

uniformly over all $N \geq 1$, all $s, h \geq 0$ and all $n \geq 1$. The jumps of $M_N$ are of size at most $N^{-3/2}|e'_n| = O(nN^{-3/2})$ so that the second term is easily bounded. Regarding the first term, observe that the predictable bracket is non-decreasing in time so that it suffices to take $t = s + h$. Now observe that there exists $C > 0$ such that for all $N$ and $n$

$$\left\langle M_N(\cdot,n)-M_N(s,n)\right\rangle_{s+h} = \int_s^{s+h}\frac{1}{N}\sum_{k=1}^N\frac{1}{2}N^2\tau(n(r,k)\left(\left(e_n\left(\frac{k}{N}\right)-e_n\left(\frac{k-1}{N}\right)\right)^2 + \left(e_n\left(\frac{k+1}{N}\right)-e_n\left(\frac{k}{N}\right)\right)^2\right)dr \leq Cn^2\int_s^{s+h}\frac{1}{N}\sum_{k=1}^N\tau(n(r,k))dr,$$

where we use the notation $(\langle M(\,\cdot\,)\rangle_s)_{s\geq 0}$ for the predictable bracket of any martingale $M$. By Jensen's inequality

$$\left|\int_s^{s+h}\frac{1}{N}\sum_{k=1}^N\tau(n(r,k))dr\right|^{\frac{p}{2}} \leq h^{\frac{p}{2}-1}\int_s^{s+h}\frac{1}{N}\sum_{k=1}^N\tau(n(r,k))^{p/2}dr.$$

Noting that the law of $\{n(i), i \leq N\}$ is exchangeable under $m_N$ we get

$$\mathbb{E}^N\left[\left\langle M_N(\cdot,n)-M_N(s,n)\right\rangle_{s+h}^{\frac{p}{2}}\right]^{\frac{1}{p}} \lesssim h^{\frac{1}{2}}n\,m_N\left[(\tau(n(1)))^{p/2}\right]^{1/p}.$$

The moment on the r.h.s. is controlled by the last estimate of Lemma 3.1. Consequently,

$$\mathbb{E}^N\left[\sup_{t\in[s,s+h]}\left|M_N(t,n)-M_N(s,n)\right|^p\right]^{\frac{1}{p}} \lesssim n(h^{\frac{1}{2}}+N^{-\frac{3}{2}}).$$

We deduce that

$$\mathbb{E}^N\Big[\sup_{s\leq t\leq s+h}\left|\widehat{u}_N(t,n)-\widehat{u}_N(s,n)\right|^p\Big]^{\frac{1}{p}} \lesssim n(h^{\frac{1}{2}}+N^{-\frac{3}{2}}),$$

uniformly over all $N \geq 1$, $s \in [0, T-h]$ and $n \geq 1$. Applying the triangle inequality w.r.t. the $L^{p/2}(\Omega)$-norm at the second line we get

$$\begin{aligned}\mathbb{E}^N\left[\sup_{s\leq t\leq s+h}\left\|u_N(t)-u_N(s)\right\|_{H^{-\beta}}^p\right]^{\frac{1}{p}} &\leq \mathbb{E}^N\left[\left(\sum_{n\geq 1}n^{-2\beta}\sup_{s\leq t\leq s+h}\left|\widehat{u}_N(t,n)-\widehat{u}_N(s,n)\right|^2\right)^{p/2}\right]^{\frac{1}{p}}\\ &\leq \left(\sum_{n\geq 1}n^{-2\beta}\mathbb{E}^N\Big[\sup_{s\leq t\leq s+h}\left|\widehat{u}_N(t,n)-\widehat{u}_N(s,n)\right|^p\Big]^{\frac{2}{p}}\right)^{\frac{1}{2}}\\ &\lesssim h^{\frac{1}{2}}+N^{-\frac{3}{2}},\end{aligned}$$

if $2\beta - 1 > 1$ so that (29) follows provided $p \geq 3$. □

We have all the ingredients at hand to prove the following convergence result.

**Theorem 3.6.** *The process $(u_N(t), t \geq 0)$, starting from its stationary measure $m_N$, converges in law to the process $(u(t), t \geq 0)$, starting from $m_{c,\sigma}$, in the space*

$$\mathbb{D}([0,\infty), H^{-\beta}([0,1]))$$

*for any given $\beta > 1$.*

*Proof.* We aim at applying Theorem 2.6. Assumption 2.1 is the content of Proposition 3.2, the convergence in law required in Assumption 2.2 is checked in Lemma 3.1, Assumption 2.3 is the content of Proposition 3.5, Assumption 2.4 is discussed in Section 3.3 and Assumption 2.5 is proven in Proposition 3.4. □

## 4. Pair of reflected interfaces

The dynamics studied in this section is a special case of a dynamics of $k \geq 2$ reflected interfaces, introduced previously in [19, 28] in the context of dynamics of planar structures and tilings. See also Remark 4.8 below.

We consider the set $\Omega_N$ of all pairs $\mathtt{u} = (\mathtt{v}, \mathtt{w})$ of lattice paths from $\{0, \dots, 2N\}$ to $\mathbb{Z}$ that make $\pm 1$ steps, are pinned at height 0 at both ends and are ordered, namely

$$\mathtt{v}(k+1) - \mathtt{v}(k) \in \{-1, 1\}, \quad \mathtt{w}(k+1) - \mathtt{w}(k) \in \{-1, 1\},$$
$$\mathtt{v}(0) = \mathtt{w}(0) = \mathtt{v}(2N) = \mathtt{w}(2N) = 0,$$
$$\mathtt{v} \geq \mathtt{w}.$$

We let $\mathtt{m}$ be the uniform measure on $\Omega_N$. Let $\Delta$ be the discrete one-dimensional Laplacian, that is

$$\Delta f(k) = f(k+1) - 2f(k) + f(k-1).$$

Given a configuration $\mathtt{u} = (\mathtt{v}, \mathtt{w}) \in \Omega_N$, we say that $k \in \{1, \dots, 2N-1\}$ is a *contact point* if

$$\mathtt{v}(\ell) = \mathtt{w}(\ell), \quad \forall\, \ell \in \{k-1, k, k+1\}.$$

We also say that $\mathtt{v}$ forms an upward corner at $k$ if $\Delta\mathtt{v} = -2$ and a downward corner at $k$ if $\Delta\mathtt{v} = 2$. The same definition holds for $\mathtt{w}$. On $\Omega_N$, we introduce a partial order by saying that $\mathtt{u} = (\mathtt{v}, \mathtt{w}) \geq \mathtt{u}' = (\mathtt{v}', \mathtt{w}')$ iff $\mathtt{v}(k) \geq \mathtt{v}'(k)$ and $\mathtt{w}(k) \geq \mathtt{w}'(k)$ for every $k$. Given a lattice path $\mathtt{v}$, if $\mathtt{v}$ forms an (upward or downward) corner at $k$, we let $\mathtt{v}^k$ denote the configuration where $\mathtt{v}(k)$ is flipped to $\mathtt{v}(k) + \Delta\mathtt{v}(k)$.

Note that, if $k$ is a contact point and $\mathtt{v}, \mathtt{w}$ form an upward (resp. downward) corner there, then the configuration $(\mathtt{v}^k, \mathtt{w})$ does not belong to $\Omega_N$, but $(\mathtt{v}, \mathtt{w}^k)$ does (resp. $(\mathtt{v}, \mathtt{w}^k)$ does not belong to $\Omega_N$, but $(\mathtt{v}^k, \mathtt{w})$ does).

We consider the following corner flip dynamics on $\Omega_N$:

(1) for every $k \in \{1, \dots, 2N-1\}$ which is not a contact point, if $\mathtt{v}$ (resp. $\mathtt{w}$) forms a corner at $k$, then $\mathtt{v}(k)$ (resp. $\mathtt{w}(k)$) is flipped at rate $(2N)^2/2$, so that the resulting configuration is $(\mathtt{v}^k, \mathtt{w})$ (resp. $(\mathtt{v}, \mathtt{w}^k)$);
(2) for every contact point $k$, if $\mathtt{v}$ and $\mathtt{w}$ form an upward corner (resp. downward corner) then $\mathtt{w}(k)$ (resp. $\mathtt{v}(k)$) flips at rate $(2N)^2/2$, so that the resulting configuration is $(\mathtt{v}, \mathtt{w}^k)$ (resp. $(\mathtt{v}^k, \mathtt{w})$;
(3) for every contact point $k$, if $\mathtt{v}$ and $\mathtt{w}$ form a corner then they flip simultaneously at rate $(2N)^2/4$, so that the resulting configuration is $(\mathtt{v}^k, \mathtt{w}^k)$.

We refer to Figure 2 for an illustration. Let us emphasize that all these jumps are driven by independent Poisson clocks. It is straightforward to check that this dynamics is reversible w.r.t. the uniform measure $\mathtt{m}$ on $\Omega_N$.

**Remark 4.1.** Let us point out that the jump rates at contact points are chosen in such a way that the generator satisfies

$$\mathcal{L}_N[\mathtt{v}(k) + \mathtt{w}(k)] = \frac{(2N)^2}{2}\Delta(\mathtt{v}(k) + \mathtt{w}(k)).$$

This identity would still hold if we imposed:

(1) for every contact point $k$, if $\mathtt{v}$ and $\mathtt{w}$ form an upward corner (resp. downward corner) then $\mathtt{w}(k)$ (resp. $\mathtt{v}(k)$) flips at rate $\alpha$,
(2) for every contact point $k$, if $\mathtt{v}$ and $\mathtt{w}$ form a corner then they flip simultaneously at rate $\beta$,

with $\alpha, \beta \geq 0$ satisfying $\frac{\alpha}{2} + \beta = (2N)^2/2$. In this more general setting, the invariant measure is no longer uniform nor reversible (except if $\alpha = (2N)^2/2$ and $\beta = (2N)^2/4$).

In order to take the continuum limit $N \to \infty$, we need to suitably rescale the paths. Given $\mathtt{u} \in \Omega_N$, we define $h = (h_v, h_w)$ where $h_v : [0,1] \mapsto \mathbb{R}$ is the continuous function defined via

$$h_v(k/(2N)) := \frac{1}{\sqrt{2N}}\mathtt{v}(k), \quad k \in \{0, 1, \dots, 2N\} \tag{33}$$

and by linear interpolation in between. For lightness of notation, we do not give an index $N$ to the rescaled functions, in contrast with what we did for the zero-range process. The function $h_w$ is defined analogously. Then, $h$ is an element of the space $E$ of all pairs $h = (h_v, h_w)$ of continuous functions from $[0,1]$ into $\mathbb{R}$ such that $h_v \geq h_w$. On $E$, we consider the distance $\|h - h'\|_E := \|h_v - h'_v\|_{L^1} + \|h_w - h'_w\|_{L^1}$. Note that $E$ is a separable metric space, but it is not complete, see the next subsection for a discussion on that point. For fixed $N$, $h_u$ belongs to the finite set $E_N \subset E$ (obviously defined as the image of $\Omega_N$ via the rescaling). Also, $u^{u_0}(t)$ denotes the rescaled evolving configuration at time $t$, with rescaled initial configuration $u_0$.





FIGURE 2. Dynamics at contact points.

4.1. **The SPDE.** The limiting dynamics is better described in terms of the sum and difference of the interfaces. Let us introduce the associated stochastic PDEs. Given a space-time white noise $\xi^S$ on $\mathbb{R}_+ \times [0,1]$ we consider the solution $S$ of the following stochastic PDE

$$\begin{cases} \partial_t S(t,x) = \frac{1}{2}\partial_x^2 S(t,x) + \xi^S, \quad x \in [0,1], \quad t \geq 0, \\ S(t,0) = S(t,1) = 0, \end{cases}$$

where the initial condition $S(0,\cdot)$ is any continuous function on $[0,1]$. This equation should be understood in the weak sense: for any $C^2$ function $h : [0,1] \to \mathbb{R}$ that vanishes at 0 and 1, we have

$$\langle S(t,\cdot), h\rangle = \langle S(0,\cdot), h\rangle + \frac{1}{2}\int_0^t \langle S(t,\cdot), h''\rangle + \int_0^t h(x)\xi^S(ds,dx). \tag{34}$$

It is well-known that there exists a unique solution $S$ which is continuous in time with values in $C([0,1])$ (endowed with the sup-norm, i.e. the solution is continuous in both variables on $[0,+\infty)\times[0,1]$). Moreover the solution $(S(t,\cdot))_{t\geq 0}$ is a Markov process which admits a unique invariant (reversible) measure, given by the law $m_{\text{bridge}}$ of the standard Brownian bridge on $[0,1]$.

Given a space-time white noise $\xi^D$, independent of $\xi^S$, we consider the solution $(D,\eta)$ of the following stochastic PDE

$$\begin{cases} \partial_t D(t,x) = \frac{1}{2}\partial_x^2 D(t,x) + \xi^D + \eta, \quad x \in [0,1], \quad t \geq 0, \\ D(t,x) \geq 0, \quad \int_{t\in\mathbb{R}_+}\int_{x\in(0,1)} D(t,x)\eta(dt,dx) = 0, \\ D(t,0) = D(t,1) = 0, \end{cases} \tag{35}$$

where the initial condition $D(0,\cdot)$ is any given non-negative continuous function. Here $\eta$ is a random non-negative Radon measure on $\mathbb{R}_+ \times (0,1)$ and the equation is understood similarly as in (34) except that one considers the additional term $\int_0^t h(x)\eta(ds,dx)$. The random variable $\eta$ is a reflection measure: it is supported by the space time points $(t,x)$ where the interface $D$ hits 0, and at these points, it pushes the interface upwards to keep it non-negative.

Nualart and Pardoux [26] showed that almost surely, there exists a unique pair $(D,\eta)$ satisfying the system above (note that the construction therein is purely deterministic). The process $(D(t,\cdot))_{t\geq 0}$ is Markov with values in $C_+([0,1]) := \{f \in C([0,1]) : f \geq 0\}$ and admits a unique invariant (reversible) measure, given by the law $m_{\text{exc}}$ of the normalised Brownian excursion, see [33, 29].

Since $(S,D)$ is an independent pair of Markov processes, then the pair $(S,D)$ is itself a Markov process on $E := C([0,1]) \times C_+([0,1])$ that admits an invariant and reversible invariant measure $m_{\text{bridge}} \otimes m_{\text{exc}}$.

In this situation, $E$ has a natural metric given by the sup-norm, which makes it a Polish space and $(S,D)$ an a.s. continuous process. However we have allowed ourselves in the black box result of Section 2 to endow our state space $E$ with a distance $d$ which makes it separable but not necessarily complete. Here we exploit this freedom and we choose $d$ as the $L^1([0,1],dx)$ distance: for $u = (u^1,u^2), v = (v^1,v^2) \in E = C([0,1]) \times C_+([0,1])$

$$\|u - v\|_E := \sum_{i=1}^{2} \|u^i - v^i\|_{L^1([0,1],dx)}.$$

The reason for this choice is that the contraction property (39) for the discrete processes can be proved in this weaker distance.

We then define $u = (v,w)$ through

$$v := \frac{S+D}{\sqrt{2}}, \quad w = \frac{S-D}{\sqrt{2}}.$$

It solves

$$\begin{cases} \partial_t v = \frac{1}{2}\partial_x^2 v + \xi^v + \frac{1}{\sqrt{2}}\eta, & x \in [0,1], \quad t \geq 0, \\ \partial_t w = \frac{1}{2}\partial_x^2 w + \xi^w - \frac{1}{\sqrt{2}}\eta, & x \in [0,1], \quad t \geq 0, \\ v \geq w, \quad \eta(dt,dx) \geq 0, & \int_{(0,1)\times\mathbb{R}_+} (v-w)(t,x)\eta(dt,dx) = 0, \end{cases} \tag{36}$$

where $\xi^v := \frac{\xi^S + \xi^D}{\sqrt{2}}$ and $\xi^w := \frac{\xi^S - \xi^D}{\sqrt{2}}$ are still independent space-time white noises.

We let $m$ be the pushforward of $m_{\text{bridge}} \otimes m_{\text{exc}}$ through the map $(S,D) \mapsto (\frac{S+D}{\sqrt{2}}, \frac{S-D}{\sqrt{2}})$. The process $u = (v,w)$ is a Markov process taking values in $E$ with invariant (reversible) measure $m$. Our next proposition provides a characterization of its resolvents and thus establishes Assumption 2.1 in the present model. This proposition takes advantage of the independence of $S$ and $D$, and therefore mixes the two sets of coordinates ($(S,D)$ and $(v,w)$).

For any pair of continuous functions $(\varphi_v, \varphi_w)$ from $[0,1]$ into $\mathbb{R}$, we define for any $h \in E$

$$\psi_\varphi(h) := e^{i\langle h_v, \varphi_v\rangle + i\langle h_w, \varphi_w\rangle}.$$

We call $(\mathcal{R}_\lambda)_{\lambda>0}$ the resolvent family of the Markov process $(S,D)$.

**Proposition 4.2.** *Let $f$ be a bounded Lipschitz function from $E$ to $\mathbb{R}$. For any $\lambda > 0$, $\mathcal{R}_\lambda f$ is the unique bounded Lipschitz-continuous function $F : E \to \mathbb{R}$ such that for any pair of $C^2$ functions $(\varphi_v, \varphi_w)$ from $[0,1]$ into $\mathbb{R}$ with compact supports in $(0,1)$, the following holds:*

$$\int F(h)(\lambda\psi_\varphi(h)m(dh) - \Sigma^{\psi_\varphi}(dh)) = \int f(h)\psi_\varphi(h)m(dh), \tag{37}$$

*where, writing $h_S = \frac{h_v+h_w}{\sqrt{2}}$ and $h_D = \frac{h_v-h_w}{\sqrt{2}}$*

$$\begin{aligned} \Sigma^{\psi_\varphi}(dh) = \frac{1}{2}\psi_\varphi(h)\Big(i\langle h_v, \varphi_v''\rangle + i\langle h_w, \varphi_w''\rangle - \|\varphi_v\|_{L^2}^2 - \|\varphi_w\|_{L^2}^2\Big)m(dh) \\ + \frac{i}{2}\psi_\varphi(h)\int_0^1 dr \frac{1}{\sqrt{2\pi r^3(1-r)^3}} \frac{(\varphi_v - \varphi_w)}{\sqrt{2}}(r) \\ m_{\text{exc}}(dh_D \mid h_D(r) = 0) m_{\text{bridge}}(dh_S). \end{aligned} \tag{38}$$

In this case, we let $\mathcal{C}$ denote the set of all functions $\psi_\varphi(\,\cdot\,)$ with $\varphi$ as in the statement of the proposition. Let us point out that the probability measure $m_{\text{exc}}(dh_D \mid h_D(r) = 0)$ can be defined by passing to the limit on a regular conditioning of the form $\{h_D(r) < \varepsilon\}$. This probability measure then coincides with the law of the concatenation of two Brownian excursions rescaled on $(0,r)$ and $(r,1)$ respectively: we refer to Step 3 in the proof of Proposition 4.4 for more details in this direction.

*Proof.* By [1, chapter V], the Markov process $(S,D)$ is associated with the tensor product $\mathcal{E}$ of the Dirichlet forms $\mathcal{E}^S, \mathcal{E}^D$ of $S$ and $D$ respectively. Moreover $\mathcal{R}_\lambda f$ is the unique function in $\mathcal{D}(\mathcal{E})$ such that (26) holds for all $g \in \mathcal{D}(\mathcal{E})$. Then if we choose $g = \psi_\varphi$, we obtain by integration by parts that $\mathcal{R}_\lambda f$ does satisfy (37), using the explicit form of the integration by parts for $\mathcal{E}^D$ given in [30]. On the other hand, if $F : E \to \mathbb{R}$ is bounded Lipschitz-continuous then $F \in \mathcal{D}(\mathcal{E})$ and if $F$ also satisfies (37) then by integration by parts $F$ satisfies (26) for $g = \psi_\varphi$. Since exponential functions are a core in the domains $\mathcal{D}(\mathcal{E}^S)$ and $\mathcal{D}(\mathcal{E}^D)$, then $F$ satisfies (26) for all $g \in \mathcal{D}(\mathcal{E})$. □

Let $m_N$ be the pushforward of $\mathtt{m}$ through the map $(\mathtt{v}, \mathtt{w}) \mapsto (h_v, h_w)$, see (33).

**Lemma 4.3.** *As $N \to \infty$, $m_N$ converges weakly to the measure $m$.*

*Proof.* It is known, see for instance [13, Theorem 2.7], that $m_N$ converges weakly to a probability measure on $C([0,1], \mathbb{R}^2)$ (and a fortiori on $E$) under which $(h_v, h_w)$ solves

$$\begin{aligned} dh_v(t) &= -\frac{h_v(t)}{1-t}dt + \frac{dt}{h_v(t) - h_w(t)} + dB^v(t), \qquad h_v(0) = 0, \\ dh_w(t) &= -\frac{h_w(t)}{1-t}dt + \frac{dt}{h_w(t) - h_v(t)} + dB^w(t), \quad h_w(0) = 0, \end{aligned}$$

where $B^v$ and $B^w$ are independent standard Brownian motions. It is then straightforward to check that $h_S := (h_v + h_w)/\sqrt{2}$ and $h_D := (h_v - h_w)/\sqrt{2}$ solve

$$\begin{aligned} dh_S(t) &= -\frac{h_S(t)}{1-t}dt + dB^S(t), \qquad\qquad h_S(0) = 0, \\ dh_D(t) &= -\frac{h_D(t)}{1-t}dt + \frac{dt}{h_D(t)} + dB^D(t), \quad h_D(0) = 0, \end{aligned}$$

where $B^S$ and $B^D$ are independent standard Brownian motions. As a consequence $h_S$ is a Brownian bridge and $h_D$ an independent normalised Brownian excursion. □

This lemma ensures that Assumption 2.2 is satisfied.

4.2. **Equicontinuity.** We follow the same strategy as in Section 3.3 in order to establish Assumption 2.4. It suffices to show that for any $u_0, u_0' \in E_N$, there exists a coupling under which

$$\mathbb{E}\Big[\Big\|u^{u_0}(t) - u^{u_0'}(t)\Big\|_E\Big] \leq \|u_0 - u_0'\|_E. \tag{39}$$

From now on, we write $u$ and $u'$ for $u^{u_0}$ and $u^{u_0'}$.

There exists a global monotone coupling of the processes starting from all initial configurations $\mathtt{u}_0 \in \Omega_N$ such that if $\mathtt{u}_0 \geq \mathtt{u}_0'$ then almost surely $\mathtt{u}^{u_0}(t) \geq \mathtt{u}^{u_0'}(t)$ for all $t \geq 0$. Indeed, the dynamics we are considering is a special case of a dynamics of $k$ reflected interfaces (see Remark 4.8 below), for which the existence of a global monotone coupling is known [28].

With this coupling at hand, let us consider first the case $\mathtt{u}_0 \geq \mathtt{u}_0'$. Then at all times $t \geq 0$

$$(2N)^{3/2}\|u(t) - u'(t)\|_E = \sum_{k=1}^{2N-1} \mathtt{v}(t,k) - \mathtt{v}'(t,k) + \mathtt{w}(t,k) - \mathtt{w}'(t,k).$$

Consequently,

$$(2N)^{3/2}\partial_t\mathbb{E}[\|u(t) - u'(t)\|_E] = \sum_{k=1}^{2N-1} \mathbb{E}\big[\mathcal{L}_N(\mathtt{v}(t,k) - \mathtt{v}'(t,k) + \mathtt{w}(t,k) - \mathtt{w}'(t,k))\big].$$

Thanks to Remark 4.1,

$$\mathbb{E}\big[\mathcal{L}_N(\mathtt{v}(t,k) + \mathtt{w}(t,k))\big] = \frac{(2N)^2}{2}\mathbb{E}\big[\Delta\mathtt{v}(t,k) + \Delta\mathtt{w}(t,k)\big],$$

and similarly for $\mathtt{v}' + \mathtt{w}'$. Since the interfaces are pinned at height 0 at $k = 0$ and $k = 2N$, an integration by parts gives

$$\begin{aligned}(2N)^{3/2}\partial_t\mathbb{E}\big[\|u_N(t) - u_N'(t)\|_E\big] \\ = -\frac{(2N)^2}{2}\sum_{k\in\{1,2N-1\}} \mathbb{E}\big[\mathtt{v}(t,k) - \mathtt{v}'(t,k) + \mathtt{w}(t,k) - \mathtt{w}'(t,k)\big],\end{aligned}$$

which is non-positive, and therefore (39) is proved.

To treat the general case where $\mathtt{u}_0$ and $\mathtt{u}_0'$ are not ordered, we introduce

$$\mathtt{v}^{\max}(k) = \max(\mathtt{v}_0(k), \mathtt{v}_0'(k)), \quad \mathtt{v}^{\min}(k) = \min(\mathtt{v}_0(k), \mathtt{v}_0'(k)),$$

together with

$$\mathtt{w}^{\max}(k) = \max(\mathtt{w}_0(k), \mathtt{w}_0'(k)), \quad \mathtt{w}^{\min}(k) = \min(\mathtt{w}_0(k), \mathtt{w}_0'(k)).$$

Our coupling implies that almost surely at all times $t \geq 0$

$$\mathtt{v}^{\max}(k,t) \geq \mathtt{v}(k,t), \quad \mathtt{v}'(k,t) \geq \mathtt{v}^{\min}(k,t),$$

and

$$\mathtt{w}^{\max}(k,t) \geq \mathtt{w}(k,t), \quad \mathtt{w}'(k,t) \geq \mathtt{w}^{\min}(k,t),$$

with the notation that $(\mathtt{v}^{\max}(t), \mathtt{w}^{\max}(t))$ is the configuration at time $t$ started from $(\mathtt{v}^{\max}, \mathtt{w}^{\max})$, and analogously for the evolution starting from the minimal configuration. Consequently,

$$\mathbb{E}[\|u(t) - u'(t)\|_E] \leq \mathbb{E}[\|u_N^{\max}(t) - u_N^{\min}(t)\|_E] \leq \|u^{\max} - u^{\min}\|_E.$$

To conclude, one observes that

$$\begin{aligned}\big\|u^{\max} - u^{\min}\big\|_E &= \big\|v^{\max} - v^{\min}\big\|_{L^1} + \big\|w^{\max} - w^{\min}\big\|_{L^1} \\ &= \|v_0 - v_0'\|_{L^1} + \|w_0 - w_0'\|_{L^1} \\ &= \|u_0 - u_0'\|_E.\end{aligned}$$

4.3. **Discrete characterization and convergence.** In this subsection, we check that Assumption 2.5 is satisfied. Given two functions $\varphi_v, \varphi_w$ from $[0,1]$ into $\mathbb{R}$, recall that

$$\psi_\varphi(h) = e^{i\langle h_v, \varphi_v\rangle + i\langle h_w, \varphi_w\rangle}.$$

We define the finite complex measure $\Sigma_N^{\psi_\varphi}$ on $E$ by setting

$$\Sigma_N^{\psi_\varphi}(dh) = \mathcal{L}_N \psi_\varphi(h) m_N(dh). \tag{40}$$

**Proposition 4.4.** *Let $\varphi_v, \varphi_w$ be two $C^2$ functions from $[0,1]$ into $\mathbb{R}$ with compact supports in $(0,1)$. As $N \to \infty$, $\Sigma_N^{\psi_\varphi}$ converges weakly to the finite complex measure $\Sigma^{\psi_\varphi}$ which was defined in (38).*

*Proof of Proposition 4.4.* We write $v^k, w^k$, $k = 1, \dots, 2N-1$, for the configurations $\mathtt{v}^k, \mathtt{w}^k$ after rescaling. Since $\mathtt{v}, \mathtt{w}$ are obtained by affine interpolation in between their values at lattice sites, one can check that

$$\langle h_v, \varphi_v\rangle = \frac{1}{(2N)^{3/2}} \sum_{k=1}^{2N-1} \mathtt{v}(k) \Phi_v(k/(2N)),$$

with

$$\begin{aligned}&\Phi_v(k/(2N))\\ &= 2N \int_{(k-1)/(2N)}^{k/(2N)} \varphi_v(x)(2Nx - (k-1))dx + 2N \int_{k/(2N)}^{(k+1)/(2N)} \varphi_v(x)(k+1-2Nx)dx,\end{aligned}$$

and similarly for $\langle h_w, \varphi_w\rangle$ with a function $\Phi_w$ defined analogously.

A direct computation shows that the generator $\mathcal{L}_N$ can be written as a sum of $\mathcal{L}_N^A$ and $\mathcal{L}_N^R$ where the former deals with non-reflected terms in the generator, that is

$$\mathcal{L}_N^A \psi_\varphi(h) = \sum_{k=1}^{2N-1} (2N)^2 \psi_\varphi(h) A_k(h),$$

with

$$A_k(h) = \frac{1}{2}\left(\frac{\psi_\varphi(h_v^k, h_w)}{\psi_\varphi(h_v, h_w)} - 1\right)\mathbf{1}_{\Delta \mathtt{v}(k) \neq 0} + \frac{1}{2}\left(\frac{\psi_\varphi(h_v, h_w^k)}{\psi_\varphi(h_v, h_w)} - 1\right)\mathbf{1}_{\Delta \mathtt{w}(k) \neq 0},$$

while $\mathcal{L}_N^R$ is the part of the generator coming from the reflection, that is

$$\mathcal{L}_N^R \psi_\varphi(h) = \sum_{k=1}^{2N-1} (2N)^2 \psi_\varphi(h) R_k(h),$$

with

$$\begin{aligned}R_k(h) = &-\frac{1}{2}\left(\frac{\psi_\varphi(h_v^k, h_w)}{\psi_\varphi(h_v, h_w)} - 1\right)\mathbf{1}_{\wedge(k)} - \frac{1}{2}\left(\frac{\psi_\varphi(h_v, h_w^k)}{\psi_\varphi(h_v, h_w)} - 1\right)\mathbf{1}_{\vee(k)}\\ &+ \frac{1}{4}\left(\frac{\psi_\varphi(h_v^k, h_w^k)}{\psi_\varphi(h_v, h_w)} - 1\right)\big(\mathbf{1}_{\wedge(k)} + \mathbf{1}_{\vee(k)}\big).\end{aligned}$$

Here $\wedge(k)$ (resp. $\vee(k)$) is the event that $k$ is a contact point and the interfaces $\mathtt{v}, \mathtt{w}$ have an upward (resp. downward) corner there. We control these terms separately. From now on $O(N^{-a})$ will denote a quantity that is bounded by $CN^{-a}$ where $C > 0$ is independent of $N$ and $k$.

*Step 1: Non-reflected terms.* One can check that

$$\begin{aligned}A_k(h) = &\frac{i}{2(2N)^{3/2}}\big[\Phi_v(k/(2N))\Delta \mathtt{v}(k) + \Phi_w(k/(2N))\Delta \mathtt{w}(k)\big]\\ &- \frac{1}{(2N)^3}\left[\left(\Phi_v\left(\frac{k}{2N}\right)\right)^2 \mathbf{1}_{\Delta \mathtt{v}(k)\neq 0} + \left(\Phi_w\left(\frac{k}{2N}\right)\right)^2 \mathbf{1}_{\Delta \mathtt{w}(k)\neq 0}\right] + O(N^{-9/2}).\end{aligned}$$

Via a summation by parts and recalling that $\varphi_v, \varphi_w$ are $C^2$ functions that have compact support in $(0,1)$, we thus deduce that, up to negligible terms, $\mathcal{L}_N^A \psi_\varphi(u)$ coincides with

$$\frac{1}{2}\psi_\varphi(h)\Big(i\langle h_v, \varphi_v''\rangle + i\langle h_w, \varphi_w''\rangle - \|\varphi_v\|^2 - \|\varphi_w\|^2\Big) + \mathcal{S}_N(h), \tag{41}$$

where

$$\mathcal{S}_N(h) := \psi_\varphi(h)\frac{1}{N}\sum_{k=1}^{2N-1}\left(\Phi_v\left(\frac{k}{2N}\right)^2\left(\frac{1}{2}-\mathbf{1}_{\Delta\mathtt{v}(k)\neq 0}\right)+\Phi_w\left(\frac{k}{2N}\right)^2\left(\frac{1}{2}-\mathbf{1}_{\Delta\mathtt{w}(k)\neq 0}\right)\right). \tag{42}$$

Let us show that $\mathcal{S}_N(h)m_N(dh)$ converges weakly to 0 as $N\to\infty$. Set

$$a_k := \Phi_v^2(k/(2N)).$$

By symmetry, and since $\psi_\varphi$ is a bounded function, it is enough to prove that

$$\frac{1}{N}m_N\left[\left|\sum_{k=1}^{2N-1}a_k\left(\frac{1}{2}-\mathbf{1}_{\Delta\mathtt{v}(k)\neq 0}\right)\right|\right] \tag{43}$$

tends to zero as $N\to\infty$. Let $\mathbb{P}^{\otimes 2}$ denote the law of two iid simple random walks $\mathtt{v},\mathtt{w}$ started at zero, and set

$$D_N := \big\{(\mathtt{v},\mathtt{w}) : \forall\, k, \mathtt{v}(k)\geq\mathtt{w}(k) \quad\text{and}\quad \mathtt{v}(2N)=\mathtt{w}(2N)\big\}.$$

We then have $m_N(\,\cdot\,)=\mathbb{P}^{\otimes 2}(\,\cdot\,|D_N)$.

Using Jensen's inequality for $p>1$ and the definition of $m_N$, (43) is smaller than

$$\begin{aligned}&\frac{1}{\sqrt{N}}\left(m_N\left[\left|\frac{1}{\sqrt{N}}\sum_{k=1}^{2N-1}a_k\left(\frac{1}{2}-\mathbf{1}_{\Delta\mathtt{v}(k)\neq 0}\right)\right|^p\right]\right)^{1/p}\\ &\qquad\leq\frac{1}{\sqrt{N}(\mathbb{P}^{\otimes 2}(D_N))^{1/p}}\left(\mathbb{E}^{\otimes 2}\left[\left|\frac{1}{\sqrt{N}}\sum_{k=1}^{2N-1}a_k\left(\frac{1}{2}-\mathbf{1}_{\Delta\mathtt{v}(k)\neq 0}\right)\right|^p\right]\right)^{1/p}.\end{aligned} \tag{44}$$

Since $\mathbb{P}^{\otimes 2}(D_N)$ is of order $1/N^2$, the constants $a_k$ are bounded uniformly in $N$ and the random variables $\frac{1}{2}-\mathbf{1}_{\Delta\mathtt{v}(k)\neq 0}$ are i.i.d., centered and of finite variance under $\mathbb{P}^{\otimes 2}$, this expression is upper bounded by a $p$-dependent constant times $N^{-1/2+2/p}$, that tends to zero as soon as $p>4$.

*Step 2: Simplifying the reflection terms.* It is straightforward to check that

$$R_k(h)=\frac{i}{2(2N)^{3/2}}\big(\mathbf{1}_{\vee(k)}+\mathbf{1}_{\wedge(k)}\big)\left(\Phi_v\left(\frac{k}{2N}\right)-\Phi_w\left(\frac{k}{2N}\right)+O(N^{-3/2})\right). \tag{45}$$

Since $\varphi_v,\varphi_w$ have compact supports in $(0,1)$, we can restrict ourselves to those $k\in[\delta 2N,(1-\delta)2N]$ for some $\delta>0$ small enough but fixed w.r.t. $N$. By Lemma 4.6(1) below, the quantity $O(N^{-3/2})$ has a negligible contribution in $\mathcal{L}_N^R\psi_\varphi(u)m_N(du)$. Furthermore, by Lemma 4.6(2) it suffices to show that, as $N\to\infty$ and

then $a \uparrow \infty$ the measure

$$(46)\quad (2N)^2\psi_\varphi(h) \sum_{k\in[\delta 2N,(1-\delta)2N]} \sum_{j\in[-a\sqrt{2N},a\sqrt{2N}]} \frac{i}{2(2N)^{3/2}}\big(\mathbf{1}_{\vee(k,j)}+\mathbf{1}_{\wedge(k,j)}\big) \times\left(\Phi_v\left(\frac{k}{2N}\right)-\Phi_w\left(\frac{k}{2N}\right)\right)m_N(dh),$$

converges weakly to

$$(47)\quad \frac{i}{2}\psi_\varphi(h)\int_0^1 dr \frac{1}{\sqrt{2\pi r^3(1-r)^3}}\frac{(\varphi_v-\varphi_w)}{\sqrt{2}}(r) \, dm_{\mathrm{exc}}(h_D \mid h_D(r)=0)dm_{\mathrm{bridge}}(h_S),$$

where $h_S=(h_v+h_w)/\sqrt{2}$ and $h_D=(h_v-h_w)/\sqrt{2}$. Here, $\vee(k,j)$ (resp. $\wedge(k,j)$) denotes the event that $k$ is a contact point, the interfaces $\mathtt{v},\mathtt{w}$ have a downward (resp. upward) corner there and $\mathtt{v}(k\pm1)=\mathtt{w}(k\pm1)=j$.

*Step 3: Concatenation maps and decomposition of the invariant measure - the continuous case.* For any functions $h_1,h_2$ from $[0,1]$ into $\mathbb{R}^2$ and for any given $r\in(0,1)$, we define the concatenation map

$$T_r(h_1,h_2)(x)=\sqrt{r}\,h_1\Big(\frac{x}{r}\Big)\mathbf{1}_{\{x\le r\}}+\sqrt{1-r}\,h_2\left(\frac{x-r}{1-r}\right)\mathbf{1}_{\{x>r\}},\quad x\in[0,1].$$

Note that this is a function from $[0,1]$ into $\mathbb{R}^2$.

We observe that under $m$, $h_S$ is a Brownian bridge on $[0,1]$ so that for any $r\in(0,1)$, $h_S(r)\sim\mathcal{N}(0,r(1-r))$, and $h_D$ is an independent normalised Brownian excursion. Let $m_{(0,0)\to(1,y)}$ denote the probability measure on $E$ under which $h_S$ is a Brownian bridge between $(0,0)$ and $(1,y)$ and $h_D$ is an independent Brownian excursion. In particular $m_{(0,0)\to(1,0)}=m$. Let also $m_{(0,y)\to(1,0)}$ denote the probability measure on $E$ under which $h_S$ is a Brownian bridge between $(0,y)$ and $(1,0)$ and $h_D$ is an independent Brownian excursion.

The law of $T_r(h_1,h_2)$ under the probability measure (on $E\times E$)

$$\int_{y\in\mathbb{R}}\frac{1}{\sqrt{2\pi r(1-r)}}e^{-\frac{y^2}{2r(1-r)}}m_{(0,0)\to(1,y)}(dh_1)m_{(0,y)\to(1,0)}(dh_2)\,dy,$$

coincides with the law of $h$ under $m_{\mathrm{bridge}}(dh_S)m_{\mathrm{exc}}(dh_D\mid h_D(r)=0)$.

*Step 4: Concatenation maps and decomposition of the invariant measure – the discrete case.* For any functions $h_1,h_2$ from $[0,1]$ into $\mathbb{R}^2$ and for any given integer $k\in\{2,\dots,N-2\}$ we let $T^{\wedge}_{N,k}(h_1,h_2)$ be the pair $(h_v,h_w)$ defined by

$$(h_v,h_w)(x):=\begin{cases}\sqrt{\frac{k-1}{2N}}\,h_1\Big(\frac{x2N}{k-1}\Big) & \text{if } x\le\frac{k-1}{2N},\\ \sqrt{\frac{k-1}{2N}}\,h_1(1)+\Big(\frac{1-|x2N-k|}{\sqrt{2N}},\frac{1-|x2N-k|}{\sqrt{2N}}\Big) & \text{if } \frac{k-1}{2N}<x\le\frac{k+1}{2N},\\ \sqrt{1-\frac{k+1}{2N}}\,h_2\Big(\frac{x-\frac{k+1}{2N}}{1-\frac{k+1}{2N}}\Big) & \text{if } \frac{k+1}{2N}<x\le 1.\end{cases}$$

$T^{\vee}_{N,k}(h_1,h_2)$ is defined similarly except that $\frac{1-|x2N-k|}{\sqrt{2N}}$ is replaced by $\frac{|x2N-k|-1}{\sqrt{2N}}$.

Let $\nu_{(0,a)\to(n,b)}$ denote the uniform measure on the set of pairs $(\mathtt{v},\mathtt{w})$ of lattice paths of length $n$ such that $\mathtt{v}\ge\mathtt{w}$, $\mathtt{v}(0)=\mathtt{w}(0)=a$ and $\mathtt{v}(n)=\mathtt{w}(n)=b$. We consider the rescalings $h_v=\mathtt{v}(\cdot\,n)/\sqrt{n}$ and $h_w=\mathtt{w}(\cdot\,n)/\sqrt{n}$. The law of $T^{\wedge}_{N,k}(h_1,h_2)$ under the probability measure $\nu_{(0,0)\to(k-1,j)}(du_1)\nu_{(0,j)\to(2N-k-1,0)}(du_2)$ coincides with the law of $h$ under $m_N(dh\mid\wedge(k,j))$. The same assertion holds with $\wedge$ replaced by $\vee$.

*Step 5: End of proof.* Let $F:E\to\mathbb{R}$ be some bounded and continuous function and set $G:=F\psi_\varphi$. Note that $m_N(\vee(k,j)))=m_N(\wedge(k,j))$ (by symmetry, or see (55) below). Using the identity in law from Step 4, we deduce that the integral of $F(h)$ against (46) can be written

$$(48)\qquad \sqrt{2N}\sum_{k\in[\delta 2N,(1-\delta)2N]}\sum_{j\in[-a\sqrt{2N},a\sqrt{2N}]}\frac{i}{2}m_N(\vee(k,j))\left(\Phi_v\left(\frac{k}{2N}\right)-\Phi_w\left(\frac{k}{2N}\right)\right)$$

$$(49)\qquad \times\int_E\sum_{X\in\{\wedge,\vee\}}G(T^X_{N,k}(h_1,h_2))\nu_{(0,0)\to(k-1,j)}(dh_1)\nu_{(0,j)\to(2N-k-1,0)}(dh_2).$$

Note that only those $j$ whose parity matches with that of $k-1$ contribute in the sum. Using in turn Lemma 4.6(3) together with the change of variables $r = k/(2N)$ and $y = \sqrt{2}j/\sqrt{2N}$ this equals

$$\frac{i}{2}\int_{r=0}^{1}\frac{dr}{\sqrt{2\pi r^3(1-r)^3}}\int_{y=-\sqrt{2}a}^{\sqrt{2}a}\frac{e^{-\frac{y^2}{2r(1-r)}}}{\sqrt{2\pi r(1-r)}}dy\frac{\Big(\Phi_v\Big(\frac{\lfloor r2N\rfloor}{2N}\Big)-\Phi_w\Big(\frac{\lfloor r2N\rfloor}{2N}\Big)\Big)}{\sqrt{2}} \tag{50}$$

$$\times J_N\left(\lfloor r2N\rfloor,\left\lfloor\frac{\sqrt{2N}y}{\sqrt{2}}\right\rfloor,G\right)(1+o(1)) \tag{51}$$

where

$$J_N(k,j,G) := \frac{1}{2}\int_{E\times E}\sum_{X\in\{\wedge,\vee\}}G\big(T^X_{N,k}(h_1,h_2)\big)\nu_{(0,0)\to(k-1,j)}(dh_1)\nu_{(0,j)\to(2N-k-1,0)}(dh_2). \tag{52}$$

Since $|J_N(r,y,G)|$ is bounded uniformly over all $N$, $r$ and $y$, Lemma 4.5 below and the dominated convergence theorem ensure that the whole expression we started from converges to

$$\frac{i}{2}\int_{E\times E}\int_{r=0}^{1}\frac{dr}{\sqrt{2\pi r^3(1-r)^3}}\int_{y=-\sqrt{2}a}^{\sqrt{2}a}\frac{e^{-\frac{y^2}{2r(1-r)}}}{\sqrt{2\pi r(1-r)}}dy \times G(T_r(h_1,h_2))\frac{(\varphi_v(r)-\varphi_w(r))}{\sqrt{2}}m_{(0,0)\to(1,y)}(dh_1)m_{(0,y)\to(1,0)}(dh_2).$$

Using the identity in law of Step 3, we deduce that this measure converges as $a\uparrow\infty$ towards

$$\frac{i}{2}\int_E\int_{r\in[\delta,1-\delta]}\frac{dr}{\sqrt{2\pi r^3(1-r)^3}}\frac{(\varphi_v(r)-\varphi_w(r))}{\sqrt{2}}G(u)m_{\mathrm{bridge}}(h_S)m_{\mathrm{exc}}(h_D\mid h_D(r)=0),$$

as required. □

**Lemma 4.5.** *For any bounded continuous function $G : E\to\mathbb{R}$, any $r\in(0,1)$ and $y\in\mathbb{R}$, $J_N(\lfloor r2N\rfloor,\lfloor\sqrt{2N}y/\sqrt{2}\rfloor,G)$ converges as $N\to\infty$ towards*

$$\int_E G(T_r(h_1,h_2))m_{(0,0)\to(1,y)}(dh_1)m_{(0,y)\to(1,0)}(dh_2).$$

*Proof.* Let $\widetilde{E}_N$ be the set of all pairs $(h_v,h_w)$ obtained by rescaling ordered lattice paths $\mathtt{v}\geq\mathtt{w}$. Let us point out that, contrary to $E_N$, we do not pin these lattice paths at sites 0 and $2N$ so that $E_N$ is a strict subset of $\widetilde{E}_N$. Let $\pi_N : E\to\widetilde{E}_N$ be a projection map as defined in Section 2. For $X\in\{\wedge,\vee\}$ and $r\in(0,1)$, it is straightforward to check that the maps $T^X_{N,\lfloor r2N\rfloor}$, $N\geq1$ are uniformly (over $N$) Lipschitz from $\widetilde{E}_N\times\widetilde{E}_N$ into $\widetilde{E}_N$. Of course, the Lipschitz constant depends on $r$ and blows up as $r$ approaches 0 or 1. Furthermore, for any given $y\in\mathbb{R}$, $T^X_{N,\lfloor r2N\rfloor}(\pi_Nh_1,\pi_Nh_2)$ converges to $T_r(h_1,h_2)$ $m_{(0,0)\to(1,y)}(dh_1)m_{(0,y)\to(1,0)}(dh_2)$-almost everywhere.

A slight generalisation of Lemma 4.3 shows the following weak convergences hold as $N\to\infty$

$$\nu_{(0,0)\to\left(\lfloor r2N\rfloor-1,\sqrt{2N}y/\sqrt{2}\right)}(dh_1)\longrightarrow m_{(0,0)\to(1,y)}(dh_1),$$
$$\nu_{\left(0,\sqrt{2N}y/\sqrt{2}\right)\to(2N-\lfloor r2N\rfloor-1,0)}(dh_2)\longrightarrow m_{(0,y)\to(1,0)}(dh_2).$$

We have all the ingredients to apply Lemma 2.10 and deduce that $J_N(\lfloor r2N\rfloor,\lfloor\sqrt{2N}y/\sqrt{2}\rfloor,G)$ converges to

$$\int_E G(T_r(h_1,h_2))m_{(0,0)\to(1,y)}(dh_1)m_{(0,y)\to(1,0)}(dh_2).$$

□

**Lemma 4.6.** *Fix $\delta>0$.*

$$\limsup_{N\to\infty}\sum_{k\in[\delta2N,(1-\delta)2N]}\big(m_N(\vee(k))+m_N(\wedge(k))\big)\sqrt{2N}<\infty. \tag{1}$$

$$\limsup_{a\to\infty}\limsup_{N\to\infty}\sum_{k\in[\delta2N,(1-\delta)2N]}\sum_{j\notin[-a\sqrt{2N},a\sqrt{2N}]} \times\big(m_N(\vee(k,j))+m_N(\wedge(k,j))\big)\sqrt{2N}=0. \tag{2}$$

(3) *Uniformly over all* $k \in [\delta 2N, (1-\delta)2N]$ *and all* $j \in [-(2N)^{2/3}, (2N)^{2/3}]$ *with the same parity as* $k-1$*, as* $N \to \infty$

$$m_N(\vee(k,j)) = m_N(\wedge(k,j) = \frac{1}{2\pi(2N)^2}\left(\frac{(2N)^2}{k(2N-k)}\right)^2 e^{-\frac{j^2}{2N}\frac{(2N)^2}{k(2N-k)}}(1+o(1)).$$

*Proof.* By the Karlin–McGregor formula, see for instance [15, Theorem 2.5], the number of configurations in $\Omega_N$ that have a contact point at $k$ with an upward (resp. downward) corner at $k$ and are such that $\mathfrak{v}(k-1) = j$ equals

$$C(k-1,j)C(2N-k-1,j), \tag{53}$$

where

$$C(k,j) = \binom{k}{(k+j)/2}^2 - \binom{k}{(k+j)/2-1}\binom{k}{(k+j)/2+1} \tag{54}$$

is the number of pairs or ordered simple random walk trajectories of length $k$, starting at 0 and ending at $j$. As a consequence

$$m_N(\vee(k,j)) = m_N(\wedge(k,j)) = \frac{C(k-1,j)C(2N-k-1,j)}{C(2N,0)}. \tag{55}$$

We fix $\delta > 0$ and we always work with $k \in [\delta 2N, (1-\delta)2N]$ so that $k \to \infty$ as $N \to \infty$. Define the function

$$f(x) = (1+x)\ln(1+x) + (1-x)\ln(1-x), \quad x \in [-1,1],$$

and note that $f(x) = x^2 + O(x^4)$ as $x \to 0$. Note also that it is a convex function so that $f(x) \geq x^2$ over $[-1,1]$. A tedious computation shows that, provided $(k-1) - j \to \infty$ as $N \to \infty$, we have

$$C(k-1,j) \sim 2^{2k-1}\frac{4k(k-1)}{\pi((k-1)^2-j^2)((k+1)^2-j^2)}e^{-(k-1)f\left(\frac{j}{k-1}\right)}, \quad N \longrightarrow \infty.$$

A straightforward computation then leads to (3), noticing that in this statement $j$ is taken to be negligible compared to $N$ and therefore to $k$.

To establish (1) and (2) we need to deal with values $j$ that can be arbitrarily close to $\pm(k-1)$ and $\pm(2N-k-1)$. Actually the contributions coming from $j \in \{-k+1, k-1, -2N+k+1, 2N-k-1\} =: \mathcal{J}$ are easy to deal with since, in any of these four cases, either $C(k-1,j) = 1$ or $C(2N-k-1,j) = 1$. We therefore exclude these four values from now on. The same arguments as above show that there exists some constant $c > 0$ such that for all $N$ large enough

$$C(k-1,j) \leq c2^{2k-1}\frac{4k(k-1)}{\pi((k-1)^2-j^2)((k+1)^2-j^2)}e^{-(k-1)f\left(\frac{j}{k-1}\right)},$$

and

$$C(2N-k-1,j) \leq c2^{4N-2k-1}\frac{4(2N-k)(2N-k-1)}{\pi((2N-k-1)^2-j^2)((2N-k+1)^2-j^2)}e^{-(2N-k-1)f\left(\frac{j}{2N-k-1}\right)},$$

and we deduce that (recall that $f(x) \geq x^2$)

$$\sum_{k\in[\delta 2N,(1-\delta)2N]}\sum_{j\notin[-a\sqrt{2N},a\sqrt{2N}]\cup\mathcal{J}}\big(m_N(\vee(k,j)) + m_N(\wedge(k,j))\big)\sqrt{2N}$$
$$\lesssim \sum_{k\in[\delta 2N,(1-\delta)2N]}\frac{1}{2N}\sum_{j\notin[-a\sqrt{2N},a\sqrt{2N}]}\frac{1}{\sqrt{2N}}e^{-\frac{j^2}{2N}\frac{1}{\delta(1-\delta)}},$$

which is bounded uniformly over all $N$ large and its limsup in $N$ goes to 0 as $a \to \infty$. This establishes (2). The bound (1) can be deduced from (2) and (3). □

4.4. **Tightness.** We aim at checking Assumption 2.3. Recall the Sobolev space $H^{-\beta}([0,1],\mathbb{R})$ introduced in Section 3.5. We define $H^{-\beta}([0,1],\mathbb{R}^2)$ as the set of all pairs $(h_v, h_w)$ such that both $h_v$ and $h_w$ belong to $H^{-\beta}([0,1],\mathbb{R})$. To show that the sequence $(u)_N$ is tight in $\mathbb{D}([0,\infty), H^{-\beta}([0,1],\mathbb{R}^2))$, it suffices to prove that the sequences $(v)_N$ and $(w)_N$ separately are tight in $\mathbb{D}([0,\infty), H^{-\beta}([0,1],\mathbb{R}))$ for any$\beta > 1$. The proof follows from the Lyons–Zheng decomposition [23], exactly like in Section 3.5 for the zero-range process so we do not present the details.

We can now prove the following result.

**Theorem 4.7.** *The discrete process $(u(t), t \geq 0)$, starting from its stationary measure $m_N$, converges in law to the process $(u(t), t \geq 0)$, starting from $m$, in the space*

$$\mathbb{D}\big([0,\infty),\ H^{-\beta}([0,1],\mathbb{R}^2)\big)$$

*for any given $\beta > 1$.*

*Proof.* We aim at applying Theorem 2.6. Assumption 2.1 is the content of Proposition 4.2, the convergence in law required in Assumption 2.2 is checked in Lemma 4.3, Assumption 2.3 is discussed right above the statement of the theorem, Assumption 2.4 is discussed in Section 4.2 and Assumption 2.5 is proven in Proposition 4.4. □

**Remark 4.8.** The dynamics of the two reflected interfaces can be easily generalised [19, 28] to $k \geq 2$ reflected interfaces, $\mathtt{v}_{(1)} \geq \cdots \geq \mathtt{v}_{(k)}$ pinned at zero at their endpoints: $\mathtt{v}_{(i)}(0) = \mathtt{v}_{(i)}(2N) = 0$ for all $1 \leq i \leq k$. Assign the flip rates as follows. If interface $\mathtt{v}_{(i)}$ has an upward (resp. downward) corner at $1 \leq x \leq 2N-1$, call $N_i(x) \geq 0$ the number of interfaces with index $j$ strictly larger (resp. smaller) than $i$, such that $\mathtt{v}_{(j)}(y) = \mathtt{v}_{(i)}(y)$, $y = x, x+1, x-1$. In other words, the $j$ interfaces below (resp. above) $\mathtt{v}_{(i)}$ have a contact point with it at position $x$. Then, the corner of $\mathtt{v}_{(i)}$ is flipped, together with that of the other $j$ mentioned interfaces, with rate $(2N)^2/(2(N_i(x)+1))$. See Figure 3 for an illustration. The uniform measure over the collections of ordered interfaces $\mathtt{v}_{(i)}, 1 \leq i \leq k$, is stationary and reversible for the process. Also, as already mentioned above, the existence of a monotone global coupling under which the $L^1$ distance contracts is known [28]. It is very likely that, modulo a certain amount of technical work (especially for the "discrete characterization and convergence", which requires applying the Karlin–McGregor formula for $k$ paths) one can use our method to show that, for $k$ fixed and $N \to \infty$, the stationary dynamics converges to a collection of $k$ reflected SPDEs, that generalises (36). A much more challenging situation is that where $k$ is of order $N$. In this case, the process can be seen as a dynamics for rhombus tilings of an $N \times N$ planar domain [28] (the interfaces $\mathtt{v}_{(i)}$ representing the level lines of the height function), and we expect that the scaling limit of the stationary process is a two-dimensional stochastic heat equation with additive noise.





FIGURE 3. An example of allowed configuration for $k = 3$ reflected interfaces. At positions $x_1, x_3$ there is a triple contact with a downward corner, and at $x_2$ a double contact with an upward corner. The rate at which $v_{(3)}$ flips upward at position $x_1$ (together with the other two interfaces) is $(2N)^2/6$, since $N_3(x_1) = 2$. On the other hand, $v_{(2)}$ can also flip upward at $x_1$ (together with $v_{(1)}$) and it does so with rate $(2N)^2/4$.

## 5. $\nabla\phi$ INTERFACES WITH CONVEX POTENTIAL IN ONE DIMENSION

To conclude, let us present yet another example of interesting Markov process for which our general method should apply. Given a convex function $V : \mathbb{R} \mapsto \mathbb{R}$ bounded from below, consider the one-dimensional $\nabla\phi$ interface model described by the height function $h = (h_i)_{i=0,1,\dots,N}$ with boundary conditions $h_0 = h_N = 0$ and with Boltzmann–Gibbs measure given as

$$m_N(dh) \propto e^{-\sum_{i=1}^{N} V(h_i - h_{i-1})} \prod_{i=1}^{N-1} dh_i \tag{56}$$

(the potential $V$ should increase at least linearly at $\pm\infty$, so that the measure is normalisable). One can consider two other variants of this model:

- the one with "hard wall", where $m_N$ is restricted to the subspace where $h_i \geq 0$ for every $i$;
- the discrete version (with or without the "hard wall" constraint $h_i \geq 0$) where the heights $h_i$ are restricted to integer values and the Lebesgue measure $\prod_{i=1}^{N-1} dh_i$ is replaced by the counting measure.

The following dynamics (Gibbs sampler) is reversible with respect to $m_N$: with rate $N^2$, the height $h_i$, $1 \le i \le N-1$ is updated to a new value sampled from $m_N(\cdot|h_{i\pm1})$, i.e., the equilibrium measure conditioned to the value of the heights in neighboring positions.

Let us point out that the dynamics with continuous heights and hard wall constraint can be seen as a Poissonian version of the dynamics considered by Funaki and Olla [11].

For the dynamics with continuous heights and no hard-wall constraint, [5] identified precisely the spectral gap and proved the occurrence of total variation cut-off, under the assumption that $V$ is of at most polynomial growth at infinity. For the case with discrete heights and hard-wall constraint, the mixing time was identified in [25, Section 3], up to a global prefactor.

We believe that the method of the present paper allows to prove that, under the same assumptions on $V$, the stationary process of suitably rescaled height fluctuations $\widetilde{h}_N(t,x) = N^{-1/2}h_{\lfloor xN \rfloor}(t), x \in [0,1], t \ge 0$ converges in distribution to either the stochastic heat equation with additive noise (if the hard wall constraint is absent) or the Nualart–Pardoux equation (35) (if the hard wall constraint is present). We do not expect the actual proof to be harder than or significantly different from that of either the zero-range process or of the pair of reflected interfaces, given above, but we preferred not to work it out, in order to keep this work within a reasonable length.

Let us just mention that it is easy to check that, because of the assumed convexity of $V$, these interface dynamics admit a global monotone coupling, and that under this coupling, the $L^1$ distance between configurations contracts (this holds both with and without hard-wall constraint). This allows to obtain equicontinuity of the sequence of resolvents. The technical work that is left concerns the discrete integration by parts, that involves arguments based on the local limit theorem for random walk (and for random walks conditioned to be positive) [3, 4].

**Data availability.** We do not analyse or generate any datasets.

**Competing interests.** The authors have no competing interests to declare that are relevant to the content of this article.

## References


[1] Bouleau, Nicolas; Hirsch, Francis Dirichlet forms and analysis on Wiener space, De Gruyter Studies in Mathematics, 14, Walter de Gruyter, 1991, x+325 pages | DOI | MR | Zbl

[2] Billingsley, Patrick Convergence of probability measures, Wiley Series in Probability and Statistics, John Wiley & Sons, 1999, x+277 pages (A Wiley-Interscience Publication) | DOI | MR | Zbl

[3] Bryn-Jones, A.; Doney, Ronald A. A functional limit theorem for random walk conditioned to stay non-negative, J. Lond. Math. Soc. (2), Volume 74 (2006) no. 1, pp. 244-258 | DOI | Zbl | MR

[4] Caravenna, Francesco; Chaumont, Loïc Invariance principles for random walks conditioned to stay positive, Ann. Inst. Henri Poincaré, Probab. Stat., Volume 44 (2008) no. 1, pp. 170-190 | DOI | MR | Zbl | Numdam

[5] Caputo, Pietro; Labbé, Cyril; Lacoin, Hubert Spectral gap and cutoff phenomenon for the Gibbs sampler of $\nabla\varphi$ interfaces with convex potential, Ann. Inst. Henri Poincaré, Probab. Stat., Volume 58 (2022) no. 2, pp. 794-826 | DOI | MR | Zbl

[6] Chang, Chih-Chung; Landim, Claudio; Olla, Stefano Equilibrium fluctuations of asymmetric simple exclusion processes in dimension $d \ge 3$, Probab. Theory Relat. Fields, Volume 119 (2001) no. 3, pp. 381-409 | DOI | MR | Zbl

[7] Caputo, Pietro; Martinelli, Fabio; Toninelli, Fabio Lucio Mixing times of monotone surfaces and SOS interfaces: a mean curvature approach, Commun. Math. Phys., Volume 311 (2012) no. 1, pp. 157-189 | DOI | Zbl | MR

[8] Debussche, Arnaud; Zambotti, Lorenzo Conservative stochastic Cahn–Hilliard equation with reflection, Ann. Probab., Volume 35 (2007) no. 5, pp. 1706-1739 | DOI | MR | Zbl

[9] Etheridge, Alison M.; Labbé, Cyril Scaling limits of weakly asymmetric interfaces, Commun. Math. Phys., Volume 336 (2015) no. 1, pp. 287-336 | DOI | MR | Zbl

[10] Fisher, Michael E. Walks, walls, wetting, and melting, J. Stat. Phys., Volume 34 (1984), pp. 667-729 | DOI | Zbl | MR

[11] Funaki, Tadahisa; Olla, Stefano Fluctuations for $\nabla\phi$ interface model on a wall, Stochastic Processes Appl., Volume 94 (2001) no. 1, pp. 1-27 | DOI | MR | Zbl

[12] Giacomin, Giambattista Random polymer models, Imperial College Press, 2007 | Zbl | DOI | MR

[13] Gillet, Florent Asymptotic behaviour of watermelons (2003) | arXiv | Zbl

[14] Gonçalves, Patrícia; Jara, Milton Nonlinear fluctuations of weakly asymmetric interacting particle systems, Arch. Ration. Mech. Anal., Volume 212 (2014) no. 2, pp. 597-644 | DOI | MR | Zbl

[15] Gorin, Vadim Lectures on random lozenge tilings, Cambridge Studies in Advanced Mathematics, 193, Cambridge University Press, 2021 | DOI | Zbl | MR

[16] Jara, Milton; Menezes, Otávio Non-equilibrium fluctuations of interacting particle systems (2018) | arXiv

[17] Kipnis, Claude; Landim, Claudio Scaling limits of interacting particle systems, Grundlehren der Mathematischen Wissenschaften, 320, Springer, 1999, xvi+442 pages | DOI | MR | Zbl

[18] Lenglart, Erik; Lépingle, Dominique; Pratelli, Maurizio Présentation unifiée de certaines inégalités de la théorie des martingales, Seminar on Probability, XIV (Paris, 1978/1979) (French) (Lecture Notes in Mathematics), Volume 784, Springer, 1980, pp. 26-52 (with an appendix by Lenglart) | DOI | MR | Zbl

[19] Luby, Michael; Randall, Dana; Sinclair, Alistair Markov chain algorithms for planar lattice structures, SIAM J. Comput., Volume 31 (2001) no. 1, pp. 167-192 | DOI | Zbl | MR

[20] Laslier, Benoît; Toninelli, Fabio Lucio Hydrodynamic limit equation for a lozenge tiling Glauber dynamics, Ann. Henri Poincaré, Volume 18 (2017) no. 6, pp. 2007-2043 | DOI | MR | Zbl

[21] Laslier, Benoît; Toninelli, Fabio Lucio Lozenge tiling dynamics and convergence to the hydrodynamic equation, Commun. Math. Phys., Volume 358 (2018) no. 3, pp. 1117-1149 | DOI | MR | Zbl
[22] Laslier, Benoît; Toninelli, Fabio Lucio The mixing time of the lozenge tiling Glauber dynamics, Ann. Henri Lebesgue, Volume 6 (2023), pp. 907-940 | Zbl | Numdam | DOI | MR
[23] Lyons, Terence J.; Zheng, Weian A crossing estimate for the canonical process on a Dirichlet space and a tightness result, Colloque Paul Lévy sur les processus stochastiques (Astérisque), Volume 157-158, Société Mathématique de France, 1988, pp. 249-271 | MR | Zbl | Numdam
[24] Ma, Zhi Ming; Röckner, Michael Introduction to the theory of (nonsymmetric) Dirichlet forms, Universitext, Springer, 1992, vi+209 pages | DOI | MR | Zbl
[25] Martinelli, Fabio; Sinclair, Alistair Mixing time for the solid-on-solid model, Proceedings of the forty-first annual ACM symposium on Theory of computing, ACM Press (2009), pp. 571-580 | DOI | Zbl
[26] Nualart, David; Pardoux, Étienne White noise driven quasilinear SPDEs with reflection, Probab. Theory Relat. Fields, Volume 93 (1992) no. 1, pp. 77-89 | DOI | Zbl | MR
[27] Petrov, Valentin V. Sums of independent random variables, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer, 1975, x+346 pages (Translated from the Russian by A. A. Brown) | MR | Zbl
[28] Wilson, David Bruce Mixing times of lozenge tiling and card shuffling Markov chains, Ann. Appl. Probab., Volume 14 (2004) no. 1, pp. 274-325 | DOI | Zbl | MR
[29] Zambotti, Lorenzo A Reflected Stochastic Heat Equation as Symmetric Dynamics with Respect to the 3-d Bessel Bridge, J. Funct. Anal., Volume 180 (2001) no. 1, pp. 195-209 | DOI | Zbl | MR
[30] Zambotti, Lorenzo Integration by parts formulae on convex sets of paths and applications to SPDEs with reflection, Probab. Theory Relat. Fields, Volume 123 (2002) no. 4, pp. 579-600 | DOI | Zbl | MR
[31] Zambotti, Lorenzo Fluctuations for a $\nabla\phi$ interface model with repulsion from a wall, Probab. Theory Relat. Fields, Volume 339 (2004) no. 3, pp. 315-339 | DOI | Zbl | MR
[32] Zambotti, Lorenzo Convergence of approximations of monotone gradient systems, J. Evol. Equ., Volume 6 (2006) no. 4, pp. 601-619 | DOI | MR | Zbl
[33] Zambotti, Lorenzo Random obstacle problems. Lecture notes from the 45th Probability Summer School held in Saint-Flour, 2015, Lecture Notes in Mathematics, 2181, Springer, 2017, ix+162 pages | DOI | MR | Zbl