

for all $\gamma \in \Gamma$ and $p \in P$. Here $|\gamma|$ denotes word length in Γ with respect to S and, for $1 \leq i \leq d$, $s_i(\gamma)$ denotes the i -th largest singular value of $\gamma \in \mathrm{SL}(d, \mathbb{R})$ with respect to the usual inner product on $\mathbb{R}^d$. A representation is called *Borel-Anosov* if it is $\{1, \dots, d-1\}$ -Anosov.

Anosov representations have been introduced by F. Labourie ([Lab06]) and have been the subject of many studies (cf. in particular [GGKW17], [KLP16], [KLP17], [KLP18a], [BPS19] and [Can] for a recent survey). It is widely accepted that non-trivial Anosov representations present the right analog of convex cocompact representations in higher rank. Many properties are suggested by this analogy. In another direction, we present in this note a phenomenon which is purely higher rank: there is a dimension gap for minimal invariant subsets for the action of $\rho(\Gamma)$ on the spaces of flags. The proof uses a new variational principle for the action of $\rho(\Gamma)$ on the spaces of partial flags (see below Theorem 1.7).

Let Q be a non-empty subset of $\{1, \dots, d-1\}$. We consider the space $\mathcal{F}_Q$ of partial flags with signature Q , endowed with a rotationally invariant Riemannian metric. Let us recall the definition of the Minkowski dimension of a set $\Lambda \subset \mathcal{F}_Q$: for $\varepsilon > 0$, let $N(\Lambda, \varepsilon)$ be the covering number of the set Λ by balls of radius ε in the metric of $\mathcal{F}_Q$. The *Minkowski (or Box) dimension* $\dim_B(\Lambda)$ is defined by

$$\dim_B(\Lambda) := \limsup_{\varepsilon \rightarrow 0} \frac{\log N(\Lambda, \varepsilon)}{\log(1/\varepsilon)}.$$

Recall that the Hausdorff dimension of a metric set is bounded above by its Box dimension. Our result is

THEOREM 1.1 (Dimension gap for minimal sets of Anosov representations)

With the above notations, let Λ be a minimal invariant set for the action of $\rho(\Gamma)$ on $\mathcal{F}_Q$. Assume that the representation ρ is P -Anosov for some $P \subset \{1, \dots, d-1\}$ and such that $\rho(\Gamma)$ is Zariski dense in $\mathrm{SL}(d, \mathbb{R})$. Then, with $M := \#(Q \cap P)$,

$$\dim_B(\Lambda) \leq \dim(\mathcal{F}_Q) - \frac{M-1}{2}.$$

In the case $d = 2$ the bound given by the above theorem is trivial. And, indeed, uniform discrete subgroups of $\mathrm{SL}(2, \mathbb{R})$ have full limit set. A dimension gap for the limit set of classical Schottky groups was proved in [Doy88]. We note, however, that there exist uniform discrete (and thus convex co-compact) subgroups of isometries of rank one symmetric spaces and thus there is no dimension gap for convex cocompact representations in rank one. On the other hand, there are many conditions that ensure that the limit set is a Lipschitz submanifold of the flag space (see the discussions in [PSW23]). Our result is easy in those cases.

If ρ is P -Anosov then it is also $P \cup P^*$ -Anosov for $P^* = \{d-p : p \in P\}$. Theorem 1.1 gives a non-trivial upper bound for the dimension of any minimal invariant set in $\mathcal{F}_Q$ for all Q with $\#(Q \cap (P \cup P^*)) \geq 2$. In particular, as soon as the representation is k -Anosov for some $k \neq d/2$ and Zariski dense, then the codimension of the limit set in the full flag space is at least $1/2$.

There are a few simple reductions which allow us to state our core result. Firstly, we observe that the bundle $\pi_{Q,Q \cap P} : \mathcal{F}_Q \rightarrow \mathcal{F}_{Q \cap P}$ is smooth, so that for any closed $\Lambda \subset \mathcal{F}_{Q \cap P}$, we have

$$\dim_B((\pi_{Q,Q \cap P})^{-1}(\Lambda)) = \dim_B(\Lambda) + \dim \mathcal{F}_Q - \dim \mathcal{F}_{Q \cap P}.$$

So, it suffices to show Theorem 1.1 for the case $Q = P$. In that case, the unique minimal invariant subset of $\mathcal{F}_P$ is the limit set Λ_ρ .

By considering $\Gamma/\text{Ker} \rho$, we may assume that the representation ρ is faithful. Furthermore, we may assume that Γ is non-elementary hyperbolic since, on the one hand, an Anosov subgroup of $\text{SL}(d, \mathbb{R})$ is hyperbolic ([KLP18b], [BPS19, Th. 3.2]) and, on the other hand, the limit set of an Anosov representation of an elementary (i.e., virtually cyclic) hyperbolic group is finite. Taking all this into account, we only have to prove

THEOREM 1.2 (Dimension gap for limit sets of Anosov representations). — *Let Γ be a non-elementary, hyperbolic, finitely generated group, $P \subset \{1, \dots, d-1\}$ with $M := \#P \geq 2$, ρ a faithful P -Anosov representation of Γ in $\text{SL}(d, \mathbb{R})$ with $\rho(\Gamma)$ Zariski dense in $\text{SL}(d, \mathbb{R})$ and $\Lambda_\rho \subset \mathcal{F}_P$ the limit set of the representation ρ . Then,*

$$\dim_B(\Lambda_\rho) \leq \dim(\mathcal{F}_P) - \frac{M-1}{2}.$$

It seems natural to conjecture that a dimension gap will exist between the limit set and the corresponding space of flags, for any Zariski dense representation into a reductive algebraic Lie group of rank $r \geq 2$ which is Anosov with respect to at least two simple roots.

Let $g \in \text{SL}(d, \mathbb{R})$. The numbers $\lambda_i(g) := \lim_n \frac{1}{n} \log s_i(g^n)$ are the logarithms of the moduli of the eigenvalues of g . Apply the Oseledets theorem in the trivial case of one matrix $g \in \text{SL}(d, \mathbb{R})$. The Oseledets decomposition is obtained from the Jordan decomposition by grouping together the spaces corresponding to the eigenvalues with the same modulus. These moduli are the $\exp \lambda_i(g)$.

In our setting, if $\gamma \in \Gamma$ (but a finite number), for all $p \in P$, $\lambda_p(\rho(\gamma)) - \lambda_{p+1}(\rho(\gamma)) \geq c|\gamma|$. We associate to every such $\gamma \in \Gamma$ an element $\eta(\rho(\gamma))$ of the space X_P of decompositions of $\mathbb{R}^d = E_1 \oplus \dots \oplus E_{M+1}$, with $\dim E_k = p_k - p_{k-1}$, namely

$$\eta(\rho(\gamma)) = E_1(\rho(\gamma)) \oplus \dots \oplus E_{M+1}(\rho(\gamma)),$$

with the property that for all $k = 1, \dots, M+1$, the space $E_k(\rho(\gamma))$ is generated by the Jordan spaces of $\rho(\gamma)$ such that the modulus $\exp \lambda$ of the eigenvalue satisfies⁽¹⁾

$$\exp \lambda_{p_{k-1}}(\rho(\gamma)) > \exp \lambda \geq \exp \lambda_{p_k}(\rho(\gamma)).$$

Observe that the space X_P can also be seen as the open $\text{SL}(d, \mathbb{R})$ -invariant set of pair of flag spaces in general position in $\mathcal{F}_P \times \mathcal{F}_{P^*}$.

⁽¹⁾With the convention that $p_0 = 0, \lambda_{p_0} = +\infty$ and $p_{M+1} := d$.

COROLLARY 1.3. — *Let Γ be a hyperbolic finitely generated group, $P \subset \{1, \dots, d-1\}$ with $M := \#P$. With the above notations, if ρ is a P -Anosov, Zariski dense representation of Γ in $\mathrm{SL}(d, \mathbb{R})$, set $\Omega_\rho \subset X_P$ the closure of the set $\{\eta(\rho(\gamma)); \gamma \in \Gamma\}$. Then,*

$$\dim_B(\Omega_\rho) \leq \dim X_P - M + 1.$$

Indeed, Corollary 1.3 is trivial for $d = 2$ and for $M = 1$. It is trivial as well if Γ is elementary. For $d \geq 3$ and $M \geq 2$, we show in Section 2.3 why Corollary 1.3 follows from Theorem 1.2.

The first non-trivial case for our results is when $d = 3$ and $P = \{1, 2\}$. So, let ρ be a Borel-Anosov representation of the finitely generated non-elementary group (Γ, S) in $\mathrm{SL}(3, \mathbb{R})$ and we can consider the limit set Λ_ρ of the action of $\rho(\Gamma)$ on the space $\mathcal{F}$ of complete flags in $\mathbb{R}^3$. Besides, for all $\gamma \in \Gamma$, the matrix $\rho(\gamma)$ admits three distinct eigenspaces written $(E_1(\rho(\gamma)), E_2(\rho(\gamma)), E_3(\rho(\gamma)))$ in the order of the absolute values of the eigenvalues. Let Ω_ρ denote the closure in $\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$ of the $\{(E_1(\rho(\gamma)), E_2(\rho(\gamma)), E_3(\rho(\gamma))), \gamma \in \Gamma\}$. We obtain, from Theorem 1.2 and Corollary 1.3:

COROLLARY 1.4

$$\dim_B(\Lambda_\rho) \leq \frac{5}{2} < 3 = \dim(\mathcal{F}), \quad \dim_B(\Omega_\rho) \leq 5 < 6 = \dim(\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2).$$

Let Γ be a surface group. For the Hitchin component, the dimension of the projective limit set is 1 (see [Lab06]) and our result is not new. It is new and unexpected, as far as the authors are aware, for the Zariski dense representations that lie in the same connected component as the Teichmüller times Identity representations, i.e., in the Barbot component (cf. [Bar10]).

1.2. STRATEGY OF THE PROOF OF THEOREM 1.2

1.2.1. Random walk entropy. — Let $\mathcal{M}$ be the set of probability measures μ on Γ with countable support and with finite first moment, meaning

$$\sum_{\gamma \in \Gamma} \mu(\gamma) |\gamma| < +\infty,$$

such that the semi-group Γ_μ generated by the support of μ is non-elementary (i.e., contains two independent loxodromic elements). By [KV83], if $\mu \in \mathcal{M}$ then μ has finite Shannon entropy, meaning

$$H(\mu) = - \sum_{\gamma \in \Gamma} \mu(\gamma) \log(\mu(\gamma)) < +\infty.$$

The random walk entropy h_μ of $\mu \in \mathcal{M}$ is defined by

$$(1) \quad h_\mu = \lim_{n \rightarrow +\infty} \frac{1}{n} H(\mu^{*n}),$$

where μ^{*n} is the n -fold convolution of μ with itself.

1.2.2. *Lyapunov exponents.* — Denote by $\mathfrak{a}^+$ the cone

$$\mathfrak{a}^+ := \{a \in \mathbb{R}^d : a_1 \geq \dots \geq a_d, \sum a_i = 0\}.$$

For $g \in \mathrm{SL}(d, \mathbb{R})$ one has $\log S(g) \in \mathfrak{a}^+$, where

$$\log S(g) := (\log s_1(g), \dots, \log s_d(g)).$$

The Lyapunov exponents $\lambda(\rho_*\mu) = \{\lambda_i(\rho_*\mu), 1 \leq i \leq d\} \in \mathfrak{a}^+$ of the random walk (Γ, μ) in the representation ρ are defined by

$$(2) \quad \lambda(\rho_*\mu) := \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{\gamma \in \Gamma} \mu^{*n}(\gamma) \log(S(\rho(\gamma))).$$

The limits exist by the sub-additivity of the sequences

$$\{\sum_{j \leq i} \sum_{\gamma \in \Gamma} \mu^{*n}(\gamma) \log(s_j(\rho(\gamma)))\}_{n \in \mathbb{N}}.$$

1.2.3. *Lyapunov and Falconer dimension.* — We say a pair (i, j) , $0 < i < j \leq d$, is separated by $p \in P$ if $i \leq p < j$. For each $p \in P$ let S_p be the set of pairs separated by p , and define $S(P) = \bigcup_{p \in P} S_p$. On the cone $\mathfrak{a}^+$ we define the roots

$$\alpha_{i,j}(a) = a_i - a_j,$$

for $i < j$ and notice that they are non-negative on $\mathfrak{a}^+$. Lyapunov and Falconer dimensions are special values of Lyapunov and Falconer functionals on $\mathfrak{a}^+$.

We define, for $h \geq 0$ the Lyapunov functional on $\mathfrak{a}^+$ by

$$\begin{aligned} L_h^P(a) = \text{maximum of } & \sum_{(i,j) \in S(P)} r_{i,j} \\ \text{subject to } & 0 \leq r_{i,j} \leq 1 \quad \text{and} \quad \sum_{(i,j) \in S(P)} r_{i,j} \alpha_{i,j}(a) \leq h. \end{aligned}$$

For $r \geq 0$ we also define the Falconer functional on $\mathfrak{a}^+$ by

$$\begin{aligned} F_r^P(a) = \text{minimum of } & \sum_{(i,j) \in S(P)} r_{i,j} \alpha_{i,j}(a) \\ \text{subject to } & 0 \leq r_{i,j} \leq 1 \quad \text{and} \quad \sum_{(i,j) \in S(P)} r_{i,j} \geq r. \end{aligned}$$

LEMMA 1.5 (Duality between Falconer and Lyapunov functionals). — *Given $a \in \mathfrak{a}^+$, for all $r, h \geq 0$ one has $F_r^P(a) \leq h$ if and only if $L_h^P(a) \geq r$.*

Furthermore, let $d(a)$ be the number of pairs (i, j) in $S(P)$ with $\alpha_{i,j}(a) = 0$. Then, $F_r^P(a) = 0$ for $r \in [0, d(a)]$ and $r \mapsto F_r^P(a)$ is an increasing homeomorphism from $[d(a), D]$ to $[0, F_D^P(a)]$ where $D = \#S(P) = \dim(\mathcal{F}_P)$, and $h \mapsto L_h^P(a)$ is its inverse.

Proof. — To establish the first claim observe that directly from the definitions, both conditions $F_r^P(a) \leq h$ and $L_h^P(a) \geq r$ are equivalent to there being some choice of $r_{i,j} \in [0, 1]$ for $(i, j) \in S(P)$ such that

$$\sum_{(i,j) \in S(P)} r_{i,j} \alpha_{i,j}(a) \leq h \quad \text{and} \quad \sum_{(i,j) \in S(P)} r_{i,j} \geq r.$$

Moreover, in computing $F_r^P(a)$, we can always take $r_{i,j} = 1$ for the pairs (i, j) in $S(P)$ with $\alpha_{i,j}(a) = 0$. It follows that $F_r^P(a) = 0$ for $r \in [0, d(a)]$. Then, there exists $\varepsilon > 0$ with $\alpha_{i,j}(a) > \varepsilon$ for all the other pairs (i, j) . It follows immediately that $F_{r+s}^P(a) > F_r^P(a) + \varepsilon s$ for all $d(a) \leq r < r + s \leq D$. Hence $r \mapsto F_r^P(a)$ is increasing on $[d(a), D]$. Since $F_{d(a)}^P(a) = 0$, $r \mapsto F_r^P(a)$ it is an increasing homeomorphism as claimed.

Similarly, picking $\varepsilon > 0$ small enough so $\alpha_{i,j}(a) < \varepsilon^{-1}$ for all $(i, j) \in S(P)$, it follows that $L_{h+s}^P(a) > L_h^P(a) + \varepsilon s$ for all $0 \leq h < h + s < F_D^P(a)$. Hence, $h \mapsto L_h^P(a)$ is an increasing homeomorphism as well.

Let $f : I \rightarrow J$ and $g : J \rightarrow I$ be the two homeomorphisms under consideration where $I = [d(a), D]$ and $J = [0, F_D^P(a)]$. By our first claim we have $g(f(r)) \geq r$ for all $r \in I$ and $f(g(h)) \leq h$ for all $h \in J$. However, substituting $r = f^{-1}(h)$, the first inequality implies $g(h) \geq f^{-1}(h)$ for all $h \in J$, from which we obtain $f(g(h)) \geq h$ for all $h \in J$. Hence, we have $f(g(h)) = h$ for all $h \in J$ as required. $\square$

Following [DO80] and [KY79], we define the Lyapunov dimension of $\rho_*\mu$ on $\mathcal{F}_P$ by

$$(3) \quad \dim_{LY}^P(\rho_*\mu) := L_{h_\mu}^P(\lambda(\rho_*\mu)).$$

In the spirit of [Fal88], we define the Falconer dimension $\dim_F^P(\rho)$ of ρ relative to $\mathcal{F}_P$ as the critical parameter $r \geq 0$ for convergence of the series

$$(4) \quad \Phi_\rho^P(r) := \sum_{\gamma \in \Gamma} \exp(-(F_r^P \circ \log S(\rho(\gamma)))).$$

1.2.4. Proof of Theorem 1.2. — Theorem 1.2 is a direct consequence of the three following results, all under the assumptions of Theorem 1.2 and using the above notations:

THEOREM 1.6 (Upper bound on Lyapunov dimension). — *For all $\mu \in \mathcal{M}$, one has*

$$\dim_{LY}^P(\rho_*\mu) \leq \dim(\mathcal{F}_P) - \frac{M-1}{2}.$$

THEOREM 1.7 (Variational principle). — *One has*

$$(5) \quad \dim_F^P(\rho) = \sup_{\mu \in \mathcal{M}} \dim_{LY}^P(\rho_*\mu).$$

THEOREM 1.8 (Inequality between Minkowski and Falconer dimension). — *One has*

$$\dim_B(\Lambda_\rho) \leq \dim_F^P(\rho).$$

The key observation behind Theorem 1.6 is that the entropy h_μ is realized as the Furstenberg entropy of the action of $\rho(\Gamma)$ on the Grassmannian $\text{Gr}(p, \mathbb{R}^d)$ for any $p \in P$. This follows from the P -Anosov property and the identification of the Poisson boundary and geometric boundary for μ -random walks on Γ ([Kai00]). The inequality in Theorem 1.6 is then a consequence of the inequality between Furstenberg entropy and the sum of the relevant exponents (see [LL24] and Section 3).

Concerning Theorem 1.7, the fact that

$$(6) \quad \dim_{LY}^P(\rho_*\mu) \leq \dim_F^P(G)$$

is classical. We recall the proof in Section 2.2.2.

The converse is due to Yuxiang Jiao, Jialun Li, Wenyu Pan and Disheng Xu ([JLPX24]) in the case when the representation is Borel-Anosov. In [JLPX24], given $\varepsilon > 0$, the authors construct a free semi-group in Γ that is rich enough that the uniform probability measure μ_ε on the generators satisfies $\dim_{LY}(\rho_*\mu_\varepsilon) \geq \dim_F(\rho) - \varepsilon$. The complete dominated splitting is used to control the almost additivity of the singular values. Here, we consider a general subset P and we assume that the representation $\rho(\Gamma)$ is Zariski dense in $\mathrm{SL}(d, \mathbb{R})$. In the spirit of [Gou22], [AGG⁺23] and [JLPX24], we prove in the appendix a general proposition about free sub-semigroups in hyperbolic groups (see the appendix for the definition of a quasi-geodesic set; see also [Yan19] for a construction of free sub-semi-groups with convexity properties in a more general context.)

PROPOSITION 1.9 (= Proposition A.1). — *Let (Γ, S) be a non-elementary hyperbolic group with a finite symmetric generator S . Then there exists a semigroup $S \subset \Gamma$, a finite set $F \subset \Gamma$ and mappings $L, R : \Gamma \rightarrow F$ with the following properties:*

- (1) S generates Γ as a group,
- (2) S is quasi-geodesic, and
- (3) $L(\gamma)\gamma R(\gamma) \in S$ for all $\gamma \in \Gamma$.

Then, by Zariski density and [AMS95], there is a measurable complete splitting and, following [Ben96] and [BS21], we are able to control the almost additivity of the singular values. Observe that, given a recent result of Dey ([Dey22]), only groups that are virtually free or virtually surface groups can have Borel Anosov representations in $\mathrm{SL}(d, \mathbb{R})$, whereas a priori, many hyperbolic groups may have partially Anosov representations

Finally, the proof of Theorem 1.8 is done in Section 5. Similarly to other previous works (see [Zha97], [BCH10], [PSW21], [GMT23], [FS23]), it rests on covering the limit set in $\partial\Gamma$ by shadows and counting the covering numbers of the images of these shadows in Λ_ρ .

1.3. RELATED FACTS. — For convex cocompact groups in $\mathrm{SO}(n, 1)$, D. Sullivan proved in [Sul79] that the Hausdorff dimension and the Minkowski dimension of the limit set coincide with the Poincaré exponent of the group (which is the Falconer dimension in that case). This result can be extended to more general cases if one uses a conformally invariant metric on the boundary (see [Lin04], [DK22]). The problem is more delicate for the rotation invariant Riemannian metric (see e.g. [Duf17] for the $\mathrm{SU}(n, 1)$ case).

Following the pioneer work of Falconer ([Fal88]), there are many results comparing Hausdorff dimension and Falconer dimension for affine IFS. The variational principle (5) for IFS can be found in [Fal88] and [Käe04]. Under the form (5), it is due to

Ian D. Morris and Pablo Shmerkin ([MS19]) in the case of dimension 2, Ian D. Morris and Çağrı Sert [MS21] in general.

Given Theorems 1.7 and 1.8, a natural step to prove $\dim_H(\Lambda_\rho) = \dim_B(\Lambda_\rho) = \dim_F^P(\rho)$ for P -Anosov representations is finding necessary conditions on a given probability μ on Γ and a $\rho_*(\mu)$ -stationary probability measure ν on $\mathcal{F}_P$ for having

$$(7) \quad \dim_{\text{loc}} \nu = \dim_{LY}^P(\rho_*\mu).$$

Here, $\dim_{\text{loc}} \nu$ is the ν -a.e. constant value of $\lim_{\varepsilon \rightarrow 0} \log \nu(B(x, \varepsilon)) / \log \varepsilon$ (cf. [LL24]). Relation (7) for IFS is a famous problem, with many deep contributions (most relevant for us are [Fal88], [JPS07], [Hoc14], [BV19], [Var19], [BHR19], [HR22], [FS22], [Rap24], [MS21]). For random walks on discrete subgroups of $\text{SL}(d, \mathbb{R})$, relation (7) is classical when $d = 2$ (see [HS17] for the general non-discrete case). [LL23] prove relation (7) for the Hitchin component of the representation of a surface group in $\text{SL}(d, \mathbb{R})$. In [LPX23], Jialun Li, Wenyu Pan and Disheng Xu prove relation (7) for $d = 3$ when the representation ρ is Borel-Anosov and $\rho(\Gamma)$ is Zariski dense. They also observe that relation (7) may be wrong when the Zariski closure of $\rho(\Gamma)$ is conjugated to $\text{SL}(2, \mathbb{R})$. See [DS22] for the case of $\text{SU}(2, 1)$.

Acknowledgments. — The authors would like to thank Jairo Bochi, León Carvajales, Jialun Li, Rafael Potrie and Andrés Sambarino for many helpful conversations. We also thank the referees for their numerous remarks.

2. RANDOM WALK ENTROPY

Let Γ be a non-elementary finitely generated word hyperbolic group, $P \subset \{1, \dots, d-1\}$ with $M := \#P \geq 2$, ρ a P -Anosov representation of Γ in $\text{SL}(d, \mathbb{R})$. We let $\mathcal{M}$ denote the set of probability measures μ on Γ with finite first moment

$$\sum_{\gamma} |\gamma| \mu(\gamma) < +\infty,$$

and such that the semi-group Γ_μ generated by the support of μ contains two independent loxodromic elements.

In this section, we prove that the random walk entropy h_μ given by (1) coincides with the Furstenberg entropy (see below) of the stationary measure on the Gromov boundary of Γ and its images on adequate flag spaces. We also relate the Falconer dimension and the random walk entropy to the growth indicator defined by [Qui02]. To conclude the section, we prove Corollary 1.3.

2.1. FURSTENBERG ENTROPY. — Recall that a probability measure ν on a compact space X on which Γ acts continuously is said to be μ -stationary, where μ is a probability measure on Γ , if and only if

$$\nu = \sum_{\gamma \in \Gamma} \mu(\gamma) \gamma_* \nu.$$

The Furstenberg entropy of a μ -stationary measure is defined as

$$\kappa(\mu, \nu) = \sum_{\gamma \in \Gamma} \mu(\gamma) \int_X \log \left(\frac{d\gamma_* \nu}{d\nu}(\gamma x) \right) d\nu(x),$$

or $+\infty$ if the Radon-Nikodym derivative in the integral does not exist for some γ in the support of μ .

In this section we show that for $\mu \in \mathcal{M}$, the Furstenberg entropy of the natural μ -stationary measures on the Gromov boundary $\partial\Gamma$, the Grassmannian manifolds of p -dimensional subspaces of $\mathbb{R}^d$ for $p \in P$, and the space of partial flags $\mathcal{F}_P$ all coincide with the random walk entropy h_μ .

This particular feature of Anosov representations is useful since the Furstenberg entropy is what occurs in the dimension formulas of [LL24].

2.1.1. Stationary measure on the Gromov boundary $\partial\Gamma$

THEOREM 2.1. — *For each $\mu \in \mathcal{M}$ there exists a unique μ -stationary measure ν_μ on the Gromov boundary $\partial\Gamma$ of Γ , and its Furstenberg entropy is given by $\kappa(\mu, \nu_\mu) = h_\mu$.*

Proof. — If $\gamma_{-1}, \dots, \gamma_{-n}, \dots$ are i.i.d. random elements in Γ with common distribution μ . Then by [Kai00, Th. 7.6] there exists a random limit point

$$X = \lim_{n \rightarrow +\infty} \gamma_{-1} \cdots \gamma_{-n} \in \partial\Gamma,$$

almost surely, and the distribution ν_μ of X is the unique μ -stationary measure on $\partial\Gamma$.

Furthermore, $(\partial\Gamma, \nu_\mu)$ is isomorphic to the Poisson boundary of (Γ, μ) by [Kai00, Th. 7.7].

The Furstenberg entropy $\kappa(\mu, \nu_\mu)$ is equal to the difference between h_μ and the entropy of the random walk $\gamma_1, \gamma_1\gamma_2, \dots$ conditioned on X (see [KV83, Cor. 2]). However, because $(\partial\Gamma, \nu)$ is the Poisson boundary this conditional entropy is zero ([Kai00, Th. 4.3 and 4.5]) and therefore $h_\mu = \kappa(\mu, \nu_\mu)$ as claimed. $\square$

2.1.2. Boundary maps. — Let $p \in P$ and $\gamma \in \Gamma$ be such that $s_p(\rho(\gamma)) > s_{p+1}(\rho(\gamma))$. There exists a unique p -dimensional subspace $\xi^p(\gamma) \subset \mathbb{R}^d$ on which p -dimensional volume is most contracted by $\rho(\gamma)^{-1}$.

We let $\xi(\gamma) = (\{0\} \subset \xi^{p_1}(\gamma) \subset \dots \subset \xi^{p_M}(\gamma)) \subset \mathbb{R}^d \in \mathcal{F}_P$, where $P = \{p_1 < \dots < p_M\}$, whenever all $\xi^p(\gamma)$ are well defined. Since ρ is P -Anosov this is the case outside of a finite subset of Γ .

We refer to [GdlH90, Chap. 7] for basic properties of the Gromov compactification of Γ . We briefly recall that Γ with its word metric is a proper geodesic metric space. The group Γ is word hyperbolic, so there exists $\delta > 0$ such that for every geodesic triangle each side is contained in the δ -neighborhood of the other two. The Gromov boundary $\partial\Gamma$ is the set of equivalence classes of word geodesic rays in Γ , where two rays are equivalent if they are at bounded Hausdorff distance. A basis of neighborhoods of a point $x \in \partial\Gamma$ in the Gromov compactification $\bar{\Gamma} = \Gamma \cup \partial\Gamma$ is defined by taking for each $C > 0$ the set

$$N(x, C) = \{x\} \cup \{y \in \bar{\Gamma} : \min_n |\alpha_n| > C \text{ for all geodesics } \alpha \text{ joining } x \text{ and } y\}.$$

PROPOSITION 2.2 ([GGKW17]). — *The map ξ defined above extends continuously to the Gromov boundary $\partial\Gamma$. Furthermore, $\xi^p : \partial\Gamma \rightarrow \text{Gr}(p, \mathbb{R}^d)$ is injective for each $p \in P$.*

See [Can, Prop. 30.3] for a direct proof.

2.1.3. *Dynamical stationary measures.* — Fix $\mu \in \mathcal{M}$, let $\gamma_{-1}, \dots, \gamma_{-n}, \dots$ be i.i.d. random elements of Γ with common distribution μ , $\chi_1 > \dots > \chi_N$ be the distinct Lyapunov exponents of μ , and $d_1, \dots, d_N$ their multiplicities. Let

$$A = \{d_1, d_1 + d_2, \dots, d_1 + \dots + d_{N-1}\}.$$

By Oseledets theorem ([Ose68]), for each $a \in A$, there is a random limit

$$U_a = \lim_{n \rightarrow +\infty} \xi^a(\gamma_{-1} \cdots \gamma_{-n}),$$

almost surely.

The collection U of U_a for $a \in A$ is a random element in the space of flags of signature A . Its distribution, and the projections of its distribution to coarser partial flag spaces are what were called dynamical stationary measures in [LL24]. Recall from Section 1.2.2 the definition of the Lyapunov exponents $\lambda_i(\rho_*\mu)$. Since $\lambda_p(\rho_*\mu) > \lambda_{p+1}(\rho_*\mu)$ for all $p \in P$ we have that $\xi_*\nu_\mu$ is the dynamical stationary measure on $\mathcal{F}_P$, and $\xi_*^p\nu_\mu$ is the dynamical stationary measure on the Grassmannian of p -dimensional subspaces of $\mathbb{R}^d$ for each $p \in P$.

2.1.4. *Furstenberg entropy*

THEOREM 2.3. — *For each $\mu \in \mathcal{M}$ one has*

$$h_\mu = \kappa(\mu, \xi_*\nu_\mu) = \kappa(\mu, \xi_*^p\nu_\mu),$$

for all $p \in P, p \neq 0, d$.

Proof. — This is immediate from the injectivity of the maps ξ^p given by Proposition 2.2. $\square$

2.2. GROWTH INDICATOR FUNCTION

2.2.1. *Growth function and Falconer dimension.* — Given $a \in \mathfrak{a}^+$ we define the growth indicator at a by

$$(8) \quad \psi_\rho(a) = \|a\| \inf_{\mathcal{C} \ni a} \limsup_{T \rightarrow \infty} \frac{1}{T} \log \#\{\gamma \in \Gamma, \log S(\rho(\gamma)) \in \mathcal{C}, \|\log S(\rho(\gamma))\| \leq T\},$$

where the infimum is over all open subcones $\mathcal{C} \subset \mathfrak{a}^+$ that contain a . The definition does not depend on the choice of the norm $\|\cdot\|$ on $\mathfrak{a}^+$ ([Qui02]).

THEOREM 2.4 (Quint, [Qui02]). — *The growth indicator function ψ_ρ is concave and $\{\psi_\rho > 0\}$ is the interior of the limit cone defined by*

$$\mathcal{L}_\rho = \lim_{T \rightarrow +\infty} \frac{1}{T} \log S(\rho(\Gamma)),$$

where the limit is taken in the Hausdorff topology on closed sets.

Recall that we defined the Falconer dimension $\dim_F^P(\rho)$ as the critical value of the series $r \mapsto \sum_{\gamma \in \Gamma} \exp(-F_r^P \circ \log S(\rho(\gamma)))$. We have (see [Sam14, Lem. 4.2], [Qui02, Lem. III.1.3]):

LEMMA 2.5. — Assume $r \geq 0$ is such that there is $a \in \mathfrak{a}^+$ inside the interior of the limit cone $\mathcal{L}_\Gamma$ with $\psi_\rho(a) > F_r^P(a)$. Then,

$$r \leq \dim_F^P(\rho).$$

2.2.2. *Inequality between Lyapunov and Falconer dimensions.* — Recall that, given $\mu \in \mathcal{M}$ we defined the Lyapunov dimension by

$$\dim_{LY}^P(\rho_*\mu) = L_{h_\mu}(\lambda(\rho_*\mu)).$$

We will now prove relation (6), which is the following statement:

$$\sup_{\mu \in \mathcal{M}} \dim_{LY}^P(\rho_*\mu) \leq \dim_F^P(\rho).$$

For this purpose we first need the following version of the fundamental inequality for random walks, which is essentially due to Guivarc'h.

LEMMA 2.6 (Fundamental inequality). — For all $\mu \in \mathcal{M}$ one has

$$h_\mu \leq \psi_\rho(\lambda(\rho_*\mu)).$$

Proof. — Consider the sequence space $\Omega = \Gamma^{\mathbb{Z}}$ with the probability measure $m = \mu^{\mathbb{Z}}$ and let $g_n : \Omega \rightarrow \Gamma$ be the projection onto the n -th coordinate. By the subadditive ergodic theorem [Ose68] one has

$$\lambda(\rho_*\mu) = \lim_{n \rightarrow +\infty} \frac{1}{n} \log S(\rho(g_n(\omega) \cdots g_1(\omega))),$$

for m -almost every $\omega \in \Omega$. From the Shannon-McMillan-Breiman theorem for random walk entropy [KV83] we have

$$h_\mu = - \lim_{n \rightarrow +\infty} \frac{1}{n} \log \mu^{*n}(g_n(\omega) \cdots g_1(\omega)),$$

for m -a.e. $\omega \in \Omega$, where μ^{*n} is the n -fold self-convolution of μ and also the distribution of the random product $g_1 \cdots g_n$.

Fix $\varepsilon > 0$ and let $A_n \subset \Gamma$ be the set of elements satisfying

$$n\lambda(\rho_*\mu) - n\varepsilon < \log S(\gamma) < n\lambda(\rho_*\mu) + n\varepsilon,$$

and $B_n \subset \Gamma$ consist of those elements satisfying

$$\mu^{*n}(\gamma) < \exp(-n(h_\mu - \varepsilon)).$$

From the almost sure equalities above we have

$$\lim_{n \rightarrow +\infty} \mu^{*n}(A_n \cap B_n) = 1.$$

In particular

$$\#(A_n \cap B_n) > \mu(A_n \cap B_n) e^{n(h_\mu - \varepsilon)} > \frac{1}{2} e^{n(h_\mu - \varepsilon)}$$

for all n large enough.

This implies that for all $\varepsilon > 0$ and n large enough, there are at least $\frac{1}{2}e^{n(h_\mu - \varepsilon)}$ elements γ of Γ with $\log(S(\rho(\gamma)))$ $n\varepsilon$ -close to $n\lambda(\rho_*\mu)$. Which immediately yields the desired lower bound on the value of the growth indicator $\psi_\rho(\lambda(\rho_*\mu))$. $\square$

We now complete the proof of the inequality between the Lyapunov and Faloner dimensions.

Proof of inequality (6). — Fix $\mu \in \mathcal{M}$ and let $r_0 = \dim_{LY}^P(\rho_*\mu) = L_{h_\mu}(\lambda(\rho_*\mu))$. If $r_0 = 0$ there is nothing to prove, so we suppose $r_0 > 0$.

From the duality between Lyapunov and Faloner functionals (Lemma 1.5) we have $F_{r_0}(\lambda(\rho_*\mu)) = h_\mu$. Furthermore, $h_\mu \leq \psi_\rho(\lambda(\rho_*\mu))$ by the fundamental inequality. Hence

$$F_{r_0}^P(\lambda(\rho_*\mu)) \leq \psi_\rho(\lambda(\rho_*\mu)).$$

Because ρ is P -Anosov, $\lambda_i(\rho_*(\mu)) - \lambda_j(\rho_*(\mu)) > 0$ for all $(i, j) \in S(P)$. In particular, for each $0 \leq r < r_0$ we have $F_r^P(\lambda(\rho_*\mu)) < F_{r_0}^P(\lambda(\rho_*\mu))$ whenever $0 \leq r < r_0$ and hence $r \leq \dim_F^P(\rho)$ by Lemma 2.5, which concludes the proof. $\square$

2.3. EIGENSPACE SPLITTING. — In this section, we show that Theorem 1.2 implies Corollary 1.3.

Consider, for $\gamma \in \Gamma$, the P -decomposition $\eta(\rho(\gamma)) = E_1(\rho(\gamma)) \oplus \cdots \oplus E_{M-1}(\rho(\gamma))$. By the P -Anosov property, the angles between the spaces $E_1(\rho(\gamma)) \oplus \cdots \oplus E_k(\rho(\gamma))$ and $E_{k+1}(\rho(\gamma)) \oplus \cdots \oplus E_{M-1}(\rho(\gamma))$ are uniformly bounded from below by some positive number for all $1 \leq k \leq M-2$ and for all γ but a finite number.

Recall that we denote, for all γ but a finite number,

$$\xi(\gamma) = \{0\} \subset \xi^{p_1}(\gamma) \subset \cdots \subset \xi^{p_{M-2}}(\gamma) \subset \mathbb{R}^d \in \mathcal{F}_P.$$

In the same way,

$$\xi(\gamma^{-1}) = \{0\} \subset \xi^{d-p_{M-2}}(\gamma^{-1}) \subset \cdots \subset \xi^{d-p_1}(\gamma^{-1}) \subset \mathbb{R}^d \in \mathcal{F}_{P^*}.$$

It follows from [GGKW17, Lem. 2.26],⁽²⁾ that the unstable flag $f(\rho(\gamma)) \in \mathcal{F}_P$,

$$f(\rho(\gamma)) := \{0\} \subset E_1(\rho(\gamma)) \subset \cdots \subset E_1(\rho(\gamma)) \oplus \cdots \oplus E_{M-2}(\rho(\gamma)) \subset \mathbb{R}^d$$

and the stable flag $f'(\rho(\gamma)) \in \mathcal{F}_{P^*}$,

$$f'(\rho(\gamma)) := \{0\} \subset E_{M-1}(\rho(\gamma)) \subset \cdots \subset E_2(\rho(\gamma)) \oplus \cdots \oplus E_{M-1}(\rho(\gamma)) \subset \mathbb{R}^d$$

are given by

$$f(\rho(\gamma)) = \lim_{n \rightarrow +\infty} \xi(\gamma^n), \quad f'(\rho(\gamma)) = \lim_{n \rightarrow -\infty} \xi(\gamma^n).$$

In particular, $f(\rho(\gamma)) \in \Lambda_\rho \subset \mathcal{F}_P$, and $f'(\rho(\gamma)) \in \Lambda_\rho^*$, where Λ_ρ^* is the limit set associated to the representation ρ in $\mathcal{F}_{P^*}$. Thus, $(f, f') \in (\Lambda_\rho \times \Lambda_\rho^*)' \subset (\mathcal{F}_P \times \mathcal{F}_{P^*})'$, where $'$ indicates that the pairs of flags are in general position.

⁽²⁾The argument in the non-hyperbolic case goes back to David Ruelle's proof of Oseledets theorem ([Rue79], cf. [Led84, Prop. 3.2] or [Sim15, p. 141–142] for details).

and set $A_i = \{p_1, \dots, p_i\}$ for $i = 1, \dots, M$. The desired lower bound will follow from the following two claims:

- (1) For each $i = 1, \dots, M$ one has $a_k(A_i) \geq b_k(p_i)$.
- (2) For each $i = 1, \dots, M - 1$ one has $a_k(A_{i+1}) \geq a_k(A_i) + 1$.

Claim (1) is trivial since $\{p_i\} \subset A_i$. To establish Claim (2) we first suppose that $p_{i+1} \leq k$. If this is the case, then $(p_{i+1}, k+1) \in S(A_{i+1})$. If $(p_{i+1}, k+1) \in S(A_i)$, there would have to be some $j \leq i$ with $p_{i+1} < p_j < k+1$ contradicting the choice of enumeration of P . Hence $a_k(A_{i+1}) \geq a_k(A_i) + 1$ in this case. Similarly, if $p_{i+1} > k$, then $(k, p_{i+1}) \in S(A_{i+1}) \setminus S(A_i)$ and Claim (2) follows. To conclude the proof we observe that from Claim (1) we have $a_k(A_1) - b_k(p_1) \geq 0 = \frac{1 \times 0}{2}$.

Using Claims (1) and (2) we show inductively for $i = 1, \dots, M - 1$ that

$$\begin{aligned} (i+1)a_k(A_{i+1}) - \sum_{j=1}^{i+1} b_k(p_j) &= a_k(A_{i+1}) - b_k(p_{i+1}) + ia_k(A_{i+1}) - \sum_{j=1}^i b_k(p_j) \\ &\geq i + ia_k(A_i) - \sum_{j=1}^i b_k(p_j) \\ &\geq i + \frac{i(i-1)}{2} = \frac{(i+1)i}{2}, \end{aligned}$$

which concludes the proof setting $i = M - 1$. $\square$

4. PROOF OF THEOREM 1.7

The inequality (6), $\sup_{\mu \in \mathcal{M}} \dim_{LY}^P(\rho_*\mu) \leq \dim_F^P(\rho)$, was proved in Section 2.2.2.

4.1. PROOF OF $\dim_F^P(\rho) \leq \sup_{\mu \in \mathcal{M}} \dim_{LY}^P(\rho_*\mu)$. — We recall that the limit cone of ρ is defined as

$$\mathcal{L}_\rho = \lim_{T \rightarrow +\infty} \frac{1}{T} (\log S(\rho(\Gamma))),$$

in the Hausdorff topology. It is a closed convex cone in $\mathfrak{a}^+$. By the P -Anosov property if $a \in \mathcal{L}_\rho \setminus \{0\}$ we obtain

$$\alpha_{i,j}(a) > 0 \text{ for all } (i, j) \in S(P).$$

Assume that $r < r' < \dim_F^P(\rho)$ so that the series (4) diverges at r' . By compactness there exists $a \in \mathcal{L}_\rho \setminus \{0\}$ such that for all open cones $\mathcal{C}$ containing a we have

$$\sum_{\gamma \in \Gamma: \log S(\rho(\gamma)) \in \mathcal{C}} \exp(-F_{r'}^P \circ \log S(\rho(\gamma))) = +\infty.$$

We fix such a choice of a and notice that $\psi(a) \geq F_{r'}^P(a)$ by [Qui02, Lem. III.1.3] and [Sam14, Lem. 4.2]. Thus, by Lemma 1.5, $\psi(a) > F_r^P(a)$. We will prove:

PROPOSITION 4.1. — *There exists a sequence $T_k \rightarrow +\infty$ and $\mu_k \in \mathcal{M}$ such that $h_{\mu_k} = T_k \psi(a) + o(T_k)$ and $\lambda(\rho_*\mu_k) = T_k a + o(T_k)$ when $k \rightarrow +\infty$.*

Since $\psi(a) > F_r^P(a)$, we have from Lemma 1.5 that $L_{\psi(a)}^P(a) > r$. Assuming Proposition 4.1, we obtain

$$\begin{aligned} \dim_{LY}^P(\rho_*\mu_k) &= L_{h_{\mu_k}}^P(\rho_*\mu_k) = L_{T_k\psi(a)+o(T_k)}^P(T_ka + o(T_k)) \\ &= L_{\psi(a)+o(1)}^P(a + o(1)) = L_{\psi(a)}^P(a) + o(1) > r, \end{aligned}$$

for k large enough. Which concludes the proof of the inequality and of Theorem 1.7. $\square$

4.2. PROOF OF PROPOSITION 4.1. — Fix $\beta : \mathfrak{a} \rightarrow \mathbb{R}$ linear such that $\beta(a) = F_r^P(a)$, and a decreasing sequence of open cones $\mathcal{C}_k$, $k = 1, 2, 3, \dots$ whose intersection is $\mathbb{R}_+a$.

For a given k set

$$\psi_k(a) := \limsup_{T \rightarrow +\infty} \frac{1}{T} \log \# \{ \gamma \in \Gamma : \log S(\rho(\gamma)) \in \mathcal{C}_k \cap \{ \beta \leq T\beta(a) \} \}.$$

We have $\psi(a) = \lim_{k \rightarrow \infty} \psi_k(a)$ and

$$\psi_k(a) = \limsup_{T \rightarrow +\infty} \frac{1}{T} \log \# \{ \gamma \in \Gamma : \log S(\rho(\gamma)) \in \mathcal{C}_k \cap \{ (T-1)\beta(a) \leq \beta \leq T\beta(a) \} \}.$$

Hence, we may choose $T_k \rightarrow +\infty$ such that the sets

$$A_k = \{ \gamma \in \Gamma : \log S(\rho(\gamma)) \in \mathcal{C}_k \cap \{ (T_k-1)\beta(a) \leq \beta \leq T_k\beta(a) \} \},$$

satisfy

$$\psi(a) = \lim_{k \rightarrow +\infty} \frac{1}{T_k} \log \# A_k.$$

Moreover, by the Anosov property, there are constants C_1, C_2 such that for $\gamma \in A_k$, $C_1 T_k \leq |\gamma| \leq C_2 T_k$. So there exist $\ell_k \rightarrow \infty$ as $k \rightarrow \infty$, such that the sets

$$A'_k = \{ \gamma \in \Gamma : \log S(\rho(\gamma)) \in \mathcal{C}_k \cap \{ (T_k-1)\beta(a) \leq \beta \leq T_k\beta(a) \}, |\gamma| = \ell_k \},$$

still satisfy

$$\psi(a) = \lim_{k \rightarrow +\infty} \frac{1}{T_k} \log \# A'_k.$$

We now fix a semi-group S of Γ , a finite subset F , and maps $L, R : \Gamma \rightarrow F$ given by Proposition A.1. Observe that $\rho(S)$ is Zariski dense since the Zariski closure of a semi-group is a group (see [GM89]) and since S generates Γ .

Let $B_k \subset S$ be defined by

$$B_k = \{ L(\gamma)\gamma R(\gamma) : \gamma \in A'_k \}.$$

Since F is finite, there exists $C > 0$ such that

$$\#B_k \geq C^{-1} \#A'_k, \quad ||\gamma| - \ell_k| \leq C \text{ for } \gamma \in B_k$$

and fixing some norm on $\mathfrak{a}$ we have

$$\| \log S(\rho(\gamma)) - \log S(\rho(L(\gamma)\gamma R(\gamma))) \| \leq C,$$

for all $\gamma \in \Gamma$.

Fix a parameter $\varepsilon > 0$, we say a splitting $\mathbb{R}^d = E_1 \oplus \dots \oplus E_d$ into one dimensional subspaces is ε -non-degenerate if the angle between E_i and $\bigoplus_{j \neq i} E_j$ is at least ε for all i . We say two splittings $(E_1, \dots, E_d), (E'_1, \dots, E'_d)$ are ε -close if the angle between E_i and E'_i is at most ε for all i .

We say that an element $g \in \mathrm{SL}(d, \mathbb{R})$ is ε -diagonalizable if it has d distinct real eigenvalues $|\lambda_1| > \cdots > |\lambda_d|$ satisfying $|\lambda_i|/|\lambda_{i+1}| > e^\varepsilon$ for $i = 1, \dots, d-1$, and the corresponding splitting into eigenspaces is ε -non-degenerate.

Since $\rho(S)$ is Zariski dense we obtain from [AMS95, Th. 6.8]:

LEMMA 4.2 (Abels-Margulis-Soifer). — *There exists $\varepsilon_0 > 0$ and a finite set $S_0 \subset S$ such that for all $\gamma \in \Gamma$ the set $\rho(\gamma S_0)$ contains at least one ε_0 -diagonalizable element.*

For what follows we fix S_0 and $\varepsilon_0 > 0$ given by the previous lemma. We also fix $\delta_0 > 0$ given by the following:

LEMMA 4.3. — *There exists $\delta_0 > 0$ such that if B is a set of ε_0 -diagonalizable elements of $\mathrm{SL}(d, \mathbb{R})$ whose eigenspace splittings are δ_0 -close, then $\gamma_1 \cdots \gamma_n$ is $\varepsilon_0/2$ -diagonalizable for all n and $\gamma_1, \dots, \gamma_n \in B$.*

Proof. — This follows directly from [BS21, Lem. 2.17]. $\square$

We now refine the family B_k so as to obtain images under ρ which are diagonalizable with nearby splittings.

LEMMA 4.4. — *There exists a decreasing positive sequence $\varepsilon_k \downarrow 0$ and a sequence $\ell'_k \uparrow \infty$ such that for all k large there exists $C_k \subset B_k \cdot S_0$ with the following properties:*

- (1) C_k freely generates a free semi-group in Γ .
- (2) For all $\gamma \in C_k$ the element $\rho(\gamma)$ is ε_0 -diagonalizable.
- (3) For all $\gamma_1, \gamma_2 \in C_k$ the eigenspace splittings of $\rho(\gamma_1)$ and $\rho(\gamma_2)$ are δ_0 -close.
- (4) One has $\psi(a) = \lim_{k \rightarrow +\infty} \frac{1}{T_k} \log \#C_k$.
- (5) $|\gamma| = \ell'_k$ for $\gamma \in C_k$.
- (6) For all $\gamma \in C_k$ one has $\|(\log S(\rho(\gamma))) - T_k a\| \leq \varepsilon_k T_k$.

Proof. — The space of ε_0 -non-degenerate splittings is compact and hence we may cover it by some finite number N of subsets with the property that if two splittings belong to the same subset they are δ_0 -close.

From Lemma 4.2 it follows that for all $\gamma \in B_k$ there exists $f_\gamma \in S_0$ such that $\rho(\gamma f_\gamma)$ is ε_0 -diagonalizable. By the pigeonhole principle we may choose a set C'_k of at least $\#B_k/N$ of the elements γf_γ such that the eigenspace splitting of ρ applied to any two are δ_0 -close. Hence, we have constructed C'_k with properties (2), (3) and (4) above.

Since $F \cup S_0$ is finite there exists $C > 0$ such that, for all k and $\gamma \in A_k$,

$$\begin{aligned} \|L(\gamma)\gamma R(\gamma)f_{L(\gamma)\gamma R(\gamma)} - \ell_k\| &\leq C \quad \text{and} \\ \|(\log S(\rho(L(\gamma)\gamma R(\gamma)f_{L(\gamma)\gamma R(\gamma)})) - (\log S(\rho(\gamma)))\| &\leq C. \end{aligned}$$

By the pigeonhole principle again, we can find $\ell'_k, |\ell'_k - \ell_k| \leq C$, such that $C''_k := \gamma \in C'_k, |\gamma| = \ell'_k$ satisfies properties (2), (3), (4) and (5). Moreover, if $\gamma \in A_k$ then by definition we have

$$\left(\frac{1}{T_k}(\log S(\rho(\gamma))) - a\right) \in \mathcal{C}_k \cap \left\{\frac{T_k - 1}{T_k}\beta(a) \leq \beta \leq \beta(a)\right\},$$

hence setting ε_k to be C/T_k plus the diameter of the set on the right-hand side, we have

$$\|(\log S(\rho(L(\gamma)\gamma R(\gamma)f_\gamma)) - T_k a)\| \leq \varepsilon_k T_k,$$

for all $\gamma \in A_k$, which establishes property (6) for C'_k and C''_k .

To establish (1) we notice that S is a quasi-geodesic semi-group. Therefore, by the Morse lemma, there exists some $R > 0$ such that if $\gamma_1, \dots, \gamma_n, \eta_1, \dots, \eta_n \in S$ and $\gamma_1 \cdots \gamma_n = \eta_1 \cdots \eta_n$, then there is a geodesic ray $\alpha = \{\alpha_n\}_{n \geq 0}$, such that for all $j \in [0, n]$,

$$(11) \quad \text{dist}(\gamma_1 \cdots \gamma_j, \alpha), \text{dist}(\eta_1 \cdots \eta_j, \alpha) < R.$$

We claim that if we assume that the elements of $C_k \subset C''_k$ are $6R$ -separated, equation (11) forces $\gamma_j = \eta_j$ for all $j \in [0, n]$. Hence it suffices to refine C''_k so that it is a $6R$ -separated set to obtain that it freely generates a free sub-semi-group of Γ . Doing this decreases the number of elements at most by a factor equal to the number of elements in a ball of radius $6R$ in Γ , so property (4) still holds.

To prove the claim, we prove by induction on j that $\gamma_i = \eta_i$ for $i \leq j$. We have $\gamma_0 = \eta_0 = e$. Assume that $\gamma_i = \eta_i$ for $i \leq j$. Then $\gamma_1 \cdots \gamma_j = \eta_1 \cdots \eta_j$ and there exist N_1, N_2 and N_3 such that

$$\text{dist}(\gamma_1 \cdots \gamma_j, \alpha_{N_1}), \text{dist}(\gamma_1 \cdots \gamma_j \gamma_{j+1}, \alpha_{N_2}) \text{ and } \text{dist}(\gamma_1 \cdots \gamma_j \eta_{j+1}, \alpha_{N_3}) \leq R.$$

Setting $\beta_t := (\gamma_1 \cdots \gamma_j)^{-1} \alpha_{N_t}$ for $t = 1, 2, 3$, we have β_{N_1}, β_{N_2} and β_{N_3} aligned on the same geodesic and satisfying:

$$|\beta_1|, \text{dist}(\gamma_{j+1}, \beta_2), \text{dist}(\eta_{j+1}, \beta_3) < R \quad \text{and} \quad |\gamma_{j+1}| = |\eta_{j+1}| = \ell'_k.$$

If ℓ'_k is large enough, β_1 is outside the interval $[\beta_2, \beta_3]$ (see Lemma A.7) and

$$\text{dist}(\beta_1, \beta_2), \text{dist}(\beta_1, \beta_3) \in [\ell'_k - 2R, \ell'_k + 2R].$$

It follows that $\text{dist}(\beta_2, \beta_3) \leq 4R$ and therefore that $\text{dist}(\alpha_{j+1}, \eta_{j+1}) < 6R$. Since C_k is $6R$ -separated, $\alpha_{j+1} = \eta_{j+1}$ as claimed. $\square$

COROLLARY 4.5. — *If μ_k is the uniform probability measure on C_k then one has*

$$\lim_{k \rightarrow +\infty} \frac{1}{T_k} h_{\mu_k} = \psi(a).$$

Proof. — This follows immediately from properties (1) and (4) of the previous lemma. $\square$

We now use the fact that for ε_0 -diagonalizable elements with close splittings the singular values and eigenvalues are close and they almost multiply under composition.

LEMMA 4.6. — *There exists a decreasing positive sequence $\varepsilon'_k \downarrow 0$ such that for all k large enough, all n and $\gamma_1, \dots, \gamma_n \in C_k$ one has*

$$\left\| \frac{1}{n} (\log S(\rho(\gamma_1 \cdots \gamma_n))) - T_k a \right\| \leq \varepsilon'_k T_k.$$

Proof. — Let $L(g)$ denote the vector of absolute values of eigenvalues of a diagonalizable element $g \in \mathrm{SL}(d, \mathbb{R})$ in decreasing order. By [BS21, Lem. 2.16] there exists $C > 0$ such that for all $\varepsilon_0/2$ -diagonalizable $g \in \mathrm{SL}(d, \mathbb{R})$ one has

$$\|\log S(g) - \log L(g)\| \leq C.$$

By [BS21, Prop. 2.17] there exists $C > 0$ independent of k and n such that we have

$$\|(\log L(\rho(\gamma_1 \cdots \gamma_n)) - \sum_{i=1}^n (\log L(\rho(\gamma_i)))\| \leq Cn.$$

Combining this with Lemma 4.4 we obtain

$$\begin{aligned} \|(\log S(\rho(\gamma_1 \cdots \gamma_n)) - nT_k a)\| &\leq C + \|(\log L(\rho(\gamma_1 \cdots \gamma_n)) - nT_k a)\| \\ &\leq C(n+1) + \|\sum_{i=1}^n (\log L(\rho(\gamma_i)) - nT_k a)\| \\ &\leq C(2n+1) + \|\sum_{i=1}^n (\log S(\rho(\gamma_i)) - nT_k a)\| \\ &\leq C(2n+1) + n\varepsilon_k T_k. \end{aligned}$$

This proves the desired result with $\varepsilon'_k = 3C/T_k + \varepsilon_k$. $\square$

The following immediate corollary concludes the proof of Proposition 4.1.

COROLLARY 4.7. — *For any sequence of probability measures μ_k with $\mu_k(C_k) = 1$ one has*

$$\lim_{k \rightarrow +\infty} \frac{1}{T_k} \lambda(\rho_* \mu_k) = a.$$

5. COVERING BY BALLS AND PROOF OF THEOREM 1.8

Fix $\eta > 0$ and set $s := \dim_F^P(\rho) + \eta$. We may assume $s < \dim(\mathcal{F}_P)$, otherwise $\dim_B(\Lambda_\rho) \leq \dim_F^P(\rho) + \eta$ holds trivially. Since $s > \dim_F^P(\rho)$, the series $\Phi_\rho^P(s) := \sum_{\gamma \in \Gamma} \varphi_s^P(\rho(\gamma))$ converges. We are going to construct covers $\mathcal{U}$ of Λ_ρ by balls of arbitrarily small radius ε with less than $\Phi_\rho^P(s)\varepsilon^{-s+o(\varepsilon)}$ elements. This shows that the ratio $N(\Lambda_\rho, \varepsilon)/\varepsilon^{-s+o(\varepsilon)}$ is bounded from above by $\Phi_\rho^P(s)$ uniformly in ε . Therefore, $\dim_B(\Lambda_\rho) \leq s = \dim_F^P(\rho) + \eta$ for all positive η and Theorem 1.8 follows.

5.1. SHADOWS AND ANOSOV REPRESENTATIONS. — Recall that Γ is a hyperbolic group and that we chose the generating set $\mathcal{S}$ to be symmetric. The distance $d(\gamma, \gamma')$ on Γ is given by the word length of $\gamma^{-1}\gamma'$. A *geodesic* $\sigma = \{\gamma_n\}_{n \in \mathbb{Z}}$ in Γ is a sequence such that for all (i, j) , $d(\gamma_i, \gamma_j) = |j - i|$. Any point $x \in \partial\Gamma$ is the limit point of (at least) one geodesic ray $\sigma = \{\gamma_j\}_{j \geq 0}$ with $\gamma_0 = e$.

For a geodesic ray $\sigma = \{\gamma_j\}_{j \geq 0}$ with $\gamma_0 = e$, we call the R -shadow of the geodesic ray the image by ξ of the set of limit points of geodesic rays $\sigma' = \{\gamma'_j\}_{j \geq 0}$ satisfying $\gamma'_j = \gamma_j$ for $j \leq R$. By definition, the R -shadow of a geodesic ray is a subset of $\mathcal{F}_P$.

For $(i, j) \in S(P)$, $\gamma \in \Gamma$, write $\zeta_{i,j}(\gamma) := \log(s_i(\rho(\gamma))/s_j(\rho(\gamma)))$. The main step for the proof of Theorem 1.8 is the following proposition:

APPENDIX. QUASI-GEODESIC SEMI-GROUPS

A.1. GOAL. — If Γ is the free group with generators a, b then the semi-group S consisting of reduced words which begin and end with a has the following nice properties:

- (1) S generates Γ as a group.
- (2) For every n and $\gamma_1, \dots, \gamma_n \in S$ the geodesic joining the neutral element e to $\gamma_1 \cdots \gamma_n$ passes through all partial products $\gamma_1 \cdots \gamma_k$ for $k = 1, \dots, n$.
- (3) For every $\gamma \in \Gamma$ there exist two letter words $L(\gamma)$ and $R(\gamma)$ such that $L(\gamma)\gamma R(\gamma) \in S$.

We may for instance set $L(\gamma)$ to be a if the reduced form of γ does not begin with a^{-1} and ab otherwise, and $R(\gamma)$ similarly to be a if the last letter of γ is not a^{-1} and ba otherwise.

Our goal in this appendix is to give a similar construction for a general hyperbolic group. For this purpose we fix a group Γ with a finite symmetric generating set $\mathcal{S}$. We let $|\gamma|$ be the word length of $\gamma \in \Gamma$ with respect to $\mathcal{S}$ and consider the word distance $\text{dist}(\gamma, \eta) = |\gamma^{-1}\eta|$. We assume that there exists $\delta > 0$ (fixed from now on) such that Γ is δ -hyperbolic with respect to dist , and non-elementary (i.e., contains two independent loxodromic elements).

Recall that given positive constants $C, D > 0$ a (C, D) -quasi-geodesic is a sequence $x : I \cap \mathbb{Z} \rightarrow \Gamma$ where I is an interval, satisfying

$$C^{-1}|m - n| - D \leq \text{dist}(x_m, x_n) \leq C|m - n| + D,$$

for all $m, n \in I \cap \mathbb{Z}$.

We say a semigroup $S \subset \Gamma$ is quasi-geodesic if there exist constants $C, D > 0$ such that for all n and $\gamma_1, \dots, \gamma_n \in S$ the sequence

$$e, \gamma_1, \gamma_1\gamma_2, \dots, \gamma_1 \cdots \gamma_n,$$

is visited in left to right order by some (C, D) -quasi-geodesic.

We now state the main result of the appendix.

PROPOSITION A.1. — *Let $(\Gamma, \mathcal{S})$ be a non-elementary hyperbolic group with a finite symmetric generating set $\mathcal{S}$. Then there exists a semigroup $S \subset \Gamma$, a finite set $F \subset \Gamma$ and mappings $L, R : \Gamma \rightarrow F$ with the following properties:*

- (1) S generates Γ as a group,
- (2) S is quasi-geodesic, and
- (3) $L(\gamma)\gamma R(\gamma) \in S$ for all $\gamma \in \Gamma$.

The arguments in what follows use the local-to-global principle for hyperbolic groups as in [Gou22, §3].

A.2. PROOF OF PROPOSITION A.1. — Recall that the Gromov product (based at the neutral element $e \in \Gamma$) is defined by

$$(\gamma, \eta) = \frac{|\gamma| + |\eta| - |\gamma^{-1}\eta|}{2}.$$

We assume $\delta > 0$ is such that

$$\min\{(\gamma_1, \gamma_2), (\gamma_2, \gamma_3)\} \leq (\gamma_1, \gamma_3) + \delta,$$

for all $\gamma_1, \gamma_2, \gamma_3 \in \Gamma$. We further assume that

$$\text{dist}(e, \alpha) \leq (p, q) + \delta,$$

for all $p, q \in \Gamma$ and geodesic α joining them (such a choice of δ is possible by [BH99, Prop. 1.22]).

LEMMA A.2. — *For each $C \geq 1$ there exist $a, b \in \Gamma$ such that setting $A = \{a, b, a^{-1}, b^{-1}\}$ and defining*

$$c_1 = \max_{x, y \in A: x \neq y} (x, y) \quad \text{and} \quad c_2 = \min_{x \in A} |x|,$$

one has $c_1 + 2\delta + r < \frac{1}{2}c_2$ for some $r > C(c_1 + 2\delta + 1)$.

Proof. — Since Γ is non-elementary we may take x, y two independent loxodromic elements (i.e., the set of boundary fixed points of x and y are disjoint). Letting $a = x^n$ and $b = y^n$ for sufficiently large n establishes the claim since $c_1(x^n, y^n)$ remains bounded, while $c_2(x^n, y^n)$ goes to infinity. $\square$

We fix from now on a set A as in the previous lemma with corresponding constants c_1, c_2, r satisfying

$$(18) \quad r > 32(c_1 + 2\delta + 1).$$

LEMMA A.3. — *The sets defined for $x \in A$ by $V(x) = \{\gamma : (\gamma, x) \geq c_1 + 2\delta + r\}$ are pairwise disjoint.*

- For all $x \in A$ and $\gamma \notin V(x^{-1})$ one has $x\gamma \in V(x)$.
- For all $x, y \in A$ with $x \neq y$, if $\gamma \in V(x)$ and $\eta \in V(y)$ then $(\gamma, \eta) \leq c_1 + 2\delta$.
- If $\gamma \in V(x)$ for some $x \in A$ then $|\gamma| \geq r$.

Proof. — By hypothesis $2\delta + r \in (2\delta, c_2/2 - c_1)$. With this property, the first three statements are claims 1–3 in the proof of [AGG⁺23, Lem. 4.1].

For the last statement observe that

$$|\gamma| = (\gamma, \gamma) \geq \min\{(\gamma, x), (x, x)\} - \delta \geq \min\{c_1 + 2\delta + r, c_2\} - \delta \geq r. \quad \square$$

LEMMA A.4. — *There exists m such that for all $\gamma \in V(a)$ and all η with $\text{dist}(\gamma, \eta) \leq 2c_2$ one has $a^m \eta \in V(a)$.*

Proof. — We first observe that by Lemma A.3 we have $(a^{-1}, a) \leq c_1 + 2\delta$ and therefore a geodesic fixed under multiplication by a is at distance at most $c_1 + 3\delta$ from a^m for all m . It follows, letting $\ell = \lim_{n \rightarrow +\infty} \frac{1}{n}|a^n|$ be the translation length of a , that

$$\ell \geq |a| - 2(c_1 + 3\delta) \geq 2(c_1 + 2\delta + r) - 2(c_1 + 3\delta) \geq 2r - 2\delta.$$

Hence we obtain for all $m \geq 1$ that

$$\begin{aligned} (a, a^m) &= \frac{|a| + |a^m| - |a^{m-1}|}{2} \\ &\geq \frac{\ell - 2(c_1 + 3\delta) + m\ell - 2(c_1 + 3\delta) - (m-1)\ell - 2(c_1 + 3\delta)}{2} \\ &= \ell - 3(c_1 + 3\delta) \geq 2r - 3c_1 - 11\delta \geq c_1 + 3\delta + r. \end{aligned}$$

In what follows we will need to switch basepoints, so we recall that the Gromov product

$$(x, y)_z = \frac{\text{dist}(x, z) + \text{dist}(y, z) - \text{dist}(x, y)}{2},$$

satisfies $(x, y)_z + (z, y)_x = \text{dist}(x, z)$. We now calculate using Lemma A.3

$$\begin{aligned} (a^m, a^m\eta) &= (e, \eta)_{a^{-m}} \\ &= |a^m| - (a^{-m}, \eta)_e \\ &\geq |a^m| - (a^{-m}, \gamma)_e - 2c_2 \geq |a^m| - c_1 - 2\delta - 2c_2 \geq c_1 + 3\delta + r, \end{aligned}$$

if we choose m large enough depending only on c_1, c_2 and δ .

We conclude the proof using the hyperbolicity property since

$$(a, a^m\eta) \geq \min\{(a, a^m), (a^m, a^m\eta)\} - \delta \geq c_1 + 2\delta + r,$$

so $a^m\eta \in V(a)$ as required. $\square$

We define

$$T = \{\gamma : \gamma \in V(a) \text{ and } \gamma^{-1} \in V(a^{-1})\}.$$

With m given by the previous lemma, for each $\gamma \in \Gamma$ let

$$L(\gamma) = \begin{cases} a^{m+1} & \text{if } \gamma \notin V(a^{-1}), \\ a^{m+1}b & \text{otherwise,} \end{cases} \quad \text{and} \quad R(\gamma) = \begin{cases} a & \text{if } (L(\gamma)\gamma)^{-1} \notin V(a), \\ ba & \text{otherwise.} \end{cases}$$

LEMMA A.5. — One has $L(\gamma)\gamma R(\gamma) \in T$ for all $\gamma \in \Gamma$.

Proof. — Since $\text{dist}(L(\gamma)\gamma, L(\gamma)\gamma R(\gamma)) \leq 2c_2$ this follows immediately from Lemma A.4. $\square$

We now let S be the semigroup generated by T .

LEMMA A.6. — The semigroup S generates Γ as a group.

Proof. — From the definition $a \in S$ and, using Lemma A.3, $aba \in S$. Therefore $L(\gamma), R(\gamma)$ belong to the group generated by S for all γ . The equation

$$\gamma = L(\gamma)^{-1} (L(\gamma)\gamma R(\gamma)) R(\gamma)^{-1}$$

now implies that S generates Γ as a group, as required. $\square$

As a first step towards showing that S is quasi-geodesic we prove the following.

LEMMA A.7. — For any n and $\gamma_1, \dots, \gamma_n \in T$, letting α be a geodesic joining e and $\gamma_1 \cdots \gamma_n$, one has

$$|\gamma_1 \cdots \gamma_n| \geq |\gamma_1 \cdots \gamma_k| + |\gamma_{k+1} \cdots \gamma_n| - r/8,$$

and $\text{dist}(\gamma_1 \cdots \gamma_k, \alpha) \leq r/8$ for all $k = 0, \dots, n$.

Proof. — Setting $x_k = \gamma_1 \cdots \gamma_k$ for $k = 0, \dots, n$, we observe that

$$(x_{k-1}, x_{k+1})_{x_k} = (\gamma_k^{-1}, \gamma_{k+1}) \leq c_1 + 2\delta,$$

and $\text{dist}(x_k, x_{k+1}) = |\gamma_{k+1}| \geq r$ by Lemma A.3. Hence the sequence is a $(c_1 + \delta, r)$ -chain as in [Gou22]. We observe that

$$r > 2(c_1 + 2\delta) + 4\delta + 1,$$

and therefore by [Gou22, Lem. 3.8] we have

$$(x_0, x_n)_{x_k} \leq c_1 + 4\delta,$$

for all k . This implies that

$$\text{dist}(x_k, \alpha) \leq c_1 + 5\delta < r/8.$$

Also since $(x_0, x_n)_{x_k} = (x_k^{-1}, x_k^{-1}x_n)_e$ we get by definition

$$\frac{|\gamma_1 \cdots \gamma_k| + |\gamma_{k+1} \cdots \gamma_n| - |\gamma_1 \cdots \gamma_n|}{2} \leq c_1 + 4\delta \leq r/16,$$

which yields the required lower bound for $|\gamma_1 \cdots \gamma_n|$. $\square$

We now conclude the proof of Proposition A.1.

LEMMA A.8. — The semigroup S is quasi-geodesic.

Proof. — Fix n and $\gamma_1, \dots, \gamma_n \in T$, and let α be a geodesic with $\alpha_0 = e$ and $\alpha_{N_n} = \gamma_1 \cdots \gamma_n$ with $N_n = |\gamma_1 \cdots \gamma_n|$. Since the word distance is integer valued, setting $C = \lfloor r/8 \rfloor$ we have by Lemma A.7 that there exist $N_0 = 0, N_1, \dots, N_n = |\gamma_1 \cdots \gamma_n|$ such that

$$\text{dist}(\gamma_1 \cdots \gamma_k, \alpha_{N_k}) \leq C \leq r/8,$$

for all k .

We also obtain from Lemma A.7 applied to $\gamma_1, \dots, \gamma_{k+1}$ and Lemma A.3 that

$$\begin{aligned} N_{k+1} - N_k &\geq |\gamma_1 \cdots \gamma_{k+1}| - r/8 - |\gamma_1 \cdots \gamma_k| - r/8 \\ &\geq |\gamma_{k+1}| - 3r/8 \geq 5r/8 \geq 5C, \end{aligned}$$

and in particular $N_k \leq N_{k+1}$ for $k = 0, \dots, n-1$.

We now construct a path β which we claim to be quasi-geodesic (with constants independent of n and $\gamma_1, \dots, \gamma_n$) by concatenating segments of α with a paths to and from each $\gamma_1 \cdots \gamma_k$ (which will have length at most $2C$ as seen above). To be precise, we choose $\beta : [0, N_n + 2(n-1)C] \cap \mathbb{Z} \rightarrow \Gamma$ such that

- (1) For each $k = 0, \dots, n-1$ one has $\beta_{m+N_k+2kC} = \alpha_{m+N_k}$ for all $m \in [0, N_{k+1} - N_k] \cap \mathbb{Z}$.
- (2) $\beta_{N_k+2(k-1)C+C} = \gamma_1 \cdots \gamma_k$ for $k = 1, \dots, n-1$.
- (3) $\text{dist}(\beta_m, \beta_{m+1}) \leq 1$ for all m .

In view of the third property above we have $\text{dist}(\beta_\ell, \beta_m) \leq |\ell - m|$. For the lower bound, given $0 \leq \ell < m \leq N_n + 2(n-1)C$ we write

$$\text{dist}(\beta_\ell, \beta_m) = \sum_{k=0}^{m-\ell} \text{dist}(\beta_{\ell+k}, \beta_{\ell+k+1}) \geq \sum_{k=0}^{m-\ell} f(k),$$

where we set $f(k) = 1$ if $k \in [N_i + 2iC, N_{i+1} + 2iC]$ for some i and $f(k) = -1$ otherwise. Since the sequence $f(0), \dots, f(\ell - m)$ consists of subsequences of runs of at most $2C$ consecutive times the value -1 , with runs of at least $5C$ times the value 1 in between, we obtain

$$\text{dist}(\beta_\ell, \beta_m) \geq 3(|m - \ell| - 4C)/7 - 4C,$$

which establishes that S is $(7/3, 6C)$ -quasi-geodesic. $\square$

REFERENCES

- [AMS95] H. ABELS, G. A. MARGULIS & G. A. SOÏFER – “Semigroups containing proximal linear maps”, *Israel J. Math.* **91** (1995), no. 1-3, p. 1–30.
- [AGG⁺23] A. AZEMAR, V. GADRE, S. GOUËZEL, T. HAETTEL, P. LESSA & C. UYANIK – “Random walk speed is a proper function on Teichmüller space”, *J. Modern Dyn.* **19** (2023), p. 815–832.
- [BCH10] J. BAN, Y. CAO & H. HU – “The dimensions of a non-conformal repeller and an average conformal repeller”, *Trans. Amer. Math. Soc.* **362** (2010), no. 2, p. 727–751.
- [BHR19] B. BÁRÁNY, M. HOCHMAN & A. RAPAPORT – “Hausdorff dimension of planar self-affine sets and measures”, *Invent. Math.* **216** (2019), no. 3, p. 601–659.
- [Bar10] T. BARBOT – “Three-dimensional Anosov flag manifolds”, *Geom. Topol.* **14** (2010), no. 1, p. 153–191.
- [Ben96] Y. BENOIST – “Actions propres sur les espaces homogènes réductifs”, *Ann. of Math. (2)* **144** (1996), no. 2, p. 315–347.
- [BPS19] J. BOCHI, R. POTRIE & A. SAMBARINO – “Anosov representations and dominated splittings”, *J. Eur. Math. Soc. (JEMS)* **21** (2019), no. 11, p. 3343–3414.
- [BS21] E. BREUILLARD & Ç. SERT – “The joint spectrum”, *J. London Math. Soc. (2)* **103** (2021), no. 3, p. 943–990.
- [BV19] E. BREUILLARD & P. P. VARJÚ – “On the dimension of Bernoulli convolutions”, *Ann. Probab.* **47** (2019), no. 4, p. 2582–2617.
- [BH99] M. R. BRIDSON & A. HAEFLIGER – *Metric spaces of non-positive curvature*, Grundlehren Math. Wissen., vol. 319, Springer-Verlag, Berlin, 1999.
- [Can] R. D. CANARY – “Anosov representations: informal lecture notes”, <https://dept.math.lsa.umich.edu/~canary/lecnotespublic.pdf>.
- [Dey22] S. DEY – “On Borel Anosov subgroups of $\text{SL}(d, \mathbb{R})$ ”, 2022, to appear in *Geom. Topol.*, [arXiv:2208.02109](https://arxiv.org/abs/2208.02109).
- [DK22] S. DEY & M. KAPOVICH – “Patterson-Sullivan theory for Anosov subgroups”, *Trans. Amer. Math. Soc.* **375** (2022), no. 12, p. 8687–8737.
- [DO80] A. DOUADY & J. OESTERLÉ – “Dimension de Hausdorff des attracteurs”, *C. R. Acad. Sci. Paris Sér. A-B* **290** (1980), no. 24, p. A1135–A1138.
- [Doy88] P. G. DOYLE – “On the bass note of a Schottky group”, *Acta Math.* **160** (1988), no. 3-4, p. 249–284.
- [Duf17] L. DUFLUX – “Hausdorff dimension of limit sets”, *Geom. Dedicata* **191** (2017), p. 1–35.
- [DS22] L. DUFLUX & V. SUOMALA – “Projections of Poisson cut-outs in the Heisenberg group and the visual 3-sphere”, *Math. Proc. Cambridge Philos. Soc.* **172** (2022), no. 1, p. 197–230.
- [Fal88] K. J. FALCONER – “The Hausdorff dimension of self-affine fractals”, *Math. Proc. Cambridge Philos. Soc.* **103** (1988), no. 2, p. 339–350.
- [FS22] D.-J. FENG & K. SIMON – “Dimension estimates for C^1 iterated function systems and repellers. Part II”, *Ergodic Theory Dynam. Systems* **42** (2022), no. 11, p. 3357–3392.

- [FS23] ———, “Dimension estimates for C^1 iterated function systems and repellers. Part I”, *Ergodic Theory Dynam. Systems* **43** (2023), no. 8, p. 2673–2706.
- [GdlH90] É. GHYS & P. DE LA HARPE (eds.) – *Sur les groupes hyperboliques d’après Mikhael Gromov*, Progress in Math., vol. 83, Birkhäuser Boston, Inc., Boston, MA, 1990.
- [GMT23] O. GLORIEUX, D. MONCLAIR & N. THOLOZAN – “Hausdorff dimension of limit sets for projective Anosov representations”, *J. Éc. polytech. Math.* **10** (2023), p. 1157–1193.
- [GM89] I. Y. GOL’DSHEĬD & G. A. MARGULIS – “Lyapunov exponents of a product of random matrices”, *Uspehi Mat. Nauk* **44** (1989), no. 5(269), p. 13–60.
- [Gou22] S. GOUËZEL – “Exponential bounds for random walks on hyperbolic spaces without moment conditions”, *Tunis. J. Math.* **4** (2022), no. 4, p. 635–671.
- [GGKW17] F. GUÉRITAUD, O. GUICHARD, F. KASSEL & A. WIENHARD – “Anosov representations and proper actions”, *Geom. Topol.* **21** (2017), no. 1, p. 485–584.
- [Hoc14] M. HOCHMAN – “On self-similar sets with overlaps and inverse theorems for entropy”, *Ann. of Math. (2)* **180** (2014), no. 2, p. 773–822.
- [HR22] M. HOCHMAN & A. RAPAPORT – “Hausdorff dimension of planar self-affine sets and measures with overlaps”, *J. Eur. Math. Soc. (JEMS)* **24** (2022), no. 7, p. 2361–2441.
- [HS17] M. HOCHMAN & B. SOLOMYAK – “On the dimension of Furstenberg measure for $\mathrm{SL}_2(\mathbb{R})$ random matrix products”, *Invent. Math.* **210** (2017), no. 3, p. 815–875.
- [JLPX24] Y. JIAO, J. LI, W. PAN & D. XU – “On the dimension of limit sets on $\mathbb{P}(\mathbb{R}^3)$ via stationary measures: variational principles and applications”, *Internat. Math. Res. Notices* (2024), no. 19, p. 13015–13045.
- [JPS07] T. JORDAN, M. POLLICOTT & K. SIMON – “Hausdorff dimension for randomly perturbed self affine attractors”, *Comm. Math. Phys.* **270** (2007), no. 2, p. 519–544.
- [Käe04] A. KÄENMÄKI – “On natural invariant measures on generalised iterated function systems”, *Ann. Acad. Sci. Fenn. Math.* **29** (2004), no. 2, p. 419–458.
- [Kai00] V. A. KAIMANOVICH – “The Poisson formula for groups with hyperbolic properties”, *Ann. of Math. (2)* **152** (2000), no. 3, p. 659–692.
- [KV83] V. A. KAIMANOVICH & A. M. VERSHIK – “Random walks on discrete groups: boundary and entropy”, *Ann. Probab.* **11** (1983), no. 3, p. 457–490.
- [KY79] J. L. KAPLAN & J. A. YORKE – “Chaotic behavior of multidimensional difference equations”, in *Functional differential equations and approximation of fixed points (Bonn, 1978)*, Lect. Notes in Math., vol. 730, Springer, Berlin, 1979, p. 204–227.
- [KLP16] M. KAPOVICH, B. LEEB & J. PORTI – “Some recent results on Anosov representations”, *Transform. Groups* **21** (2016), no. 4, p. 1105–1121.
- [KLP17] ———, “Anosov subgroups: dynamical and geometric characterizations”, *European J. Math.* **3** (2017), no. 4, p. 808–898.
- [KLP18a] ———, “Dynamics on flag manifolds: domains of proper discontinuity and cocompactness”, *Geom. Topol.* **22** (2018), no. 1, p. 157–234.
- [KLP18b] ———, “A Morse lemma for quasigeodesics in symmetric spaces and Euclidean buildings”, *Geom. Topol.* **22** (2018), no. 7, p. 3827–3923.
- [Lab06] F. LABOURIE – “Anosov flows, surface groups and curves in projective space”, *Invent. Math.* **165** (2006), no. 1, p. 51–114.
- [Led84] F. LEDRAPPIER – “Quelques propriétés des exposants caractéristiques”, in *École d’été de probabilités de Saint-Flour, XII–1982*, Lect. Notes in Math., vol. 1097, Springer, Berlin, 1984, p. 305–396.
- [LL23] F. LEDRAPPIER & P. LESSA – “Exact dimension of Furstenberg measures”, *Geom. Funct. Anal.* **33** (2023), no. 1, p. 245–298.
- [LL24] ———, “Exact dimension of dynamical stationary measures”, *J. Modern Dyn.* **20** (2024), p. 679–715.
- [LPX23] J. LI, W. PAN & D. XU – “On the dimension of limit sets on $\mathbb{P}(\mathbb{R}^3)$ via stationary measures: the theory and applications”, 2023, [arXiv:2311.10265](https://arxiv.org/abs/2311.10265).
- [Lin04] G. LINK – “Hausdorff dimension of limit sets of discrete subgroups of higher rank Lie groups”, *Geom. Funct. Anal.* **14** (2004), no. 2, p. 400–432.
- [MS21] I. D. MORRIS & Ç. SERT – “A strongly irreducible affine iterated function system with two invariant measures of maximal dimension”, *Ergodic Theory Dynam. Systems* **41** (2021), no. 11, p. 3417–3438.

- [MS19] I. D. MORRIS & P. SHMERKIN – “On equality of Hausdorff and affinity dimensions, via self-affine measures on positive subsystems”, *Trans. Amer. Math. Soc.* **371** (2019), no. 3, p. 1547–1582.
- [Ose68] V. I. OSELEDETS – “A multiplicative ergodic theorem. Characteristic Ljapunov exponents of dynamical systems”, *Trudy Moskov. Mat. Obšč.* **19** (1968), p. 179–210.
- [PSW21] M. B. POZZETTI, A. SAMBARINO & A. WIENHARD – “Conformality for a robust class of non-conformal attractors”, *J. reine angew. Math.* **774** (2021), p. 1–51.
- [PSW23] ———, “Anosov representations with Lipschitz limit set”, *Geom. Topol.* **27** (2023), no. 8, p. 3303–3360.
- [Qui02] J.-F. QUINT – “Divergence exponentielle des sous-groupes discrets en rang supérieur”, *Comment. Math. Helv.* **77** (2002), no. 3, p. 563–608.
- [Rap24] A. RAPAPORT – “On self-affine measures associated to strongly irreducible and proximal systems”, *Adv. Math.* **449** (2024), article no. 109734 (116 pages).
- [Rue79] D. RUELE – “Ergodic theory of differentiable dynamical systems”, *Publ. Math. Inst. Hautes Études Sci.* **50** (1979), p. 27–58.
- [Sam14] A. SAMBARINO – “Hyperconvex representations and exponential growth”, *Ergodic Theory Dynam. Systems* **34** (2014), no. 3, p. 986–1010.
- [Sim15] B. SIMON – *Harmonic analysis*, A comprehensive course in analysis, vol. 3, American Mathematical Society, Providence, RI, 2015.
- [Sul79] D. SULLIVAN – “The density at infinity of a discrete group of hyperbolic motions”, *Publ. Math. Inst. Hautes Études Sci.* **50** (1979), p. 171–202.
- [Var19] P. P. VARJÚ – “On the dimension of Bernoulli convolutions for all transcendental parameters”, *Ann. of Math. (2)* **189** (2019), no. 3, p. 1001–1011.
- [Yan19] W.-Y. YANG – “Statistically convex-cocompact actions of groups with contracting elements”, *Internat. Math. Res. Notices* (2019), no. 23, p. 7259–7323.
- [Zha97] Y. ZHANG – “Dynamical upper bounds for Hausdorff dimension of invariant sets”, *Ergodic Theory Dynam. Systems* **17** (1997), no. 3, p. 739–756.

Manuscript received 19th December 2023

accepted 9th December 2024

FRANÇOIS LEDRAPPIER, Sorbonne Université, UMR 8001, LPSM,
Boîte Courrier 158, 4 Place Jussieu, 75252 Paris cedex 05, France
E-mail : fledrapp@nd.edu

PABLO LESSA, CMAT, Facultad de Ciencias,
Iguá 4225, 11400 Montevideo, Uruguay
E-mail : lessa@cmat.edu.uy
Url : <https://www.cmat.edu.uy/~lessa/>