VANISHING OF MAPS FROM A PERFECT OBJECT IS UNAFFECTED BY BOUNDED TRUNCATION

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied from the pinned Stacks Project chapter file perfect.tex; no statement or proof step was rewritten, reordered or removed. The counted span is the author environment and its own proof environment. References inside the span are handed to the assembler as navigation stubs that carry the canonical tag of the cited tag only; whether each of them is load-bearing is recorded on the stub itself.

Allowed standard prerequisites (editorial).

- The canonical truncation triangle tau_{< a} E -> E -> tau_{>= a} E -> gives a long exact sequence on Hom groups. used at proof lines 10684-10689. Relates the two Hom-vanishing statements.
- Maps from a perfect object into an object of D^b_QCoh vanish in large degree. invoked at proof lines 10690-10691. Gives the needed vanishing for the lower truncation.

Source reference. Original lemma 09M4: sources/stacks-perfect/pinned-original-source.tex, source lines 3465–3465. Cited at proof lines 10690-10691 for the vanishing into the lower truncation; the reduction via the truncation triangle is inside the counted span.

Lemma 0.2. *Let X be a quasi-compact and quasi-separated scheme. Let $P \in D_{perf}(\mathcal{O}_X)$ and $E \in D_{QCoh}(\mathcal{O}_X)$. Let $a \in \mathbf{Z}$. The following are equivalent*

(1) $\mathrm{Hom}_{D(\mathcal{O}_X)}(P[-i], E) = 0$ *for* $i \gg 0$, *and*
(2) $\mathrm{Hom}_{D(\mathcal{O}_X)}(P[-i], \tau_{\geq a}E) = 0$ *for* $i \gg 0$.

Proof. Using the triangle $\tau_{<a}E \to E \to \tau_{\geq a}E \to$ we see that the equivalence follows if we can show

$$\mathrm{Hom}_{D(\mathcal{O}_X)}(P[-i], \tau_{<a}E) = \mathrm{Hom}_{D(\mathcal{O}_X)}(P, (\tau_{<a}E)[i]) = 0$$

for $i \gg 0$. As P is perfect this is true by Lemma 09M4. □

Date: stacks-project:perfect.tex:local-snapshot-20261012; GNU FDL 1.2 or later.