Spanning path plus an isolated edge has the perfect-matching anti-Ramsey number

Ali Ghalavand

arxiv:2610.11441v1; CC BY 4.0

Editorial note. Complete original author body and embedded bibliography retained; one selected primary per task. No mathematical alteration, proof repair, added assumption or correctness certification.

Allowed standard prerequisites (editorial).

- Anti-Ramsey numbers and directed path extremal theory. Original terminology, cited perfect matching formula and directed-cycle bounds, and full native auxiliary proofs retained.. The selected arc-bound or matching-label optimization argument remains counted in full.

1 Introduction

All undirected graphs considered in this paper are finite and simple. For a graph G, its vertex and edge sets are denoted by $V(G)$ and $E(G)$, and its order and size are $|V(G)|$ and $e(G) = |E(G)|$, respectively. The complete graph on n vertices is denoted by K_n, and P_n denotes the path on n vertices, so that P_n has $n-1$ edges. The symbol $G \cup H$ denotes the vertex-disjoint union of two graphs, and kG denotes the disjoint union of k copies of G. We write $G \cong H$ for graph isomorphism and $G \subseteq H$ when H contains a subgraph isomorphic to G; this subgraph need not be induced. A linear forest is a graph whose components are paths. An isolated edge means a component isomorphic to P_2. We call a linear forest *even* if every component has an even number of vertices. A perfect matching is a set of pairwise vertex-disjoint edges covering all vertices. When a matching M is contained in a graph G, we also use M for its edge set, and $G - M$ denotes deletion of those edges while retaining all vertices.

An edge-coloring c assigns a color to each edge of a graph; it is not required to be proper. A subgraph is *rainbow* if all its edges have distinct colors. For a graph F of order at most n, the anti-Ramsey number $\mathrm{AR}(n, F)$ is the maximum number of colors in an edge-coloring of K_n with no rainbow copy of F. A *representative graph* of a coloring is a spanning subgraph obtained by choosing exactly one edge of each color. It is therefore rainbow and has as many edges as the coloring has colors.

Anti-Ramsey theory was introduced by Erdős, Simonovits, and Sós [3]. Classical results on paths and cycles were obtained by Simonovits and Sós [16] and Montellano-Ballesteros and Neumann-Lara [14], respectively. For matchings, Schiermeyer [15] determined the rainbow threshold for sufficiently large host orders. Chen, Li, and Tu [2] and Fujita, Kaneko, Schiermeyer, and Suzuki [5] determined the anti-Ramsey numbers of tP_2 for $n \geq 2t+1$. Perfect matchings form the remaining spanning case, which was settled by Haas and Young [9]. Jahanbekam and West [11] studied the related packing problem of forcing edge-disjoint rainbow spanning subgraphs, including perfect matchings, cycles, and trees. These results provide a broader setting for the study of rainbow structures that cover the entire host graph.

Linear forests have also received substantial attention in anti-Ramsey theory. Bialostocki, Gilboa, and Roditty [1] determined the anti-Ramsey numbers of graphs with at most four edges, including $P_3 \cup P_2$, $P_3 \cup 2P_2$, and $P_4 \cup P_2$. Gilboa and Roditty [8] developed methods for studying the disjoint union of a fixed graph with copies of P_2 or P_3. For sufficiently large host orders, they obtained exact values for several short-path forests and bounds for a path together with a matching. Fang, Győri, Lu, and Xiao [4] obtained bounds for linear forests and determined the exact anti-Ramsey number of $2P_4$. Xie and Yuan [17] subsequently determined the exact anti-Ramsey numbers of fixed linear forests containing an even-order path for sufficiently large complete host graphs, and characterized the extremal colorings. The spanning case, in which the forest uses every vertex of the host graph, requires an exact treatment at the smallest possible host order.

Recent work has focused on forests formed by short paths and isolated edges. He and Jin [10] determined anti-Ramsey numbers for $P_3 \cup tP_2$ and $2P_3 \cup tP_2$ in specified ranges of the host order. Jie, He, and Jin [12] determined anti-Ramsey numbers of $kP_3 \cup tP_2$ for $k \geq 2$, $t \geq (k^2-k+4)/2$, and $n \geq 3k+2t+1$. Jie and Jin [13] investigated spanning forests of the form $kP_3 \cup tP_2$ under a condition relating the numbers of the two types of components. Ghalavand and Li [7] determined $\mathrm{AR}(3k+2t, kP_3 \cup tP_2)$ for all $k \geq 1$ and $t \geq 2$, without an additional relation between k and t. These spanning results concern forests with three-vertex path components. For even components, Ghalavand, Jie, Jin, Li, and Pan [6] established the following result for spanning forests consisting of paths on four vertices and isolated edges.

Theorem 1.1. [6] *If $k, t \geq 1$, then*

$$\mathrm{AR}(4k+2t, kP_4 \cup tP_2) = \mathrm{AR}(4k+2t, (2k+t)P_2).$$

The present paper extends Theorem 1.1 to arbitrary even component orders. More precisely, we prove that if

$$F \cong P_{2a_1} \cup \cdots \cup P_{2a_q} \cup tP_2, \qquad a_i \geq 2, \quad t \geq 1,$$

where $q \geq 0$ and $n = 2\sum_{i=1}^{q} a_i + 2t \geq 6$, then $\mathrm{AR}(n, F) = \mathrm{AR}(n, \frac{n}{2}P_2)$. Here the union and sum over the longer paths are empty when $q = 0$, so this statement also includes perfect matchings. No equality among the integers $a_1, \ldots, a_q$ is assumed.

Every even linear forest F on $2s$ vertices with an isolated edge satisfies $sP_2 \subseteq F \subseteq P_{2s-2} \cup P_2$. Thus it is enough to determine $\mathrm{AR}(2s, P_{2s-2} \cup P_2)$. We use a representative graph containing a rainbow perfect matching and select endpoint labels that maximize the number of arcs in an associated digraph. Comparing this labeling with the result of swapping the labels on a single matching edge shows that at least half of the nonmatching edges contribute arcs. The directed extremal results of Zhou and Li [18], together with the auxiliary bounds proved here, then yield a long directed path. Expanding its vertices back into matching edges produces a rainbow path disjoint from one unused matching edge.

We specify the directed notation used in that reduction. A digraph D has a vertex set $V(D)$ and an arc set $A(D)$, with no loops and at most one arc in each direction between two distinct vertices. Opposite arcs are allowed. Its number of arcs is $a(D) = |A(D)|$. The notation $u \to v$ means that $(u, v) \in A(D)$. For a vertex v, its outdegree and indegree are $d_D^+(v)$ and $d_D^-(v)$, and its total degree is $d_D(v) = d_D^+(v) + d_D^-(v)$. For $U \subseteq V(D)$, the induced subdigraph on U is denoted by $D[U]$; $D - v$ means $D[V(D) \setminus \{v\}]$. For disjoint sets U, W, let $a_D(U, W)$ denote the total number of arcs with one endpoint in U and the other in W, counting both directions. When P is a path, $V(P)$ is its vertex set. The complete symmetric digraph on r vertices, denoted by $\overleftrightarrow{K}_r$, contains both arcs between every pair of distinct vertices. An isolated directed two-cycle is a component on two vertices containing both opposite arcs and having no arcs to or from other vertices. A directed Hamilton cycle visits every vertex exactly once before returning to its starting vertex. The directed path on n distinct vertices, all of whose arcs point along the path, is denoted by $\overrightarrow{P}_n$. Finally, $\mathrm{ex}_{\mathrm{dir}}(s, \overrightarrow{P}_n)$ is the maximum number of arcs in a digraph on s vertices containing no copy of $\overrightarrow{P}_n$. The notation $\lfloor x \rfloor$ denotes the integer part of x.

Section 2 contains the published results used in the proof and the auxiliary directed-path bounds. Section 3 determines $\mathrm{AR}(2s, P_{2s-2} \cup P_2)$ and deduces the result for spanning even linear forests. Section 4 concludes the paper.

2 Auxiliary lemmas

In this section, we collect the auxiliary results needed to prove our main theorems. We recall relevant results from the literature and establish the additional lemmas used in our proofs.

Lemma 2.1 (Haas and Young [9])**.** *If $s \geq 2$ is an integer, then*

$$\mathrm{AR}(2s, sP_2) = \begin{cases} 3, & s = 2, \\ \dfrac{(s-2)(3s+1)}{2} + 1, & 3 \leq s \leq 6, \\ (s-2)(2s-3) + 2, & s \geq 7. \end{cases}$$

The following Hamilton-cycle bound and its equality characterization are special cases of the directed-cycle results of Zhou and Li [18].

Lemma 2.2 (Zhou and Li [18]). *Let H be a digraph on $r \geq 3$ vertices without a directed Hamilton cycle. Then $a(H) \leq (r-1)^2$. If $r \geq 4$ and $a(H) = (r-1)^2$, then H is obtained from $\overleftrightarrow{K}_{r-1}$ by adding a vertex z and either all arcs from z to the other vertices or all arcs from the other vertices to z.*

Lemma 2.3. *Let $P = v_1 \to v_2 \to \cdots \to v_r$ be a longest directed path in a digraph D. For every $x \in V(D) \setminus V(P)$,*

$$a_D(\{x\}, V(P)) \leq r-1.$$

Proof. Since P is a longest directed path, neither $x \to v_1$ nor $v_r \to x$ can be present; otherwise P could be extended to a directed path on $r+1$ vertices. It remains to consider the arcs

$$v_i \to x \quad (1 \leq i \leq r-1) \qquad \text{and} \qquad x \to v_{i+1} \quad (1 \leq i \leq r-1).$$

For each $i = 1, \ldots, r-1$, at most one of the two arcs

$$v_i \to x, \qquad x \to v_{i+1}$$

can be present. Indeed, if both were present, then

$$v_1 \to \cdots \to v_i \to x \to v_{i+1} \to \cdots \to v_r$$

would be a directed path on $r+1$ vertices, contradicting the maximality of P. These $r-1$ pairs are pairwise disjoint, and every remaining possible arc between x and $V(P)$ belongs to exactly one of them. Hence the number of arcs between x and $V(P)$ is at most $r-1$. □

Lemma 2.4 (Zhou and Li [18]). *For every integer $s \geq 3$,*

$$\mathrm{ex}_{\mathrm{dir}}(s, \overrightarrow{P}_3) = \left\lfloor \frac{s^2}{4} \right\rfloor.$$

Lemma 2.5. *Let D be a digraph on $r+2$ vertices, where $r \geq 4$. If a longest directed path in D has r vertices, then $a(D) \leq r^2$.*

Proof. Choose a longest directed path P in D, and put $H = D[V(P)]$ and $W = V(D) \setminus V(P) = \{x, y\}$. By Lemma 2.3,

$$a_D(W, V(P)) \leq 2(r-1). \tag{1}$$

Moreover,

$$a(D) = a(H) + a_D(W, V(P)) + a(D[W]), \qquad a(D[W]) \leq 2.$$

Suppose first that H contains a directed Hamilton cycle. No arc can join W to $V(H)$: an arc directed from an outside vertex to the cycle produces a directed path starting at that vertex and then visiting all r cycle vertices; an arc in the opposite direction similarly produces a directed path ending at the outside vertex. Either construction gives a directed path on $r+1$ vertices, contrary to the choice of P. Thus

$$a(D) \leq r(r-1) + 2 \leq r^2.$$

We may therefore assume that H contains no directed Hamilton cycle. Lemma 2.2 gives $a(H) \leq (r-1)^2$. If $a(H) \leq (r-1)^2 - 1$, or if $a(H) = (r-1)^2$ and $a(D[W]) \leq 1$, then

$$a(H) + a(D[W]) \leq (r-1)^2 + 1.$$

Together with (1), this yields

$$a(D) \leq (r-1)^2 + 1 + 2(r-1) = r^2.$$

It remains to consider the case $a(H) = (r-1)^2$ and $a(D[W]) = 2$. In particular, both $x \to y$ and $y \to x$ are present. By the equality statement in Lemma 2.2, there is a vertex $z \in V(H)$ such that, writing $C = V(H) \setminus \{z\}$, we have $D[C] \cong \overleftrightarrow{K}_{r-1}$, and either all arcs from z to C or all arcs from C to z are present, with no arcs in the opposite direction.

We first show that $a_D(W, C) = 0$. Take $u \in W$, and let w be the other vertex of W. If $u \to c$ is present for some $c \in C$, then $w \to u \to c$, followed by a directed Hamilton path of $D[C]$ starting at c, gives a directed path on $r+1$ vertices. If $c \to u$ is present, a directed Hamilton path of $D[C]$ ending at c, followed by $c \to u \to w$, gives the same contradiction. Hence no arc joins W to C.

Finally, suppose that all arcs between z and C are directed from z to C. For any $u \in W$, an arc $u \to z$ could be followed by a directed Hamilton path of H starting at z, giving a directed path on $r+1$ vertices. Thus only $z \to x$ and $z \to y$ are possible. If instead all arcs are directed from C to z, the analogous argument using a directed Hamilton path of H ending at z rules out $z \to x$ and $z \to y$. Consequently, in either case,

$$a_D(W, \{z\}) \le 2.$$

Therefore,

$$a(D) = a(H) + a_D(W, C) + a_D(W, \{z\}) + a(D[W]) \le (r-1)^2 + 4 < r^2,$$

where the final inequality follows from $r \ge 4$. This completes the proof. □

Lemma 2.6. *For every integer $s \ge 3$,*

$$\mathrm{ex}_{\mathrm{dir}}(s, \overrightarrow{P}_{s-1}) \le \begin{cases} 0, & s = 3, \\ 4, & s = 4, \\ 10, & s = 5, \\ (s-2)^2, & s \ge 6. \end{cases}$$

Proof. For $s = 3$, avoiding $\overrightarrow{P}_2$ means having no arcs. For $s = 4$, the required bound is Lemma 2.4.

Consider $s = 5$, and let D contain no $\overrightarrow{P}_4$. A longest path has at most three vertices. If it has at most two, Lemma 2.4 gives $a(D) \le 6$. Otherwise, let H be induced by its three vertices. If H has a Hamilton cycle, the argument in Lemma 2.5 shows that no arc joins it to the other two vertices, and $a(D) \le 6 + 2 = 8$. If it has no Hamilton cycle, Lemmas 2.2 and 2.3 give

$$a(D) \le (3-1)^2 + 2(3-1) + 2 = 10.$$

We prove the bound for $s \ge 6$ by induction on s. For the base case $s = 6$, let D contain no $\overrightarrow{P}_5$, and let $r \le 4$ be the order of a longest path P. If $r = 4$, Lemma 2.5 gives $a(D) \le 16$. If $r \le 2$, Lemma 2.4 gives $a(D) \le 9$. If $r = 3$, put $H = D[V(P)]$. When H has a Hamilton cycle, there are no arcs between H and its complement, so $a(D) \le 6 + 6 = 12$. Otherwise, Lemmas 2.2 and 2.3 give

$$a(D) \le (3-1)^2 + 3(3-1) + 3(3-1) = 16 = (6-2)^2.$$

The last term bounds all arcs among the three vertices outside P.

Now let $s \ge 7$ and assume the assertion for $s-1$. Let D contain no $\overrightarrow{P}_{s-1}$, and choose a longest path $P = v_1 \to \cdots \to v_r$. Then $r \le s-2$. If $r = s-2$, Lemma 2.5 gives the required bound directly. Suppose $r \le s-3$. The last vertex v_r has no outgoing arc to a vertex outside P. At most $2(r-1)$ arcs incident with v_r have their other endpoint on P, and at most $s-r$ have their other endpoint outside P. Thus

$$d_D(v_r) \le 2(r-1) + (s-r) = s + r - 2 \le 2s - 5.$$

Since every path in D has at most $s-3$ vertices, the digraph $D - v_r$ contains no $\overrightarrow{P}_{s-2}$. It has $s-1$ vertices, so the induction hypothesis gives $a(D - v_r) \le (s-3)^2$. Deleting a vertex removes exactly its total degree many arcs. Consequently,

$$a(D) = a(D - v_r) + d_D(v_r) \le (s-3)^2 + 2s - 5 = (s-2)^2.$$

This completes the induction. □

Lemma 2.7. *For every integer $s \ge 3$, $2\,\mathrm{ex}_{\mathrm{dir}}(s, \overrightarrow{P}_{s-1}) + s \le \mathrm{AR}(2s, sP_2)$.*

Proof. By Lemma 2.1, the values of $\mathrm{AR}(2s, sP_2)$ for $s = 3, 4, 5, 6$ are $6, 14, 25, 39$, respectively. Lemma 2.6 gives

$$2\,\mathrm{ex}_{\mathrm{dir}}(3, \overrightarrow{P}_2) + 3 \le 3 \le 6, \quad 2\,\mathrm{ex}_{\mathrm{dir}}(4, \overrightarrow{P}_3) + 4 \le 12 \le 14,$$
$$2\,\mathrm{ex}_{\mathrm{dir}}(5, \overrightarrow{P}_4) + 5 \le 25, \qquad 2\,\mathrm{ex}_{\mathrm{dir}}(6, \overrightarrow{P}_5) + 6 \le 38 \le 39.$$

For $s \ge 7$, the two lemmas give

$$2\,\mathrm{ex}_{\mathrm{dir}}(s, \overrightarrow{P}_{s-1}) + s \le 2(s-2)^2 + s = (s-2)(2s-3) + 2 = \mathrm{AR}(2s, sP_2).$$

□

3 Main results

We first consider $P_{2s-2} \cup P_2$, which contains every even linear forest on $2s$ vertices having an isolated edge as a spanning subgraph. The proof combines the directed-path bounds from Section 2 with an endpoint labeling of a rainbow perfect matching that maximizes the arc count of an auxiliary digraph. The result for all such forests then follows by subgraph monotonicity and the perfect-matching formula.

Theorem 3.1. *For every integer $s \geq 3$,*

$$\mathrm{AR}(2s, P_{2s-2} \cup P_2) = \mathrm{AR}(2s, sP_2).$$

Proof. The forest $P_{2s-2} \cup P_2$ contains a perfect matching. Consequently, every edge-coloring of K_{2s} without a rainbow sP_2 also avoids a rainbow $P_{2s-2} \cup P_2$. Thus

$$\mathrm{AR}(2s, P_{2s-2} \cup P_2) \geq \mathrm{AR}(2s, sP_2).$$

For the reverse inequality, consider an edge-coloring of K_{2s} using $q > \mathrm{AR}(2s, sP_2)$ colors. By the definition of the anti-Ramsey number, this coloring contains a rainbow perfect matching M.

Choose one representative edge of each color, choosing the corresponding edge of M whenever that color appears on M. These choices are compatible because the edges of M have distinct colors. The resulting spanning graph G contains M, satisfies $e(G) = q$, and has pairwise distinctly colored edges. Write

$$M = \{u_i v_i : 1 \leq i \leq s\}.$$

For each i, there are two ways to assign the endpoint labels a_i, b_i:

$$(a_i, b_i) = (u_i, v_i) \quad \text{or} \quad (a_i, b_i) = (v_i, u_i).$$

Hence there are 2^s possible complete labelings of M. For each labeling, define a digraph D on $\{1, \ldots, s\}$ by

$$i \to j \quad \Longleftrightarrow \quad b_i a_j \in E(G), \qquad i \neq j.$$

The direction goes from the index of the b-endpoint to the index of the a-endpoint, regardless of the numerical order of the indices. More explicitly, for two distinct matching edges indexed by i and j,

Edge of G	Contribution to D
$b_i a_j = a_j b_i$	$i \to j$
$a_i b_j = b_j a_i$	$j \to i$
$a_i a_j$	no arc
$b_i b_j$	no arc

Thus each nonmatching edge contributes exactly one arc when its endpoints have different label types, and no arc otherwise. Reversing the written order of its endpoints does not create another arc.

The matching edges contribute no arcs because $i \neq j$ is required. Moreover, for each ordered pair (i, j), only the edge $b_i a_j$ can supply the arc $i \to j$. Therefore D is a simple loopless digraph, and distinct contributing edges supply distinct arcs. Write $a(D)$ for its number of arcs.

Claim 3.2. *There exists an endpoint labeling of M for which*

$$a(D) \geq \frac{q - s}{2} > \frac{\mathrm{AR}(2s, sP_2) - s}{2}.$$

Proof of the claim. Choose a labeling maximizing $a(D)$, which is possible because there are only 2^s labelings.

For each i, let t_i be the number of edges in $E(G) \setminus E(M)$ having one endpoint in $\{a_i, b_i\}$, and let c_i count those edges that contribute arcs to D. Equivalently, c_i is the number of arcs of D incident with i.

Interchange the labels a_i and b_i, keeping all other labels fixed, and let D_i be the resulting digraph. Each nonmatching edge incident with $\{a_i, b_i\}$ has exactly one endpoint whose label type changes. After the interchange, each of the c_i previously contributing edges has either two a-endpoints or two b-endpoints and therefore contributes no arc. Each of the remaining $t_i - c_i$ edges now has one a-endpoint and one b-endpoint and consequently contributes exactly one arc. All arcs with neither endpoint i remain unchanged. Hence

$$a(D_i) = a(D) - c_i + (t_i - c_i) = a(D) + t_i - 2c_i.$$

By the maximality of $a(D)$, we have $a(D_i) \le a(D)$. Therefore

$$2c_i \ge t_i \qquad (1 \le i \le s).$$

Every arc of D is counted at each of its two ends, so

$$\sum_{i=1}^{s} c_i = 2a(D).$$

Similarly, every nonmatching edge joins two distinct matching edges and is counted in the corresponding two quantities t_i. Thus

$$\sum_{i=1}^{s} t_i = 2|E(G) \setminus E(M)| = 2(q-s).$$

Summing $2c_i \ge t_i$ gives

$$4a(D) = 2\sum_{i=1}^{s} c_i \ge \sum_{i=1}^{s} t_i = 2(q-s).$$

Consequently,

$$a(D) \ge \frac{q-s}{2} > \frac{\mathrm{AR}(2s, sP_2) - s}{2},$$

where the strict inequality follows from $q > \mathrm{AR}(2s, sP_2)$. □

Fix the labeling supplied by Claim 3.2. By the directed-path arc bound and Lemma 2.7, every digraph on s vertices containing no directed path on $s-1$ vertices has at most

$$\frac{\mathrm{AR}(2s, sP_2) - s}{2}$$

arcs. Our digraph exceeds this bound, so it contains a directed path

$$i_1 \to i_2 \to \cdots \to i_{s-1},$$

where the indices are distinct.

For each $1 \le h \le s-2$, the definition of D gives

$$b_{i_h} a_{i_{h+1}} \in E(G).$$

Combining these connecting edges with the matching edges $a_{i_h} b_{i_h}$, for $1 \le h \le s-1$, produces the path

$$a_{i_1} b_{i_1} a_{i_2} b_{i_2} \cdots a_{i_{s-1}} b_{i_{s-1}}$$

in G. Its $2s-2$ vertices are distinct because the indices are distinct and the matching edges are pairwise disjoint.

Exactly one index j is absent from $\{i_1, \ldots, i_{s-1}\}$. The matching edge $a_j b_j$ is disjoint from this path. Together, the path and this unused matching edge form a copy of $P_{2s-2} \cup P_2$.

All edges of this copy belong to G and therefore have distinct colors. Thus the copy is rainbow. We have shown that every edge-coloring of K_{2s} using more than $\mathrm{AR}(2s, sP_2)$ colors contains a rainbow $P_{2s-2} \cup P_2$. Consequently,

$$\mathrm{AR}(2s, P_{2s-2} \cup P_2) \le \mathrm{AR}(2s, sP_2).$$

Together with the reverse inequality established above, this proves the theorem. □

Theorem 3.3. *Let F be an even linear forest on $n = 2s \ge 6$ vertices, and suppose that F has at least one component isomorphic to P_2. Then*

$$\mathrm{AR}(n, F) = \mathrm{AR}(n, sP_2) = \begin{cases} \dfrac{(s-2)(3s+1)}{2} + 1, & 3 \le s \le 6, \\ (s-2)(2s-3) + 2, & s \ge 7. \end{cases}$$

Proof. Every even-order path has a perfect matching, obtained by pairing consecutive vertices along the path. Taking these matchings over all components gives $sP_2 \subseteq F$.

Choose one P_2 component of F. Since the remaining path components have total order $n-2$, partition the vertices of P_{n-2} into consecutive blocks whose orders equal those of these components. Deleting the edges between successive blocks leaves paths isomorphic to the remaining components of F. Together with the separate P_2, these paths form a spanning copy of F in $P_{n-2} \cup P_2$.

If $H \subseteq H'$, every coloring avoiding a rainbow H also avoids a rainbow H', so $\mathrm{AR}(n, H) \leq \mathrm{AR}(n, H')$. Applying this observation to the two containments and using Theorem 3.1 gives

$$\mathrm{AR}(n, sP_2) \leq \mathrm{AR}(n, F) \leq \mathrm{AR}(n, P_{n-2} \cup P_2) = \mathrm{AR}(n, sP_2).$$

The equality of the anti-Ramsey numbers follows, and their explicit values are given by Lemma 2.1. □

4 Conclusion

Every spanning even linear forest with an isolated edge has the same anti-Ramsey number as a perfect matching on the same vertex set. Thus, within this family, the value depends only on the total order, regardless of the number and orders of the path components. The proof reduces the problem to $P_{n-2} \cup P_2$ and combines a maximizing endpoint labeling with directed-path bounds. This extends the known result for $kP_4 \cup tP_2$ to arbitrary even component orders.

Conflict of interest

The authors declare that they have no conflict of interest.

Data availability

No data was used in this investigation.

Use of artificial intelligence

The author used AI tools for algebraic checks, language editing, locating relevant auxiliary results, and examining proof ideas. The author reviewed and verified all mathematical statements, calculations, proofs, and references and takes full responsibility for the manuscript.

References

[1] A. Bialostocki, S. Gilboa, Y. Roditty, Anti-Ramsey numbers of small graphs, Ars Combin. 123 (2015) 41–53.

[2] H. Chen, X. Li, J. Tu, Complete solution for the rainbow numbers of matchings, Discrete Math. 309 (2009) 3370–3380. https://doi.org/10.1016/j.disc.2008.10.002.

[3] P. Erdős, M. Simonovits, V.T. Sós, Anti-Ramsey theorems, in: Infinite and Finite Sets, Vol. II, Colloq. Math. Soc. János Bolyai, Vol. 10, North-Holland, Amsterdam, 1975, pp. 633–643.

[4] C. Fang, E. Győri, M. Lu, J. Xiao, On the anti-Ramsey number of forests, Discrete Appl. Math. 291 (2021) 129–142. https://doi.org/10.1016/j.dam.2020.08.027.

[5] S. Fujita, A. Kaneko, I. Schiermeyer, K. Suzuki, A rainbow k-matching in the complete graph with r colors, Electron. J. Combin. 16 (2009) R51. https://doi.org/10.37236/140.

[6] A. Ghalavand, Q. Jie, Z. Jin, X. Li, L. Pan, On the anti-Ramsey number under edge deletion, Bull. Malays. Math. Sci. Soc. 49 (2026) 147. https://doi.org/10.1007/s40840-026-02147-w.

[7] A. Ghalavand, X. Li, Anti-Ramsey numbers for spanning linear forests of 3-vertex paths and matchings, arXiv:2509.25949, 2025, revised 2026. https://doi.org/10.48550/arXiv.2509.25949.

[8] S. Gilboa, Y. Roditty, Anti-Ramsey numbers of graphs with small connected components, Graphs Combin. 32 (2016) 649–662. https://doi.org/10.1007/s00373-015-1581-y.

[9] R. Haas, M. Young, The anti-Ramsey number of perfect matching, Discrete Math. 312 (2012) 933–937. https://doi.org/10.1016/j.disc.2011.10.017.

[10] M. He, Z. Jin, Rainbow short linear forests in edge-colored complete graph, Discrete Appl. Math. 361 (2025) 523–536. https://doi.org/10.1016/j.dam.2024.11.002.

[11] S. Jahanbekam, D.B. West, Anti-Ramsey problems for t edge-disjoint rainbow spanning subgraphs: cycles, matchings, or trees, J. Graph Theory 82 (2016) 75–89. https://doi.org/10.1002/jgt.21888.

[12] Q. Jie, M. He, Z. Jin, Rainbow forest consisting of short paths in K_n, Discrete Appl. Math. 376 (2025) 260–269. https://doi.org/10.1016/j.dam.2025.06.041.

[13] Q. Jie, Z. Jin, Rainbow-free colorings for spanning linear forest consisting of short paths, Discrete Appl. Math. 386 (2026) 30–57. https://doi.org/10.1016/j.dam.2026.01.016.

[14] J.J. Montellano-Ballesteros, V. Neumann-Lara, An anti-Ramsey theorem on cycles, Graphs Combin. 21 (2005) 343–354. https://doi.org/10.1007/s00373-005-0619-y.

[15] I. Schiermeyer, Rainbow numbers for matchings and complete graphs, Discrete Math. 286 (2004) 157–162. https://doi.org/10.1016/j.disc.2003.11.057.

[16] M. Simonovits, V.T. Sós, On restricted colourings of K_n, Combinatorica 4 (1984) 101–110. https://doi.org/10.1007/BF02579162.

[17] T.-Y. Xie, L.-T. Yuan, On the anti-Ramsey numbers of linear forests, Discrete Math. 343 (2020) 112130. https://doi.org/10.1016/j.disc.2020.112130.

[18] W. Zhou, B. Li, The Turán number of directed paths and oriented cycles, Graphs Combin. 39 (2023) 47. https://doi.org/10.1007/s00373-023-02647-7.