AN INTEGRAL MORPHISM HAS PROPERTY A

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied from the pinned Stacks Project chapter file etale-cohomology.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. The proof is the author's complete argument; the reduction and the construction are both inside the counted span.

Editorial note. Every labelled reference occurring inside the counted spans is carried in assembly.external_references as a non-load-bearing navigation pointer (same-chapter ones at their definition site, cross-chapter ones at the invocation inside the counted span), so no load-bearing mathematics is replaced by a stub. No bibliography citation occurs inside the counted spans.

Editorial note. The checked chapter list for same-name collisions is the set of stacks chapters staged in this repository: chow, cohomology, divisors, etale-cohomology, flat, more-algebra, more-morphisms, morphisms, spaces-more-morphisms, varieties together with this card's chapter etale-cohomology.

Editorial note. The preamble is the chapter's own native preamble excerpt already used for this source: the Xy-pic setup (native preamble lines 14-18) and the native theorem declarations with the mathematical macros (native preamble lines 154-217).

Allowed standard prerequisites (editorial).

- Property (A) for a morphism of schemes. Same chapter, the formulation of property (A) delivered verbatim as a context span in this card (source lines 5708-5726). Supplies the condition verified for an integral morphism.
- Affineness of an integral morphism and its behaviour under base change. Same chapter together with the chapter Morphisms, the description of an integral morphism as affine, which reduces the verification to rings. Supplies the reduction to the affine case in which the property is verified.

Source reference. Original lemma 00U2: sources/stacks-etale-cohomology/pinned-original-source.tex, source lines 5866–5866. Navigation pointer invoked at proof line 5866; the counted span contains the pointer only and the cited result is not reproduced inside it, so the pointer is registered as a non-load-bearing navigation stub in the established form for Stacks material. No bibliography citation is involved.

Source reference. Original lemma 04DL: sources/stacks-etale-cohomology/pinned-original-source.tex, source lines 5795–5795. Navigation pointer invoked at proof line 5872; the counted span contains the pointer only and the cited result is not reproduced inside it, so the pointer is registered as a non-load-bearing navigation stub in the established form for Stacks material. No bibliography citation is involved.

Let $f : X \to Y$ be a morphism of schemes. Here is a list of conditions we will consider in the following:

Date: stacks-project:etale-cohomology.tex:local-snapshot-20261011; GNU FDL 1.2 or later.

(A) For every étale morphism $U \to X$ and $u \in U$ there exist an étale morphism $V \to Y$ and a disjoint union decomposition $X \times_Y V = W \amalg W'$ and a morphism $h : W \to U$ over X with u in the image of h.
(B) For every $V \to Y$ étale, and every étale covering $\{U_i \to X \times_Y V\}$ there exists an étale covering $\{V_j \to V\}$ such that for each j we have $X \times_Y V_j = \coprod W_{ij}$ where $W_{ij} \to X \times_Y V$ factors through $U_i \to X \times_Y V$ for some i.
(C) For every $U \to X$ étale, there exists a $V \to Y$ étale and a surjective morphism $X \times_Y V \to U$ over X.

It turns out that each of these properties has meaning in terms of the behaviour of the functor $f_{small,*}$. We will work this out in the next few sections.

Lemma 0.1. *Let $f : X \to Y$ be an integral morphism of schemes. Then property (A) holds.*

Proof. Let $U \to X$ be étale, and let $u \in U$ be a point. We have to find $V \to Y$ étale, a disjoint union decomposition $X \times_Y V = W \amalg W'$ and an X-morphism $W \to U$ with u in the image. We may shrink U and Y and assume U and Y are affine. In this case also X is affine, since an integral morphism is affine by definition. Write $Y = \mathrm{Spec}(A)$, $X = \mathrm{Spec}(B)$ and $U = \mathrm{Spec}(C)$. Then $A \to B$ is an integral ring map, and $B \to C$ is an étale ring map. By Algebra, Lemma 00U2 we can find a finite A-subalgebra $B' \subset B$ and an étale ring map $B' \to C'$ such that $C = B \otimes_{B'} C'$. Thus the question reduces to the étale morphism $U' = \mathrm{Spec}(C') \to X' = \mathrm{Spec}(B')$ over the finite morphism $X' \to Y$. In this case the result follows from Lemma 04DL. □