Upper bound for the Hausdorff dimension of the limit set

Olivier Glorieux, Daniel Monclair, Nicolas Tholozan

doi:10.5802/jep.241; CC BY 4.0

Editorial note. All delivered mathematical text is the author's own, copied from the pinned journal source file sources/geometry-fl03-hausdorff-dimension-of-limit-sets/pinned-original-source.tex (Hausdorff dimension of limit sets for projective Anosov representations); no statement or proof step was rewritten, reordered or removed.

Editorial note. The counted proof is the author's complete argument: All delivered mathematical text is the author own, copied verbatim from the pinned journal source. Provenance note: the candidate comes from the lane-local candidate table lane-s12/candidates-geometry-fl03-hausdorff-dimension-of-limit-sets.json (patch 35). Statement source lines 1283-1286 with the author own proof at 1371-1389, which the author places remotely under a proof environment naming this statement (the face was flagged as a label mismatch precisely because its proof is not adjacent). Printed number 4.1 from the source walk, cross-checked against the published anchor of this DOI (theorem 3.1 at statement 1100-1103, whose walk value 3.1 matches the article); verified against the host aux.

Editorial note. The label th - upper bound names the counted statement, and the proof environment immediately following it in the source is delivered verbatim as its proof.

Editorial note. The counted statement is carried by the author's own theorem environment in the article source.

Editorial note. Journal-unit assembly: card-local preamble (author theorem environments and macros verbatim), article class, bibliography backend none; any literature citation carried by the delivered span is disclosed in the notes of this card.

Editorial note. Unresolved citation keys not present in the author's own bibliography: Gromov.

Editorial note. The bibliographic strings were transliterated to ASCII for the card face (the journal line carries accents and typographic punctuation); the SOURCE.json metadata keeps the original form. No mathematical text is affected.

Editorial note. PRESCREEN RE-READ: Cite-flag re-read: the quality screen takes its citation list from the census row, which spans the whole surrounding section, but the counted statement and the remote proof span contain no citation

at all (checked line by line in the pinned source). The card therefore carries no unresolved citation.

Editorial note. Assembly shape note: the card-local preamble is the unit minimal preamble used by the other cards of this source (author package set, author macros, author theorem declarations with the shared counter). The verbatim retained preamble of the unit was rejected because it carries the publisher cedram class lines (datereceived / dateaccepted / dateepreuves and the smf titlepage macros); the class is not in the toolchain, and the assembler supplies the article class instead, so the first host assemble stopped on an undefined control sequence at the datereceived line. The counters are seeded so the counted statement prints the number the article gives it.

Editorial note. Assembly shape note: the card-local preamble is the unit minimal preamble used by the other cards of this source (author package set, author macros, author theorem declarations with the shared counter). The verbatim retained preamble of the unit was rejected because it carries the publisher cedram class lines (datereceived / dateaccepted / dateepreuves and the smf titlepage macros); the class is not in the toolchain, and the assembler supplies the article class instead, so the first host assemble stopped on an undefined control sequence at the datereceived line. The counters are seeded so the counted statement prints the number the article gives it.

Editorial scope. the counted statement and its own proof environment only

Allowed standard prerequisites (editorial).

- The quantification of the distortion of balls by proximal elements. Supplies background that the delivered proof uses. The entry is declared as a dependency; it is not re-proved on the card, and the load-bearing steps of the counted statement are carried out inside the delivered span.

Source reference. Original reference 4.5: sources/geometry-fl03-hausdorff-dimension-of-limit-sets/pinned-original-source.tex, source lines 1357–1357. Navigation pointer invoked at line(s) 1376: the counted text refers to this environment of the same journal unit, which is not retained in this card, so the reference is declared as a pointer carrying the number the article itself prints for it (the section and the place of the statement inside that section, derived from the shared section-scoped theorem counter of the source); the cited item is an allowed input and none of it is reproduced inside the span.

Source reference. Original reference 4.3: sources/geometry-fl03-hausdorff-dimension-of-limit-sets/pinned-original-source.tex, source lines 1317–1317. Navigation pointer invoked at line(s) 1374: the counted text refers to this environment of the same journal unit, which is not retained in this card, so the reference is declared as a pointer carrying the number the article itself prints for it (the section and the place of the statement inside that section, derived from the shared section-scoped theorem counter of the source); the cited item is an allowed input and none of it is reproduced inside the span.

Theorem 4.1. *Let* $\Gamma \subset \mathrm{SL}(n,\mathbb{R})$ *be a projective Anosov subgroup. Then*

$$\mathrm{DimH}(\Lambda_\Gamma) \leq \delta_{1,2}(\Gamma).$$

Proof of Theorem 4.1. By compactness of Λ_Γ it is sufficient to prove that, for all $x \in \Lambda_\Gamma$, there exists $r > 0$ such that the Hausdorff dimension of $\Lambda_\Gamma \cap B(x, r)$ is less than $\delta_{1,2}(\Gamma)$.

Given such and x, let us fix y, r, U, and V as before. Let $C > 0$ be given by Lemma 4.3.

By Lemma 4.5, we have $\Lambda_\Gamma \cap B(x, r) \subset \bigcup_{\gamma \in \Gamma_\varepsilon} \gamma \cdot B(x, r)$. By definition of Γ_ε, this gives a covering of $\Lambda_\Gamma \cap B(x, r)$ by balls of radius less than εr. Note that every $\gamma \in \Gamma_\varepsilon$ is ε-Lipschitz on $B(L_+(\gamma), 4r) \subset B(x, 5r)$, hence it is $(4r, \varepsilon)$-proximal. By Lemma 4.3, it is thus $Ce^{(\mu_2 - \mu_1)(\gamma)}$-Lipschitz on $B(x, r) \subset B(L_+(\gamma), 2r)$, for some constant C.

Let $s > 0$. By definition of the s-dimensional Hausdorff measure, we have:

$$\begin{aligned} H^s_{r\varepsilon}(\Lambda_\Gamma \cap B(x, r)) \leq \sum_{\gamma \in \Gamma_\varepsilon} \operatorname{diam}(\gamma \cdot B(x, r))^s &\leq C^s \sum_{\gamma \in \Gamma_\varepsilon} e^{s(\mu_2 - \mu_1)(\gamma)} \\ &\leq C^s \sum_{\gamma \in \Gamma} e^{-s(\mu_1 - \mu_2)(\gamma)}. \end{aligned}$$

Since ε can be taken arbitrarily small, we obtain :

$$H^s(\Lambda_\Gamma \cap B) \leq C^s \sum_{\gamma \in \Gamma} e^{-s(\mu_1 - \mu_2)(\gamma)}.$$

Therefore, for all $s > \delta_{1,2}$, $H^s(\Lambda_\Gamma \cap B(x, r)) < +\infty$ which in turn implies that $\operatorname{DimH}(\Lambda_\Gamma \cap B(x, r)) \leq \delta_{1,2}$. □