PULLBACK OF A MODULE: $F^{-1}\MATHCAL{G}$ CARRIES A NATURAL $F^{-1}\MATHCAL{O}$-MODULE STRUCTURE, FUNCTORIAL IN $\MATHCAL{G}$

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied verbatim from the pinned Stacks Project chapter file sites-modules.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. Span recomputation by label pairing (LANE_BRIEF 10.2): in this chapter the author's proof environment follows the statement environment as a standalone block; the spec covers the complete environment pair (statement span, proof span) as recorded in the pool candidate and re-verified against the pinned source (sm-twins-20261014.json span sanity: all OK).

Editorial note. Occupation scan: combined claim_key + statement_anchor + generic-line scan over all registration files (active and withdrawn) found hard occupation only from the 47 pre-existing lane-s11 S11-sites-mod-* cards (same pinned snapshot, delivered) and cross-chapter citation noise, 0 other-lane hits for sites-modules.tex (sm-occupied.json, 2026-10-14); the RESUME-pool dispatch unlocks the remaining face.

Lemma 0.1. *Let $\mathcal{C}$, $\mathcal{D}$ be sites. Let $f : Sh(\mathcal{C}) \to Sh(\mathcal{D})$ be a morphism of topoi. Let $\mathcal{O}$ be a sheaf of rings on $\mathcal{D}$. Let $\mathcal{G}$ be a sheaf of $\mathcal{O}$-modules. There is a natural map of sheaves of sets*

$$f^{-1}\mathcal{O} \times f^{-1}\mathcal{G} \longrightarrow f^{-1}\mathcal{G}$$

which turns $f^{-1}\mathcal{G}$ into a sheaf of $f^{-1}\mathcal{O}$-modules. This construction is functorial in $\mathcal{G}$.

Proof. Denote $\mu : \mathcal{O} \times \mathcal{G} \to \mathcal{G}$ the multiplication map. Recall that f^{-1} (on sheaves of sets) is exact and hence commutes with products. Hence $f^{-1}\mu$ is a map as indicated. This proves the lemma. □

Date: stacks-project:sites-modules.tex:local-snapshot-20261012; GNU FDL 1.2 or later.