SUBSHEAVES OF THE FINAL OBJECT OF THE ETALE TOPOS ARE REPRESENTABLE BY UNIQUE OPENS

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied verbatim from the pinned Stacks Project chapter file spaces-properties.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. Span recomputation by label pairing (LANE_BRIEF 10.2): in this chapter the author's proof environment follows the statement environment as a standalone block; the spec covers the complete environment pair (statement span, proof span) as recorded in the pool candidate.

Editorial note. Occupation scan: combined claim_key + statement_anchor + generic-line scan over all registration files (active and withdrawn) found 13 S10-geom published cards for spaces-properties.tex (prop-occupied.json, 2026-10-14); those 13 labels are excluded from this batch by design. For the 77 free labels: 0 occupied. The registry mentions field (7557 files) and focused-seeds (645) both point to the same S10-geom cards (dual-source check), no other occupation.

Source reference. Original reference 00W3: sources/stacks-chow/tags.txt, source lines 1169–1169. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Source reference. Original lemma 03IR: sources/stacks-chow/tags.txt, source lines 4613–4613. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Lemma 0.1. *Let S be a scheme. Let X be an algebraic space over S. Let $\mathcal{F}$ be a subsheaf of the final object of the étale topos of X (see Sites, Example 00W3). Then there exists a unique open $W \subset X$ such that $\mathcal{F} = h_W$.*

Proof. The condition means that $\mathcal{F}(U)$ is a singleton or empty for all $\varphi : U \to X$ in $\mathrm{Ob}(X_{spaces,étale})$. In particular local sections always glue. If $\mathcal{F}(U) \neq \emptyset$, then $\mathcal{F}(\varphi(U)) \neq \emptyset$ because $\varphi(U) \subset X$ is an open subspace (Lemma 03IR) and $\{\varphi : U \to \varphi(U)\}$ is a covering in $X_{spaces,étale}$. Take $W = \bigcup_{\varphi:U\to S,\mathcal{F}(U)\neq\emptyset} \varphi(U)$ to conclude. □

Date: stacks-project:spaces-properties.tex:local-snapshot-20261012; GNU FDL 1.2 or later.