STACKS, DEFORMATION THEORY: LEMMA-NL-REPRESENT-EXT-CLASS

THE STACKS PROJECT AUTHORS

Editorial note. Statement span is lines 2157-2167 of sources/stacks-defos/pinned-original-source.tex; proof span is lines 2169-2214, the complete author proof (44 body lines).

Editorial note. Navigation stubs: 1 cited-but-not-included labels are supplied as non-load-bearing pointers with their pinned tags where the snapshot tag index has them.

Editorial note. extra_packages is empty on purpose: the retained preamble already loads xy with options and a second declaration clashes in the assembler wrapper.

Editorial note. Text is retained verbatim, including the author's own typos and placeholders.

Editorial note. Queue index 14 of queue_defos.json (pb=2056, stmt 2157-2167, proof 2169-2214).

Allowed standard prerequisites (editorial).

- Explicit prerequisite boundary. The cited inputs are prerequisites; the argument in the retained span is not replaced by any cited result.

Source reference. Original lemma 08TL: sources/stacks-defos/pinned-original-source.tex, source lines 2214–2214. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Lemma 0.1. *Let X be a topological space. Let $\mathcal{A} \to \mathcal{B}$ be a homomorphism of sheaves of rings. Let $\mathcal{G}$ be a $\mathcal{B}$-module. Let $\xi \in \mathrm{Ext}^1_{\mathcal{B}}(NL_{\mathcal{B}/\mathcal{A}}, \mathcal{G})$. There exists a map of sheaves of sets $\alpha : \mathcal{E} \to \mathcal{B}$ such that $\xi \in \mathrm{Ext}^1_{\mathcal{B}}(NL(\alpha), \mathcal{G})$ is the class of a map $\mathcal{I}/\mathcal{I}^2 \to \mathcal{G}$ (see proof for notation).*

Proof. Recall that given $\alpha : \mathcal{E} \to \mathcal{B}$ such that $\mathcal{A}[\mathcal{E}] \to \mathcal{B}$ is surjective with kernel $\mathcal{I}$ the complex $NL(\alpha) = (\mathcal{I}/\mathcal{I}^2 \to \Omega_{\mathcal{A}[\mathcal{E}]/\mathcal{A}} \otimes_{\mathcal{A}[\mathcal{E}]} \mathcal{B})$ is canonically isomorphic to $NL_{\mathcal{B}/\mathcal{A}}$, see Modules, Lemma 08TL. Observe moreover, that $\Omega = \Omega_{\mathcal{A}[\mathcal{E}]/\mathcal{A}} \otimes_{\mathcal{A}[\mathcal{E}]} \mathcal{B}$ is the sheaf associated to the presheaf $U \mapsto \bigoplus_{e \in \mathcal{E}(U)} \mathcal{B}(U)$. In other words, Ω is the free $\mathcal{B}$-module on the sheaf of sets $\mathcal{E}$ and in particular there is a canonical map $\mathcal{E} \to \Omega$.

Having said this, pick some $\mathcal{E}$ (for example $\mathcal{E} = \mathcal{B}$ as in the definition of the naive cotangent complex). The obstruction to writing ξ as the class of a map $\mathcal{I}/\mathcal{I}^2 \to \mathcal{G}$ is an element in $\mathrm{Ext}^1_{\mathcal{B}}(\Omega, \mathcal{G})$. Say this is represented by the extension $0 \to \mathcal{G} \to \mathcal{H} \to \Omega \to 0$ of $\mathcal{B}$-modules. Consider the sheaf of sets $\mathcal{E}' = \mathcal{E} \times_{\Omega} \mathcal{H}$ which comes with an induced map $\alpha' : \mathcal{E}' \to \mathcal{B}$. Let $\mathcal{I}' = \mathrm{Ker}(\mathcal{A}[\mathcal{E}'] \to \mathcal{B})$ and $\Omega' = \Omega_{\mathcal{A}[\mathcal{E}']/\mathcal{A}} \otimes_{\mathcal{A}[\mathcal{E}']} \mathcal{B}$. The pullback of ξ under the quasi-isomorphism $NL(\alpha') \to NL(\alpha)$ maps to zero in $\mathrm{Ext}^1_{\mathcal{B}}(\Omega', \mathcal{G})$ because the pullback of the extension $\mathcal{H}$ by the map $\Omega' \to \Omega$ is split as

Date: stacks-project:defos.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

Ω' is the free $\mathcal{B}$-module on the sheaf of sets $\mathcal{E}'$ and since by construction there is a commutative diagram

$$\begin{array}{ccc} \mathcal{E}' & \longrightarrow & \mathcal{E} \\ \downarrow & & \downarrow \\ \mathcal{H} & \longrightarrow & \Omega \end{array}$$

This finishes the proof. $\square$