STACKS, DEFORMATION THEORY: LEMMA-INF-OBS-EXT-RINGED-TOPOI

THE STACKS PROJECT AUTHORS

Editorial note. Statement span is lines 2946-2961 of sources/stacks-defos/pinned-original-source.tex; proof span is lines 2963-3077, the complete author proof (113 body lines).

Editorial note. Navigation stubs: 3 cited-but-not-included labels are supplied as non-load-bearing pointers with their pinned tags where the snapshot tag index has them.

Editorial note. extra_packages is empty on purpose: the retained preamble already loads xy with options and a second declaration clashes in the assembler wrapper.

Editorial note. Text is retained verbatim, including the author's own typos and placeholders.

Editorial note. Queue index 3 of queue_defos.json (pb=3982, stmt 2946-2961, proof 2963-3077).

Allowed standard prerequisites (editorial).

- Explicit prerequisite boundary. The cited inputs are prerequisites; the argument in the retained span is not replaced by any cited result.

Source reference. Original lemma 08MB: sources/stacks-defos/pinned-original-source.tex, source lines 2770–2772. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 08ME: sources/stacks-defos/pinned-original-source.tex, source lines 2912–2922. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 03EW: sources/stacks-defos/pinned-original-source.tex, source lines 3077–3077. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Lemma 0.1. *Let $i : (Sh(\mathcal{C}), \mathcal{O}) \to (Sh(\mathcal{D}), \mathcal{O}')$ be a first order thickening of ringed topoi. Assume given $\mathcal{O}$-modules $\mathcal{F}$, $\mathcal{K}$ and an $\mathcal{O}$-linear map $c : \mathcal{I} \otimes_{\mathcal{O}} \mathcal{F} \to \mathcal{K}$. Then there exists an element*

$$o(\mathcal{F}, \mathcal{K}, c) \in \mathrm{Ext}^2_{\mathcal{O}}(\mathcal{F}, \mathcal{K})$$

whose vanishing is a necessary and sufficient condition for the existence of a sequence (08MB) with $c_{\mathcal{F}'} = c$.

Proof. We first show that if $\mathcal{K}$ is an injective $\mathcal{O}$-module, then there does exist a sequence (08MB) with $c_{\mathcal{F}'} = c$. To do this, choose a flat $\mathcal{O}'$-module $\mathcal{H}'$ and a

Date: stacks-project:defos.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

surjection $\mathcal{H}' \to \mathcal{F}$ (Modules on Sites, Lemma 03EW). Let $\mathcal{J} \subset \mathcal{H}'$ be the kernel. Since $\mathcal{H}'$ is flat we have

$$\mathcal{I} \otimes_{\mathcal{O}'} \mathcal{H}' = \mathcal{I}\mathcal{H}' \subset \mathcal{J} \subset \mathcal{H}'$$

Observe that the map

$$\mathcal{I}\mathcal{H}' = \mathcal{I} \otimes_{\mathcal{O}'} \mathcal{H}' \longrightarrow \mathcal{I} \otimes_{\mathcal{O}'} \mathcal{F} = \mathcal{I} \otimes_{\mathcal{O}} \mathcal{F}$$

annihilates $\mathcal{I}\mathcal{J}$. Namely, if f is a local section of $\mathcal{I}$ and s is a local section of $\mathcal{H}$, then fs is mapped to $f \otimes \overline{s}$ where $\overline{s}$ is the image of s in $\mathcal{F}$. Thus we obtain

$$\begin{array}{ccc} \mathcal{I}\mathcal{H}'/\mathcal{I}\mathcal{J} & \hookrightarrow & \mathcal{J}/\mathcal{I}\mathcal{J} \\ \downarrow & & \downarrow \gamma \\ \mathcal{I} \otimes_{\mathcal{O}} \mathcal{F} & \xrightarrow{c} & \mathcal{K} \end{array}$$

a diagram of $\mathcal{O}$-modules. If $\mathcal{K}$ is injective as an $\mathcal{O}$-module, then we obtain the dotted arrow. Denote $\gamma' : \mathcal{J} \to \mathcal{K}$ the composition of γ with $\mathcal{J} \to \mathcal{J}/\mathcal{I}\mathcal{J}$. A local calculation shows the pushout

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{J} & \longrightarrow & \mathcal{H}' & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle \gamma'} & & \downarrow & & \| & & \\ 0 & \longrightarrow & \mathcal{K} & \longrightarrow & \mathcal{F}' & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \end{array}$$

is a solution to the problem posed by the lemma.

General case. Choose an embedding $\mathcal{K} \subset \mathcal{K}'$ with $\mathcal{K}'$ an injective $\mathcal{O}$-module. Let $\mathcal{Q}$ be the quotient, so that we have an exact sequence

$$0 \to \mathcal{K} \to \mathcal{K}' \to \mathcal{Q} \to 0$$

Denote $c' : \mathcal{I} \otimes_{\mathcal{O}} \mathcal{F} \to \mathcal{K}'$ be the composition. By the paragraph above there exists a sequence

$$0 \to \mathcal{K}' \to \mathcal{E}' \to \mathcal{F} \to 0$$

as in (08MB) with $c_{\mathcal{E}'} = c'$. Note that c' composed with the map $\mathcal{K}' \to \mathcal{Q}$ is zero, hence the pushout of $\mathcal{E}'$ by $\mathcal{K}' \to \mathcal{Q}$ is an extension

$$0 \to \mathcal{Q} \to \mathcal{D}' \to \mathcal{F} \to 0$$

as in (08MB) with $c_{\mathcal{D}'} = 0$. This means exactly that $\mathcal{D}'$ is annihilated by $\mathcal{I}$, in other words, the $\mathcal{D}'$ is an extension of $\mathcal{O}$-modules, i.e., defines an element

$$o(\mathcal{F}, \mathcal{K}, c) \in \mathrm{Ext}^1_{\mathcal{O}}(\mathcal{F}, \mathcal{Q}) = \mathrm{Ext}^2_{\mathcal{O}}(\mathcal{F}, \mathcal{K})$$

(the equality holds by the long exact cohomology sequence associated to the exact sequence above and the vanishing of higher ext groups into the injective module $\mathcal{K}'$). If $o(\mathcal{F}, \mathcal{K}, c) = 0$, then we can choose a splitting $s : \mathcal{F} \to \mathcal{D}'$ and we can set

$$\mathcal{F}' = \mathrm{Ker}(\mathcal{E}' \to \mathcal{D}'/s(\mathcal{F}))$$

so that we obtain the following diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{K} & \longrightarrow & \mathcal{F}' & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \| & & \\ 0 & \longrightarrow & \mathcal{K}' & \longrightarrow & \mathcal{E}' & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \end{array}$$

with exact rows which shows that $c_{\mathcal{F}'} = c$. Conversely, if $\mathcal{F}'$ exists, then the pushout of $\mathcal{F}'$ by the map $\mathcal{K} \to \mathcal{K}'$ is isomorphic to $\mathcal{E}'$ by Lemma 08ME and the vanishing of higher ext groups into the injective module $\mathcal{K}'$. This gives a diagram as above, which implies that $\mathcal{D}'$ is split as an extension, i.e., the class $o(\mathcal{F}, \mathcal{K}, c)$ is zero. $\square$