STACKS, DEFORMATION THEORY: LEMMA-INF-OBS-MAP-SPECIAL-RINGED-TOPOI

THE STACKS PROJECT AUTHORS

Editorial note. Statement span is lines 3741-3756 of sources/stacks-defos/pinned-original-source.tex; proof span is lines 3758-3774, the complete author proof (15 body lines).

Editorial note. Navigation stubs: 2 cited-but-not-included labels are supplied as non-load-bearing pointers with their pinned tags where the snapshot tag index has them.

Editorial note. extra_packages is empty on purpose: the retained preamble already loads xy with options and a second declaration clashes in the assembler wrapper.

Editorial note. Text is retained verbatim, including the author's own typos and placeholders.

Editorial note. Queue index 40 of queue_defos.json (pb=670, stmt 3741-3756, proof 3758-3774).

Allowed standard prerequisites (editorial).

- Explicit prerequisite boundary. The cited inputs are prerequisites; the argument in the retained span is not replaced by any cited result.

Source reference. Original lemma 08MP: sources/stacks-defos/pinned-original-source.tex, source lines 3584–3594. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 07A6: sources/stacks-defos/pinned-original-source.tex, source lines 3774–3774. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Lemma 0.1. *Let $i : (Sh(\mathcal{C}), \mathcal{O}) \to (Sh(\mathcal{D}), \mathcal{O}')$ be a first order thickening of ringed topoi. Let $\mathcal{F}'$, $\mathcal{G}'$ be $\mathcal{O}'$-modules and set $\mathcal{F} = i^*\mathcal{F}'$ and $\mathcal{G} = i^*\mathcal{G}'$. Let $\varphi : \mathcal{F} \to \mathcal{G}$ be an $\mathcal{O}$-linear map. There exists an element*

$$o(\varphi) \in \mathrm{Ext}^1_{\mathcal{O}}(Li^*\mathcal{F}', \mathcal{I}\mathcal{G}')$$

whose vanishing is a necessary and sufficient condition for the existence of a lift of φ to an $\mathcal{O}'$-linear map $\varphi' : \mathcal{F}' \to \mathcal{G}'$.

Proof. It is clear from the proof of Lemma 08MP that the vanishing of the boundary of φ via the map

$$\mathrm{Hom}_{\mathcal{O}}(\mathcal{F}, \mathcal{G}) = \mathrm{Hom}_{\mathcal{O}'}(\mathcal{F}', \mathcal{G}) \longrightarrow \mathrm{Ext}^1_{\mathcal{O}'}(\mathcal{F}', \mathcal{I}\mathcal{G}')$$

is a necessary and sufficient condition for the existence of a lift. We conclude as

$$\mathrm{Ext}^1_{\mathcal{O}'}(\mathcal{F}', \mathcal{I}\mathcal{G}') = \mathrm{Ext}^1_{\mathcal{O}}(Li^*\mathcal{F}', \mathcal{I}\mathcal{G}')$$

the adjointness of $i_* = Ri_*$ and Li^* on the derived category (Cohomology on Sites, Lemma 07A6). □

Date: stacks-project:defos.tex:local-snapshot-20261012; GNU FDL 1.2 or later.