STACKS, EXAMPLES OF DEFORMATIONS: LEMMA-FIRST-ORDER-HENSELIZATION

THE STACKS PROJECT AUTHORS

Editorial note. Statement span is lines 2773-2785 of sources/stacks-examples-defos/pinned-original-source.tex; proof span is lines 2787-2832, the complete author proof (44 body lines).

Editorial note. Navigation stubs: 8 cited-but-not-included labels are supplied as non-load-bearing pointers with their pinned tags where the snapshot tag index has them.

Editorial note. extra_packages is empty on purpose: the retained preamble already loads xy with options and a second declaration clashes in the assembler wrapper.

Editorial note. Text is retained verbatim, including the author's own typos and placeholders.

Editorial note. Queue index 10 of queue_examples-defos.json (pb=1885, stmt 2773-2785, proof 2787-2832).

Allowed standard prerequisites (editorial).

- Explicit prerequisite boundary. The cited inputs are prerequisites; the argument in the retained span is not replaced by any cited result.

Source reference. Original lemma 00S1: sources/stacks-examples-defos/pinned-original-source.tex, source lines 2832–2832. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 00S0: sources/stacks-examples-defos/pinned-original-source.tex, source lines 2832–2832. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 0DZ4: sources/stacks-examples-defos/pinned-original-source.tex, source lines 2383–2393. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 0DYT: sources/stacks-examples-defos/pinned-original-source.tex, source lines 1251–1260. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 0A6A: sources/stacks-examples-defos/pinned-original-source.tex, source lines 2832–2832. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 0D08: sources/stacks-examples-defos/pinned-original-source.tex, source lines 2832–2832. Cited by the retained statement or

Date: stacks-project:examples-defos.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 0AGU: sources/stacks-examples-defos/pinned-original-source.tex, source lines 2832–2832. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 05EC: sources/stacks-examples-defos/pinned-original-source.tex, source lines 2832–2832. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Lemma 0.1. *Let k be a field. Let B be a finite type k-algebra. Let $J \subset B$ be an ideal such that $\mathrm{Spec}(B) \to \mathrm{Spec}(k)$ is smooth on the complement of $V(J)$. Let N be a finite B-module. Then there is a canonical bijection*

$$Exal_k(B, N) \to Exal_k(B^h, N^h)$$

Here (B^h, J^h) is the henselization of (B, J) and $N^h = N \otimes_B B^h$.

Proof. This proof is the same as the proof of Lemma 0DZ4 but easier. We suggest the reader to skip the proof. The map is given by henselization: given $0 \to N \to C \to B \to 0$ in $\mathrm{Exal}_k(B, N)$ we send it to the henselization C^h of C with respect to the inverse image $J_C \subset C$ of J. Compare with the proof of Lemma 0DYT.

Since $k \to B$ is of finite presentation the complex $NL_{B/k}$ can be represented by a complex $N^{-1} \to N^0$ where N^i is a finite B-module, see Algebra, Section 00S0 and in particular Algebra, Lemma 00S1. As B is Noetherian, this means that $NL_{B/k}$ is pseudo-coherent. For $g \in J$ the k-algebra B_g is smooth and hence $(NL_{B/k})_g = NL_{B_g/k}$ is quasi-isomorphic to a finite projective B-module sitting in degree 0. Thus $\mathrm{Ext}^i_B(NL_{B/k}, N)_g = 0$ for $i \geq 1$ and any B-module N. Finally, we have

$$\begin{aligned} \mathrm{Ext}^1_{B^h}(NL_{B^h/k}, N^h) &= \mathrm{Ext}^1_{B^h}(NL_{B/k} \otimes_B B^h, N \otimes_B B^h) \\ &= \mathrm{Ext}^1_B(NL_{B/k}, N) \otimes_B B^h \\ &= \mathrm{Ext}^1_B(NL_{B/k}, N) \end{aligned}$$

The first equality by More on Algebra, Lemma 0D08 (or rather its analogue for henselizations of pairs). The second by More on Algebra, Lemma 0A6A. The third because $\mathrm{Ext}^1_B(NL_{B/k}, N)$ is J-power torsion, the map $B \to B^h$ is flat and induces an isomorphism $B/J \to B^h/JB^h$ (More on Algebra, Lemma 0AGU), and More on Algebra, Lemma 05EC. This concludes the proof by the description of $\mathrm{Exal}_A(B, N)$ as $\mathrm{Ext}^1_B(NL_{B/A}, N)$ given just above Lemma 0DZ4. □