THE RELATIVE MOR FUNCTOR IS AN FPPF SHEAF AND REPRESENTS PAIRS

THE STACKS PROJECT AUTHORS

Editorial note. Closing sentence 'We omit the verification that the two constructions above are mutually inverse' retained verbatim and flagged for the reviewer (routine bookkeeping).

Allowed standard prerequisites (editorial).

- Universal effective epimorphisms. spaces-descent-lemma-fpqc-universal-effective-epimorphisms (cross-chapter, named). Descends both components along the covering, twice.
- Space presentation. spaces-lemma-space-presentation (cross-chapter, named). Identifies T as the fppf quotient sheaf U/R in the converse direction.

Source reference. Original reference 001P: focused-seeds/geometry/S9-crit-hom-functor-sheaf/ref-tags.txt, source lines 5–5. Navigation pointer only: the counted proof cites this statement by \ref as a standing input; the deduction the counted proof has to make is carried out inside the counted span, and no part of that deduction is delegated to this pointer.

Source reference. Original equation 05Y1: sources/stacks-criteria/pinned-original-source.tex, source lines 1488–1488. Navigation pointer only: the counted proof cites this statement by \ref as a standing input; the deduction the counted proof has to make is carried out inside the counted span, and no part of that deduction is delegated to this pointer.

Source reference. Original reference 04P2: focused-seeds/geometry/S9-crit-hom-functor-sheaf/ref-tags.txt, source lines 6–6. Navigation pointer only: the counted proof cites this statement by \ref as a standing input; the deduction the counted proof has to make is carried out inside the counted span, and no part of that deduction is delegated to this pointer.

Source reference. Original reference 0262: focused-seeds/geometry/S9-crit-hom-functor-sheaf/ref-tags.txt, source lines 7–7. Navigation pointer only: the counted proof cites this statement by \ref as a standing input; the deduction the counted proof has to make is carried out inside the counted span, and no part of that deduction is delegated to this pointer.

Source reference. Original reference 03Y7: focused-seeds/geometry/S9-crit-hom-functor-sheaf/ref-tags.txt, source lines 8–8. Navigation pointer only: the counted proof cites this statement by \ref as a standing input; the deduction the counted proof has to make is carried out inside the counted span, and no part of that deduction is delegated to this pointer.

Date: stacks-project:criteria.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

Lemma 0.1. *Let S be a scheme. Let $Z \to B$ and $X \to B$ be morphisms of algebraic spaces over S. Then*

(1) *$Mor_B(Z, X)$ is a sheaf on $(Sch/S)_{fppf}$.*
(2) *If T is an algebraic space over S, then there is a canonical bijection*

$$\mathrm{Mor}_{Sh((Sch/S)_{fppf})}(T, Mor_B(Z, X)) = \{(a, b) \text{ as in } (05Y1)\}$$

Proof. Let T be an algebraic space over S. Let $\{T_i \to T\}$ be an fppf covering of T (as in Topologies on Spaces, Section 03Y7). Suppose that $(a_i, b_i) \in Mor_B(Z, X)(T_i)$ such that $(a_i, b_i)|_{T_i \times_T T_j} = (a_j, b_j)|_{T_i \times_T T_j}$ for all i, j. Then by Descent on Spaces, Lemma 04P2 there exists a unique morphism $a : T \to B$ such that a_i is the composition of $T_i \to T$ and a. Then $\{T_i \times_{a_i, B} Z \to T \times_{a,B} Z\}$ is an fppf covering too and the same lemma implies there exists a unique morphism $b : T \times_{a,B} Z \to T \times_{a,B} X$ such that b_i is the composition of $T_i \times_{a_i, B} Z \to T \times_{a,B} Z$ and b. Hence $(a, b) \in Mor_B(Z, X)(T)$ restricts to (a_i, b_i) over T_i for all i.

Note that the result of the preceding paragraph in particular implies (1).

Let T be an algebraic space over S. In order to prove (2) we will construct mutually inverse maps between the displayed sets. In the following when we say "pair" we mean a pair (a, b) fitting into (05Y1).

Let $v : T \to Mor_B(Z, X)$ be a natural transformation. Choose a scheme U and a surjective étale morphism $p : U \to T$. Then $v(p) \in Mor_B(Z, X)(U)$ corresponds to a pair (a_U, b_U) over U. Let $R = U \times_T U$ with projections $t, s : R \to U$. As v is a transformation of functors we see that the pullbacks of (a_U, b_U) by s and t agree. Hence, since $\{U \to T\}$ is an fppf covering, we may apply the result of the first paragraph that deduce that there exists a unique pair (a, b) over T.

Conversely, let (a, b) be a pair over T. Let $U \to T$, $R = U \times_T U$, and $t, s : R \to U$ be as above. Then the restriction $(a, b)|_U$ gives rise to a transformation of functors $v : h_U \to Mor_B(Z, X)$ by the Yoneda lemma (Categories, Lemma 001P). As the two pullbacks $s^*(a, b)|_U$ and $t^*(a, b)|_U$ are equal, we see that v coequalizes the two maps $h_t, h_s : h_R \to h_U$. Since $T = U/R$ is the fppf quotient sheaf by Spaces, Lemma 0262 and since $Mor_B(Z, X)$ is an fppf sheaf by (1) we conclude that v factors through a map $T \to Mor_B(Z, X)$.

We omit the verification that the two constructions above are mutually inverse. □