An explicit logarithmic list-chromatic lower bound for fixed intersection graphs G(n,r,s)

M. M. Koshelev

arxiv:2610.13723v1; CC BY 4.0

Editorial note. Complete original author mathematical body and supporting proofs retained; one independent main conclusion per task. No proof repair, added hypotheses or independent correctness certification.

Editorial note. Mechanical source0.tex adapter uses T1 instead of unavailable T2A for this English-only body, omits the unused unavailable systeme package, and replaces native Biber setup with installed BibTeX unsrt using original reference fields. Raw main.tex and KoshelevBib.bib retained; only unescaped bibliography math expressions receive title-case protection.

Editorial note. Mechanical bibliography comment adapter omits native percent-comment-only lines from references-bibtex.bib, including those inside entries. All active original fields and their values are retained; no missing journal, year or other source metadata is filled. Raw KoshelevBib.bib unchanged.

Editorial note. Mechanical title-case protection braces the original ShKup title so BibTeX cannot lowercase the native Hungarian H-accent command into undefined h. Original title wording, all active fields and raw bibliography unchanged; no journal or year supplied.

Editorial note. Collection note: the original author text says Corollary with unresolved label coroll. This wording and unresolved locator remain in source0.tex. Only the generated rendering replaces the unresolved reference with [unresolved original author label: coroll]. No intended target is assigned; the original mainLKG statement and proof remain in context without being identified as that target.

Allowed standard prerequisites (editorial).

- List coloring, intersection graphs, supersaturation and fractional colorings. All native G(n,r,s), Kneser and fractional-coloring definitions, quantitative parameter restrictions and auxiliary counting lemmas retained.. The selected random-list lower constructions and primary/reserve upper construction remain counted in full; asymptotic statements do not erase their original parameter restrictions.

1 Introduction

In this paper we study list colorings of the graphs $G(n, r, s)$. For integers $n > r > s \geq 0$, let $[n] = \{1, \dots, n\}$ and let $\binom{[n]}{r}$ denote the family of all r-element subsets of $[n]$. The graph $G(n, r, s)$ has vertex set $\binom{[n]}{r}$, with two distinct vertices A and B adjacent if and only if $|A \cap B| = s$. This family includes the Kneser graphs $KG(n, r) = G(n, r, 0)$ and the Johnson graphs $J(n, r) = G(n, r, r-1)$. Independent sets in $G(n, r, s)$ are precisely uniform set systems with one forbidden intersection size, which connects these graphs to extremal set theory [1]. Their realizations as Euclidean distance graphs also make them useful in combinatorial geometry [2]. Other applications of these graphs include coding and Ramsey theory (see [3, 4, 5]). A lot of literature studies various properties of these graphs (see, for example, [6, 7, 8, 9]).

A substantial part of this research concerns coloring parameters. For example, the classical result of Lovász [10] determines the chromatic number for Kneser graphs.

Theorem 1. *Let $n \geq 2r$. Then*

$$\chi(G(n, r, 0)) = n - 2r + 2.$$

For general r and s, chromatic numbers have been investigated using methods from extremal set theory, algebra, and design theory; see, for example, [11, 2, 12]. Other coloring parameters have also attracted attention. In particular, the clique-chromatic number is the minimum number of colors in a vertex coloring with no monochromatic inclusion-maximal clique of size at least two. Bounds on this parameter for several families of $G(n, r, s)$ graphs were obtained in [13, 14, 15].

Here we focus on the *list chromatic number*, also known as the *choice number*. List coloring was introduced independently by Vizing [16] and Erdős, Rubin, and Taylor [17]. A *list assignment* L associates a set $L(v)$ of available colors with each vertex v of a graph G. An *L-coloring* is a proper vertex coloring c such that $c(v) \in L(v)$ for every $v \in V(G)$. The list chromatic number $\chi_L(G)$ is the least integer k such that G admits an L-coloring for every list assignment satisfying $|L(v)| \geq k$ for all $v \in V(G)$.

Note that the following proposition trivially follows from the definition.

Proposition 2.

$$\chi(G) \le \chi_L(G) \le \Delta(G) + 1,$$

where $\Delta(G)$ denotes the maximum degree of G.

List chromatic number was studied for many families of graphs including complete multipartite graphs (see [18, 19]), planar graphs (see [20]), line graphs of bipartite graphs (see [21]) and pseudorandom graphs (see [22]).

Unfortunately, little is known about the behaviour of list chromatic number of $G(n, r, s)$ graphs. For positive s we in fact are aware of only one result due to Zakharov [12].

Theorem 3. *For fixed integers $r > s \ge 0$ we have*

$$\chi_L(G(n, r, s)) = O(n^{s+1} \ln n).$$

Together with Proposition 2, this gives

$$\chi_L(G(n, r, s)) = \Theta(n^{r-s}) \qquad \text{for } r \le 2s + 1,$$

and

$$c_{r,s} n^{s+1} \le \chi_L(G(n, r, s)) \le C_{r,s} n^{s+1} \ln n \qquad \text{for } r > 2s + 1.$$

Here $c_{r,s}$ and $C_{r,s}$ are positive constants.

The Kneser case is a little bit more studied. Bulankina and Kupavskii [23] proved the following two theorems.

Theorem 4. *For every fixed $\varepsilon \in (0, 1/2)$ and any $2 \le r \le n^{1/2-\varepsilon}$ we have*

$$\chi_L(G(n, r, 0)) \ge C_\varepsilon n \ln n$$

for some $C_\varepsilon > 0$.

Theorem 5. *For any r and n such that $2r \le n$ we have*

$$\chi_L(G(n, r, 0)) \le n \ln \frac{n}{r} + n.$$

Inozemtsev and Kupavskii [24] subsequently strengthened the lower bound. To be more precise, they obtained the following theorem.

Theorem 6. *Fix $\varepsilon > 0$ and let $3 \le r \le \sqrt{\frac{n}{1+\varepsilon}}$. Then*

$$\chi_L(G(n, r, 0)) \ge \left(\frac{r-2}{8r} - \frac{\varepsilon}{2} \right) n \ln n$$

for all $n > n_0(\varepsilon)$.

In this paper we address the following two questions. The first one is the correct growth rate of $\chi_L(G(n, r, s))$ for $r > 2s + 1$. We give an answer to this question in a slightly more restraining regime $r \ge 2s + 3$.

Theorem 7. *Fix integers $s > 0$ and $r \ge 2s + 3$. Then*

$$\chi_L(G(n, r, s)) \ge (1 - o(1)) \frac{r - 2s - 1}{8 \ln 2 \binom{r}{s+1}} \binom{n}{s+1} \ln n.$$

Furthermore we give an upper bound of the same order of magnitude with explicit constant. Note that the result about the correct order of magnitude is not new, since Theorem 3 already settles this question. Our results, however, have two major advantages. First, these results can be applied to almost any graph and second, the explicit constant in our results is smaller than the implicit constant in Theorem 3.

We give two bounds. The second one is stronger but at the same time a bit less general.

Before we go on to the results, let us give a definition of fractional chromatic number.

Definition 1.1. Let $\mathcal{Q}$ be a set of all possible distributions on the collection of independent sets in graph G. Define

$$\chi_f(G) := \min_{\mathbf{Q} \in \mathcal{Q}} \max_{v \in G} \mathbf{Q}^{-1}(v \in I),$$

where I is a random independent set. We call $\chi_f(G)$ the fractional chromatic number of G.

A lot of properties of fractional chromatic number are known. For example, it is easy to prove that for vertex transitive graph G we have

$$\chi_f(G) = \frac{|V_G|}{\alpha(G)}.$$

For fixed integers $s \ge 0$ and $r > 2s + 1$, and for sufficiently large n, we have

$$\chi_f(G(n, r, s)) = \frac{1}{\binom{r}{s+1}} \binom{n}{s+1}.$$

A simple probabilistic argument allows us to prove the following theorem.

Theorem 8. *For any graph G on n vertices we have*

$$\chi_L(G) \le k = \lceil \chi_f(G)(\ln n + 1) \rceil.$$

Note that this theorem alone gives us something like

$$\chi_L(G(n,r,s)) \le (1+o(1))\frac{r}{\binom{r}{s+1}}\binom{n}{s+1}\ln n.$$

The second result is a bit more involved.

Theorem 9. *Let G be a graph on $n \ge 2$ vertices. Let $a(G) = n\chi_f^{-1}(G)$ (note that $a(G) = \alpha(G)$ for vertex transitive graphs). Assume that $\frac{\chi_f(G)}{\sqrt{\ln a(G)}} > 10 \ln n$ and $a(G) \ge 3$. Then*

$$\chi_L(G) \le k = \lceil \chi_f(G)\left(\ln a(G) + \ln\ln a(G)\right) \rceil + 3\left\lceil \frac{\chi_f(G)}{\sqrt{\ln a(G)}} \right\rceil$$

For fixed integers $s \ge 0$ and $r > 2s+1$, all conditions of this theorem are satisfied for $G(n,r,s)$ when n is sufficiently large. We can apply it and obtain a slightly better bound.

Corollary 1.1. *Fix r and s such that $r > 2s+1$ and $s \ge 0$. We have*

$$\chi_L(G(n,r,s)) \le (1+o(1))\frac{r-s-1}{\binom{r}{s+1}}\binom{n}{s+1}\ln n.$$

The second question we address is the correct growth rate of $\chi_L(G(n,r,0))$ for $r = \Omega(n^{1/2})$. To this end we prove the following theorem.

Theorem 10. *Let $n \ge 1000$, $r \ge 4\ln n$, $20r \le n$. Then*

$$\chi_L(G(n,r,0)) \ge \frac{n}{120}\ln\left(\frac{en}{r}\right).$$

Note that the combination of Theorems 4, 5, 10 and 1 together with Proposition 2 shows that

$$\chi_L(G(n,r,0)) = \Theta\left(n\ln(en/r)\right)$$

uniformly for $2 \le r \le (1/2-\varepsilon)n$.

The rest of the paper is structured as follows. In section 2 we formulate auxiliary lemmas and theorems. In section 3 we prove these auxiliary results. In section 4 we prove the lower bounds for both $r \ge 2s+3$ and $s = 0$ cases. Finally in section 5 we prove upper bounds on $\chi_L(G)$.

2 Auxiliary results

In our proofs we need a supersaturation result on the number of edges inside a set of vertices. For the case $r \ge 2s+3$ we use the following result that is a compilation of two theorems from [25].

Theorem 11. *Fix two integers, r and s, $s > 0$ and $r > 2s+2$. For any $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon,r,s) > 0$ such that for all $n \ge n_0(\varepsilon,r,s)$ and any $A \subset G$ such that $|A| > (1+\varepsilon)\binom{n-s-1}{r-s-1}$ we have $e(A) > \frac{\delta|A|^2}{n^s}$.*

For the $s = 0$ case we prove the following auxiliary theorem.

Theorem 12. *Fix r, n such that $2r \le n$. Let A be the set of vertices of $G(n,r,0)$ such that $|A| > \alpha(G(n,r,0)) = \binom{n-1}{r-1}$. Then*

$$e(A) \ge \frac{\binom{n-r-1}{r-1}}{2\binom{n-1}{r-1}}|A|\left(|A| - \binom{n-1}{r-1}\right).$$

The other ingredient we need is the result on containers in $G(n,r,s)$. The notion of the container was first introduced by Saxton and Thomason and independently by Balogh, Morris and Samotij (see [26, 27]). Before we proceed, let us give the definition. Note that here we a slightly different definition that is more convenient for us.

Definition 2.1. Consider a graph G. Let $\mathcal{C}$ be a collection of subsets of its vertices satisfying the following properties.

1. Each independent subset of G is contained in at least one of the sets $C \in \mathcal{C}$.
2. For each set $C \in \mathcal{C}$ its size does not exceed $K\alpha(G)$.
3. The cardinality $|\mathcal{C}|$ does not exceed f.

We call such a collection (K,f)-container.

For example, $\mathcal{C} = \{G\}$ is a $(|G|/\alpha(G), 1)$-container and the set of all subsets of size $\alpha(G)$ is a $\left(1, \binom{n}{\alpha(G)}\right)$-container.

We again need two auxiliary results here, one for the $r \geq 2s+3$ case and the second for $s = 0$ case.

Lemma 2.1. *Fix $s > 0$, $r \geq 2s+2$ and $\varepsilon > 0$. There exists $\delta = \delta_{\varepsilon,r,s} > 0$ such that there exists a $(1+\varepsilon, \exp\left(\delta n^s \ln^2 n\right))$-container in $G(n,r,s)$.*

Lemma 2.2. *Fix r, n such that $r \geq 10$, $n \geq 1000$, $3r \leq n$ and take any $1 \geq \varepsilon > \frac{1}{n}$. Then there exists a $\left(1+\varepsilon, 2\left(\frac{e\binom{n}{r}}{k}\right)^k\right)$-container in $G(n,r,0)$, where*

$$k = \left\lfloor \frac{2(2+\varepsilon)\binom{n-1}{r-1}\ln\binom{n}{r}}{\varepsilon\binom{n-r-1}{r-1}} \right\rfloor.$$

For instance, there exists a $\left(2, 2\left(\frac{e\binom{n}{r}}{k}\right)^k\right)$-container in $G(n,r,0)$, where

$$k = \left\lfloor \frac{6\binom{n-1}{r-1}\ln\binom{n}{r}}{\binom{n-r-1}{r-1}} \right\rfloor.$$

3 Proof of auxiliary results

3.1 Proof of Theorem 12

Proof. Recall that $G(n,r,0)$ is an (N,d,λ) graph with $N = \binom{n}{r}$, $d = \binom{n-r}{r}$ and $\lambda = \binom{n-r-1}{r-1}$. Note that $d = \frac{n-r}{r}\lambda$, $N = \frac{n}{r}\alpha$.

We thus have

$$2e(A) \geq \frac{d}{N}|A|^2 - \frac{\lambda}{N}|A|(N-|A|) = \frac{d+\lambda}{N}|A|^2 - \lambda|A| = \frac{\lambda}{\alpha}\left(\frac{\frac{n-r}{r}+1}{\frac{n}{r}}|A|^2 - |A|\alpha\right) = \frac{\lambda}{\alpha}|A|(|A|-\alpha),$$

which is exactly what we need. ■

3.2 Proof of Lemma 2.1

Proof of the lemma. Let us consider an arbitrary independent set I in $G(n,r,s)$ and perform the following algorithm. Put $V_0 = V_{G(n,r,s)}$ and $U_0 = \varnothing$. At i-th step select the vertex v_i with maximum degree in $G(n,r,s)[V_{i-1}]$. If $v_i \in I$, put $V_i = V_{i-1} \setminus (\{v_i\} \cup N(v_i))$ and $U_i = U_{i-1} \cup \{v_i\}$. If $v_i \notin I$, we put $V_i = V_{i-1} \setminus \{v_i\}$ and $U_i = U_{i-1}$. We stop whenever $|V_i| \leq (1+\varepsilon/2)\binom{n-s-1}{r-s-1}$. In the remaining part of the proof index i denotes the last step of the algorithm (that is, $|V_i| \leq (1+\varepsilon/2)\binom{n-s-1}{r-s-1}$).

Let us obtain some properties of V_i and U_i. First of all, let us notice that V_i is a function of U_i. Indeed, if two independent sets I_1 and I_2 are the same on $V_{G(n,r,s)} \setminus V_i$, the algorithm does just the same steps for them and thus the final sets will be the same. This means that the number of different $V_i \cup U_i$ over all I is exactly the same as the number of different U_i.

The second thing to notice is that whenever we add a vertex to U_{j+1}, we throw out $\Delta(G(n,r,s)[V_j]) + 1$ vertices from V_j. This can be trivially bounded from below with the use of Theorem 11:

$$\Delta(G(n,r,s)[V_j]) \geq \frac{2e(G(n,r,s)[V_j])}{|V_j|} \geq \frac{2\delta|V_j|}{n^s}.$$

We thus have

$$|V_{j+1}| \leq \left(1 - \frac{C}{n^s}\right)|V_j| \Rightarrow 1 \leq |V_{i-1}| \leq \left(1 - \frac{C}{n^s}\right)^{|U_i|-1}\binom{n}{r} \Rightarrow |U_i| \leq \tilde{C}n^s \ln n$$

for some $\tilde{C} = \tilde{C}(r,s,\varepsilon)$.

Finally, the last thing to notice is that $V_i \cup U_i$ contains I.

Using these three properties we can see that the collection of all possible $V_i \cup U_i$ is indeed a container, each set $V_i \cup U_i$ is of size at most $(1+\varepsilon/2+o_n(1))\binom{n-s-1}{r-s-1} \leq (1+\varepsilon)\binom{n-s-1}{r-s-1}$ and there are at most

$$\sum_{j=0}^{\lfloor \tilde{C}n^s \ln n \rfloor} \binom{\binom{n}{r}}{j} = \exp\left(O(n^s \ln^2 n)\right)$$

such sets, which is exactly what we need. ■

3.3 Proof of Lemma 2.2

Proof. Let us consider an arbitrary independent set I in $G(n,r,0)$ and perform the following algorithm. Put $V_0 = V_{G(n,r,0)}$ and $U_0 = \varnothing$. At i-th step select the vertex v_i with maximum degree in $G(n,r,0)[V_{i-1}]$. If $v_i \in I$, put $V_i = V_{i-1} \setminus (\{v_i\} \cup N(v_i))$ and $U_i = U_{i-1} \cup \{v_i\}$. If $v_i \notin I$, we put $V_i = V_{i-1} \setminus \{v_i\}$ and $U_i = U_{i-1}$. We stop whenever $|V_i| \le (1+\varepsilon/2)\binom{n-1}{r-1}$. In the remaining part of the proof index i denotes the last step of the algorithm (that is, $|V_i| \le (1+\varepsilon/2)\binom{n-1}{r-1}$).

Let us obtain some properties of V_i and U_i. First of all, let us notice that V_i is a function of U_i. Indeed, if two independent sets I_1 and I_2 are the same on $V_{G(n,r,0)} \setminus V_i$, the algorithm does just the same steps for them and thus the final sets will be the same. This means that the number of different $V_i \cup U_i$ over all I is exactly the same as the number of different U_i.

The second thing to notice is that whenever we add a vertex to U_{j+1}, we throw out $\Delta(G(n,r,0)[V_j]) + 1$ vertices from V_j. This can be trivially bounded from below with the use of Theorem 12:

$$\Delta(G(n,r,0)[V_j]) \ge \frac{2e(G(n,r,0)[V_j])}{|V_j|} \ge \binom{n-r-1}{r-1} \frac{|V_j| - \binom{n-1}{r-1}}{\binom{n-1}{r-1}} \ge \frac{\varepsilon/2}{1+\varepsilon/2} \frac{\binom{n-r-1}{r-1}}{\binom{n-1}{r-1}} |V_j|.$$

We thus have

$$|V_{j+1}| \le \left(1 - \frac{\varepsilon}{2+\varepsilon} \frac{\binom{n-r-1}{r-1}}{\binom{n-1}{r-1}}\right) |V_j| \Rightarrow$$

$$\Rightarrow 1 \le |V_{i-1}| \le \left(1 - \frac{\varepsilon}{2+\varepsilon} \frac{\binom{n-r-1}{r-1}}{\binom{n-1}{r-1}}\right)^{|U_i|-1} \binom{n}{r} \le$$

$$\le \exp\left\{-\frac{\varepsilon}{2+\varepsilon} \frac{\binom{n-r-1}{r-1}}{\binom{n-1}{r-1}} (|U_i| - 1)\right\} \binom{n}{r} \Rightarrow$$

$$\Rightarrow |U_i| \le 1 + \frac{(2+\varepsilon)\binom{n-1}{r-1} \ln \binom{n}{r}}{\varepsilon \binom{n-r-1}{r-1}} \le \frac{2(2+\varepsilon)\binom{n-1}{r-1} \ln \binom{n}{r}}{\varepsilon \binom{n-r-1}{r-1}}.$$

Note that since $n \ge 3r$, we have

$$\frac{2(2+\varepsilon)\binom{n-1}{r-1} \ln \binom{n}{r}}{\varepsilon \binom{n-r-1}{r-1}} \le 2\frac{3^{10}9!\binom{n-1}{r-1}}{\varepsilon n^8} \le 2\frac{3^{10}9!}{n^7}\binom{n-1}{r-1} \le \frac{100}{n^2}\binom{n-1}{r-1} \le \frac{1}{10n}\binom{n-1}{r-1}$$

and thus $|U_i \cup V_i| \le \left(1 + \varepsilon/2 + \frac{1}{10n}\right)\binom{n-1}{r-1} \le (1+\varepsilon)\binom{n-1}{r-1}$.

Finally, the last thing to notice is that $V_i \cup U_i$ contains I.

Now we put all of this together to obtain that the set of all possible $U_i \cup V_i$ is indeed a container, each $V_i \cup U_i$ has size at most $(1+\varepsilon)\binom{n-1}{r-1}$ and the number of distinct $V_i \cup U_i$ is at most

$$S = \sum_{i=0}^{k} \binom{\binom{n}{r}}{i}.$$

Since $k \le \binom{n-1}{r-1} = \frac{r}{n}\binom{n}{r} \le \frac{1}{3}\binom{n}{r}$, we have

$$\binom{\binom{n}{r}}{i}\binom{\binom{n}{r}}{i+1}^{-1} = \frac{(i+1)}{\binom{n}{r} - i} \le \frac{1}{2}$$

for $i < k$ and thus

$$S \le 2\binom{\binom{n}{r}}{k} \le 2\left(\frac{\binom{n}{r}e}{k}\right)^k.$$

■

4 Lower bounds

4.1 Proof of Theorem 7

Proof. Fix any $0 < \varepsilon < 1$ and let us take $\delta = (1-\varepsilon)\frac{r-2s-1}{8\ln 2\binom{r}{s+1}}$. The main idea is the following. We take $2q = 2\left\lfloor \delta\binom{n}{s+1}\ln n \right\rfloor$ colors and randomly assign each vertex a list containing half of these colors. Assume that we managed to color our graph properly. Let B_i be the set of vertices of color i. Then B_i is contained in some set from (any) container. This means that we can assign each color c a set $\varphi(c)$ from the container such that for each i and each $v \in B_i$ we have at least one C from the container such that $v \in C$ and $\varphi(i) = C$. It remains to prove that with positive probability for any φ there exists at least one such v that $v \notin \bigcup_{j=1}^{q} \varphi(c_j)$, where $\{c_1, \dots, c_q\}$ is the list of colors assigned to v. We call φ with this condition bad and all other φ good.

Let us now give a rigorous proof. Consider the $(2, \exp\{Cn^s \ln^2 n\})$-container $\mathcal{C}$ from Lemma 2.1. Let us fix some function φ that maps each color to some set $C \in \mathcal{C}$. Let us bound the probability that φ is good. Let $\mathcal{C}(v)$ be the collection of all

sets in $\mathcal{C}$ containing v. Let also list(v) be the list of colors for v. Finally, put $\xi_v = \mathbb{1}\left\{\text{list}(v) \cap \varphi^{-1}(\mathcal{C}(v)) = \varnothing\right\}$. Then φ is good if and only if $\sum_{i=1}^{\binom{n}{r}} \xi_i = 0$. Since all ξ_i are independent, we have

$$\mathbf{P}\left(\sum_{i=1}^{\binom{n}{r}} \xi_i = 0\right) = \prod_{i=1}^{\binom{n}{r}} \mathbf{P}(\xi_i = 0).$$

Let us now bound $\mathbf{P}(\xi_v = 0)$. In fact we will not bound all such probabilities, but only the ones with small $\varphi^{-1}(\mathcal{C}(v))$. Note that since each $C \in \mathcal{C}$ is of size at most $2\binom{n-s-1}{r-s-1}$, we have that

$$\sum_{v=1}^{\binom{n}{r}} |\varphi^{-1}(\mathcal{C}(v))| = \sum_{c=1}^{2q} |\varphi(c)| \leq 4q\binom{n-s-1}{r-s-1}.$$

We thus obtain that there are at most $\frac{1}{2}\binom{n}{r}$ vertices such that

$$|\varphi^{-1}(\mathcal{C}(v))| \geq \frac{8q\binom{n-s-1}{r-s-1}}{\binom{n}{r}} \leq 8\delta\binom{r}{s+1}\ln n.$$

Now consider any v such that $|\varphi^{-1}(\mathcal{C}(v))| \leq 8\delta\binom{r}{s+1}\ln n$. We can bound $\mathbf{P}(\xi_v = 0)$ for big enough n as follows.

$$\mathbf{P}(\xi_v = 0) = 1 - \mathbf{P}(\xi_v = 1) = 1 - \frac{\binom{2q-|\varphi^{-1}(\mathcal{C}(v))|}{q}}{\binom{2q}{q}} = 1 - \prod_{i=0}^{q-1} \frac{2q - |\varphi^{-1}(\mathcal{C}(v))| - i}{2q - i} \leq$$

$$\leq 1 - \left(\frac{q - 8\delta\binom{r}{s+1}\ln n}{2q}\right)^{8\delta\binom{r}{s+1}\ln n} \leq 1 - 2^{-(1+\varepsilon)8\delta\binom{r}{s+1}\ln n} = 1 - n^{-(1+\varepsilon)8\delta\binom{r}{s+1}\ln 2}.$$

We thus have

$$\mathbf{P}\left(\sum_{i=1}^{\binom{n}{r}} \xi_i = 0\right) \leq \left(1 - n^{-(1+\varepsilon)8\delta\binom{r}{s+1}\ln 2}\right)^{\frac{1}{2}\binom{n}{r}} \leq \exp\left\{-\frac{1}{2}\binom{n}{r} n^{-(1+\varepsilon)8\delta\binom{r}{s+1}\ln 2}\right\}.$$

It remains to notice that taking union bound over all φ yields

$$\mathbf{P}(\text{exists at least 1 good } \varphi) \leq (\exp\{Cn^s \ln^2 n\})^{2\delta n^{s+1}\ln n} \exp\left\{-\frac{1}{2}\binom{n}{r} n^{-(1+\varepsilon)8\delta\binom{r}{s+1}\ln 2}\right\} =$$

$$= \exp\left\{2\delta C n^{2s+1}\ln^3 n - \frac{1}{2}\binom{n}{r} n^{-(1+\varepsilon)8\delta\binom{r}{s+1}\ln 2}\right\} =$$

$$= \exp\{2\delta C n^{2s+1}\ln^3 n - \Omega(n^{r-(1-\varepsilon^2)(r-2s-1)})\} = o(1).$$

It remains to notice that we are done since ε can be made arbitrarily small. ■

4.2 Proof of Theorem 10

Proof. Let us take $\delta = \frac{1}{120}$. The main idea is the following. We take $2q = 2\left\lfloor \delta n \ln\left(\frac{en}{r}\right)\right\rfloor$ colors and randomly assign each vertex a list containing half of these colors. Assume that we managed to color our graph properly. Let B_i be the set of vertices of color i. Then B_i is contained in some set from (any) container. This means that we can assign each color c a set $\varphi(c)$ from the container such that for each i and each $v \in B_i$ we have at least one C from the container such that $v \in C$ and $\varphi(i) = C$. It remains to prove that with positive probability for any φ there exists at least one such v that $v \notin \bigcup_{j=1}^{q} \varphi(c_j)$, where $\{c_1, \ldots, c_q\}$ is the list of colors assigned to v. We call φ with this condition bad and all other φ good.

Let us now give a rigorous proof. Consider the container $\mathcal{C}$ from Corollary `[unresolved original author label: coroll]`. Let us fix some function φ that maps each color to some set $C \in \mathcal{C}$. Let us bound the probability that φ is good. Let $\mathcal{C}(v)$ be the collection of all sets in $\mathcal{C}$ containing v. Let also list(v) be the list of colors for v. Finally, put $\xi_v = \mathbb{1}\left\{\text{list}(v) \cap \varphi^{-1}(\mathcal{C}(v)) = \varnothing\right\}$. Then φ is good if and only if $\sum_{i=1}^{\binom{n}{r}} \xi_i = 0$. Since all ξ_i are independent, we have

$$\mathbf{P}\left(\sum_{i=1}^{\binom{n}{r}} \xi_i = 0\right) = \prod_{i=1}^{\binom{n}{r}} \mathbf{P}(\xi_i = 0).$$

Let us now bound $\mathbf{P}(\xi_v = 0)$. In fact we will not bound all such probabilities, but only the ones with small $\varphi^{-1}(\mathcal{C}(v))$. Note that since each $C \in \mathcal{C}$ is of size at most $2\binom{n-1}{r-1}$, we have that

$$\sum_{v=1}^{\binom{n}{r}} |\varphi^{-1}(\mathcal{C}(v))| = \sum_{c=1}^{2q} |\varphi(c)| \leq 4q\binom{n-1}{r-1}.$$

We thus obtain that there are at most $\frac{1}{2}\binom{n}{r}$ vertices such that

$$|\varphi^{-1}(\mathcal{C}(v))| \geq \frac{8q\binom{n-1}{r-1}}{\binom{n}{r}} \leq 8\delta r \ln\left(\frac{en}{r}\right).$$

Now consider any v such that $|\varphi^{-1}(\mathcal{C}(v))| \leq \frac{8qr}{n}$. We can bound $\mathbf{P}(\xi_v = 0)$ for big enough n as follows.

$$\mathbf{P}(\xi_v = 0) = 1 - \mathbf{P}(\xi_v = 1) = 1 - \frac{\binom{2q-|\varphi^{-1}(\mathcal{C}(v))|}{q}}{\binom{2q}{q}} = 1 - \prod_{i=0}^{q-1} \frac{2q - |\varphi^{-1}(\mathcal{C}(v))| - i}{2q - i} \leq$$

$$\leq 1 - \left(\frac{q - q/2}{2q}\right)^{8\delta r \ln\left(\frac{en}{r}\right)} \leq 1 - 4^{-8\delta r \ln\left(\frac{en}{r}\right)} \leq 1 - n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}.$$

We thus have

$$\mathbf{P}\left(\sum_{i=1}^{\binom{n}{r}} \xi_i = 0\right) \leq \left(1 - n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right)^{\frac{1}{2}\binom{n}{r}} \leq \exp\left\{-\frac{1}{2}\binom{n}{r} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right\}.$$

It remains to notice that taking union bound over all φ yields

$$\mathbf{P}(\text{exists at least 1 good } \varphi) \leq 2^{2q} \left(\frac{e\binom{n}{r}}{k}\right)^{2kq} \exp\left\{-\frac{1}{2}\binom{n}{r} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right\} =$$

$$= \exp\left(2q \ln 2 + 2kq + 2kq \ln \binom{n}{r} - 2kq \ln k - \frac{1}{2}\binom{n}{r} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right) \leq$$

$$\leq \exp\left(4kqr \ln\left(\frac{en}{r}\right) - 2kq \ln k - \frac{1}{2}\binom{n}{r} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right) \leq$$

$$\leq \exp\left(4kqr \ln\left(\frac{en}{r}\right) - \frac{1}{2}\binom{n}{r} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right) \leq$$

$$\leq \exp\left(25\delta rn \frac{\binom{n-1}{r-1} \ln\left(\frac{en}{r}\right) \ln\binom{n}{r}}{\binom{n-r-1}{r-1}} \ln\left(\frac{en}{r}\right) - \frac{1}{2}\binom{n}{r} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right) \leq$$

$$\leq \exp\left(\binom{n}{r}\left(25\delta \frac{r^3}{\binom{n-r-1}{r-1}} \ln^3\left(\frac{en}{r}\right) - \frac{1}{2} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right)\right) \leq$$

$$\leq \exp\left(\binom{n}{r}\left(25\delta \frac{r! r^2}{(n-2r)^{r-1}} \ln^3\left(\frac{en}{r}\right) - \frac{1}{2} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right)\right) \leq$$

$$\leq \exp\left(\binom{n}{r}\left(\frac{r^{r+3}}{n^{r-1}} \ln^3\left(\frac{en}{r}\right) - \frac{1}{2} n^{-12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n}}\right)\right).$$

That is, if inequality

$$-r + 1 + (r+3)\frac{\ln r}{\ln n} \leq 4 - r\left(1 - \frac{\ln r}{\ln n}\right) < -12\delta \frac{r \ln\left(\frac{en}{r}\right)}{\ln n},$$

holds we are done. A sufficient condition is

$$r\left(1 - \frac{\ln r}{\ln n} - 12\delta \frac{\ln\left(\frac{en}{r}\right)}{\ln n}\right) = r\left(1 - 12\delta - \frac{12\delta}{\ln n} - (1 - 12\delta)\frac{\ln r}{\ln n}\right) = (1 - 12\delta) r\left(1 - \frac{\ln r}{\ln n}\right) - \frac{12\delta r}{\ln n} > 4.$$

Using the fact that $4 \ln n \leq r \leq n/20$ and $\delta = \frac{1}{120}$, we can see that

$$(1 - 12\delta) r\left(1 - \frac{\ln r}{\ln n}\right) - \frac{12\delta r}{\ln n} \geq 0.9r \frac{\ln 20}{\ln n} - \frac{0.1}{\ln n} r \geq \frac{\ln 20}{2 \ln n} r \geq 2 \ln 20 > 4$$

and thus we are done. ■

5 Upper bounds

5.1 Proof of Theorem 8

Proof. Let $\mathbf{Q}$ be the distribution on independent sets that minimizes $\max_v \mathbf{Q}^{-1}(v \in I)$. For each color c let us assign it to a random set I_c independently from everything else according to distribution $\mathbf{Q}$. Now for any vertex v and color c we say that

c is good for v if $v \in I_c$ and c is in list(v). Now we need to bound the probability that for each v we have at least one good color.

Note that since $\mathbf{Q}$ is taken to minimize $\max_v \mathbf{Q}^{-1}(v \in I)$ and I_c has distribution $\mathbf{Q}$ we have

$$\max_v \mathbf{Q}^{-1}(v \in I_c) = \chi_f(G)$$

and thus

$$\mathbf{Q}(v \in I_c) \geq \frac{1}{\chi_f(G)}$$

for any $v \in G$ and any color $c \in \text{list}(v)$. Now the probability that there is no good color for v can be bounded from above as $(1 - \chi_f^{-1}(G))^k \leq \exp\{-k\chi_f^{-1}(G)\} \leq (en)^{-1}$ and it remains to apply the union bound over all $v \in G$. ■

5.2 Proof of Theorem 9

Proof. Fix some lists of size k for all vertices.

Take $l = \left\lceil \frac{\chi_f(G)}{\sqrt{\ln a(G)}} \right\rceil$ and $p = \frac{2l}{k}$. For each color in at least one of the lists we say that it is a reserve color with probability p and main color otherwise. We do this independently for all colors. Let us now notice that for each vertex v we have $\xi_v \sim Bin(k, p)$ reserve colors.

It is easy to prove that there exists a coloring such that $\xi_v \in [l, 3l]$ for all v simultaneously. Indeed, we have $\mathbf{E}\xi_v = 2l$ and thus due to Chernoff's bound

$$1 - \mathbf{P}(\xi_v \in [l, 3l]) = \mathbf{P}\left(|\xi_v - \mathbf{E}\xi_v| > \frac{1}{2}\mathbf{E}\xi_v\right) \leq 2e^{-l/5} < 2n^{-2}.$$

Taking union bound over all v gives the desired result.

So we split all the colors into main and reserve. For now, drop all reserve colors from the lists, so only main colors are left. Note that the size of each list is still at least

$$k - 3l = \lceil \chi_f(G)\,(\ln a(G) + \ln\ln a(G)) \rceil.$$

Now let us perform the proof of Theorem 8 with slight modifications. Let $\mathbf{Q}$ be the distribution on independent sets that minimizes $\max_v \mathbf{Q}^{-1}(v \in I)$. For each color c let us assign it to a random set I_c independently from everything else according to distribution $\mathbf{Q}$. Now for any vertex v and color c we say that c is good for v if $v \in I_c$ and c is in list(v). The next thing to do is to bound the probability that for a vertex v we have at least one good color c

Note that since $\mathbf{Q}$ is taken to minimize $\max_v \mathbf{Q}^{-1}(v \in I)$ and I_c has distribution $\mathbf{Q}$ we have

$$\max_v \mathbf{Q}^{-1}(v \in I_c) = \chi_f(G)$$

and thus

$$\mathbf{Q}(v \in I_c) \geq \frac{1}{\chi_f(G)}$$

for any $v \in G$ and any color $c \in \text{list}(v)$. Now the probability that there is no good color for v can be bounded from above as

$$(1 - \chi_f^{-1}(G))^{k-3l} \leq \exp\{-(k-3l)\chi_f^{-1}(G)\} \leq \exp\{-\ln a(G) - \ln\ln a(G)\} \leq \frac{1}{a(G)\ln a(G)}.$$

We can now obtain the bound on the expected number of vertices without good color. If X is this number, then $\mathbf{E}X \leq \frac{n}{a(G)\ln a(G)}$. We can now use Markov inequality to obtain

$$\mathbf{Q}(X > l) \leq \frac{\mathbf{E}X}{l} \leq \frac{1}{\sqrt{\ln a(G)}} \leq \frac{1}{\sqrt{\ln 3}} < 1.$$

We can thus choose the sets I_c in such a way that at most l vertices will remain uncolored. But we can then easily color them using the reserve colors (since each list contains at least l of those). ■

References

[1] P. Frankl and Z. Füredi. Forbidding just one intersection. *J. Combin. Theory Ser. A*, 39:160–176, 1985.

[2] P. Frankl and R. Wilson. Intersection theorems with geometric consequences. *Combinatorica*, 1:357–368, 1981.

[3] F. J. MacWilliams and N. J. A. Sloane. *The theory of error-correcting codes.* North-Holland, 1977.

[4] Tuvi Etzion. On the nonexistence of perfect codes in the johnson scheme. *SIAM Journal on Discrete Mathematics*, 9(2):201–209, 1996.

[5] P. Delsart. An algebraic approach to the association schemes of coding theory. *Philips Res. Repts. Suppl.*, 10, 1973.

[6] A. M. Raigorodskii. Cliques and cycles in distance graphs and graphs of diameters. *Discrete Geometry and Algebraic Combinatorics, AMS, Contemporary Mathematics*, 625:93–109, 2014.

[7] M. Koshelev. Spectrum of johnson graphs. *Discrete Mathematics*, 346(3):113262, 2023.

[8] Ya. Shubin. On the minimal number of edges in induced subgraphs of special distance graphs. *Math. Notes*, 111(6):961–969, 2022.

[9] N. A. Dubinin, E. A. Neustroeva, A. M. Raigorodskii, and Ya. K. Shubin. The minimum number of cliques in induced subgraphs of johnson graphs. *Mathematical Notes*, 118(1):80–94, 2025.

[10] László Lovász. Kneser's conjecture, chromatic number, and homotopy. *Journal of Combinatorial Theory, Series A*, 25(3):319–324, 1978.

[11] A. M. Raigorodskii and P. S. Sinelnikov-Murylev. Johnson graphs, their random subgraphs, and some of their extremal characteristics. *Russian Math. Surveys*, 80:3:471–530, 2025.

[12] D. Zakharov. Chromatic numbers of kneser-type graphs. *J. Combin. Theory Ser. A*, 172:105188, 2020.

[13] D. A. Zakharov and A. M. Raigorodskii. Clique chromatic numbers of intersection graphs. *Math. Notes*, 105:137–139, 2019.

[14] A. Raigorodskii and M. Koshelev. New bounds on clique-chromatic numbers of johnson graphs. *Discrete Applied Mathematics*, 283:724–729, 2020.

[15] M. Koshelev. Lower bounds on the clique-chromatic numbers of some distance graphs. *Moscow Journal of Combinatorics and Number Theory*, 10(2):141–148, 2021.

[16] V. G. Vizing. Vertex colorings with given colors. *Diskret. Analiz*, 29:3–10, 1976. In Russian.

[17] Paul Erdős, Arthur L. Rubin, and Herbert Taylor. Choosability in graphs. In *Proceedings of the West Coast Conference on Combinatorics, Graph Theory and Computing (Humboldt State University, Arcata, California, 1979)*, volume 26 of *Congressus Numerantium*, pages 125–157, Winnipeg, 1980. Utilitas Mathematica Publishing.

[18] Noga Alon. Choice numbers of graphs: a probabilistic approach. *Combinatorics, Probability and Computing*, 1(2):107–114, 1992.

[19] Dmitry A. Shabanov. On a generalization of Rubin's theorem. *Journal of Graph Theory*, 67(3):226–234, 2011.

[20] Carsten Thomassen. Every planar graph is 5-choosable. *Journal of Combinatorial Theory, Series B*, 62(1):180–181, 1994.

[21] Fred Galvin. The list chromatic index of a bipartite multigraph. *Journal of Combinatorial Theory, Series B*, 63(1):153–158, 1995.

[22] Noga Alon, Michael Krivelevich, and Benny Sudakov. List coloring of random and pseudo-random graphs. *Combinatorica*, 19:453–472, 1999.

[23] Vera Bulankina and Andrey Kupavskii. Choice number of kneser graphs. *Discrete Mathematics*, 345(11):113097, 2022.

[24] Eduard Inozemtsev and Andrey Kupavskii. Two questions on kneser colorings. *Discrete Mathematics*, 349(2):114842, 2026.

[25] A. Kupavskii and Ya. Shubin. On supersaturation for Erdős-Sós problem. arxiv:2602.10292, 2026.

[26] David Saxton and Andrew Thomason. Hypergraph containers. *Inventiones Mathematicae*, 201(3):925–992, 2015.

[27] József Balogh, Robert Morris, and Wojciech Samotij. Independent sets in hypergraphs. *Journal of the American Mathematical Society*, 28(3):669–709, 2015.