REDUCEDNESS DESCENDS AND ASCENDS ALONG ETALE HOMOMORPHISMS OF NOETHERIAN LOCAL RINGS

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied verbatim from the pinned Stacks Project chapter file etale.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. Span recomputation by label pairing (LANE_BRIEF 10.2): in this chapter the author's proof environment follows the statement environment as a standalone block; the spec covers the complete environment pair (statement span, proof span) as recorded in the pool candidate.

Source reference. Original reference 033F: sources/stacks-chow/tags.txt, source lines 4054–4054. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Source reference. Original lemma 039L: sources/stacks-chow/tags.txt, source lines 4276–4276. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Source reference. Original reference 033B: sources/stacks-chow/tags.txt, source lines 4050–4050. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Proposition 0.1. *Let A, B be Noetherian local rings. Let $f : A \to B$ be an étale homomorphism of local rings. Then A is reduced if and only if B is so.*

Proof. It is clear from the faithful flatness of $A \to B$ that if B is reduced, so is A. See also Algebra, Lemma 033F. Conversely, assume A is reduced. By assumption B is a localization of a finite type A-algebra B' at some prime $\mathfrak{q}$. After replacing B' by a localization we may assume that B' is étale over A, see Lemma 039L. Then we see that Algebra, Lemma 033B applies to $A \to B'$ and B' is reduced. Hence B is reduced. □

Date: stacks-project:etale.tex:local-snapshot-20261012; GNU FDL 1.2 or later.