LOCAL UNRAMIFIED HOMOMORPHISMS ARE QUOTIENTS OF LOCALIZATIONS OF STANDARD ETALE ALGEBRAS

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied verbatim from the pinned Stacks Project chapter file etale.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. Span recomputation by label pairing (LANE_BRIEF 10.2): in this chapter the author's proof environment follows the statement environment as a standalone block; the spec covers the complete environment pair (statement span, proof span) as recorded in the pool candidate.

Source reference. Original lemma 039G: sources/stacks-chow/tags.txt, source lines 4271–4271. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Source reference. Original reference 0395: sources/stacks-chow/tags.txt, source lines 4260–4260. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Theorem 0.1. *Let $f : A \to B$ be an unramified morphism of local rings. Then there exist $f, g \in A[t]$ such that*

(1) *$B' = A[t]_g/(f)$ is standard étale – see (a) and (b) above, and*
(2) *B is isomorphic to a quotient of a localization of B' at a prime.*

Proof. Write $B = B'_{\mathfrak{q}}$ for some finite type A-algebra B' (we can do this because B is essentially of finite type over A). By Lemma 039G we see that $A \to B'$ is unramified at $\mathfrak{q}$. Hence we may apply Algebra, Proposition 0395 to see that a principal localization of B' is a quotient of a standard étale A-algebra. □

Date: stacks-project:etale.tex:local-snapshot-20261012; GNU FDL 1.2 or later.