THE NORMALISED COMPLEX OF AN EILENBERG-MACLANE OBJECT IS CONCENTRATED IN ONE DEGREE

THE STACKS PROJECT AUTHORS

Editorial note. The counted statement is the author's own lemma environment labelled lemma-N-K (source lines 4012-4019); the proof environment begins on the second line after it and is its own proof, closing at source line 4030.

Editorial note. The proof invokes two author results of the same chapter by label; both are retained as context spans and the generator expands the context transitively (budget 800 lines for this card), while reference targets in other chapters that carry canonical project tags are auto-declared as non-load-bearing source pointers.

Editorial note. The retained preamble is the chapter's own minimal preamble (sources/stacks-simplicial/preamble.tex, taken with its real line count), a verbatim excerpt of the native Xy-pic setup, the native theorem declarations with the mathematical macros, and the project's own no-op reference environments; no macro was invented or redefined.

Editorial scope. One original result of the Stacks Project's Simplicial Methods chapter, counted once: the normalised complex of K(A,k) is A concentrated in degree k.

Allowed standard prerequisites (editorial).

- Simplicial objects and the normalised complex. For a simplicial object of an abelian category, the normalised chain complex N, whose n-th term is the quotient of the n-simplices by the degenerate ones.. It is the object computed in the statement (source lines 4014-4018).
- Eilenberg-MacLane objects. The simplicial object K(A,k) built as the induced object from the skeleton, with K(A,k)_i = 0 for i below k, equal to A in degree k, and nondegenerate simplices only in degree k (the author's definition-eilenberg-maclane and lemma-imshriek-abelian, retained as context).. It supplies the three degree computations of the proof (source lines 4022-4029).

Source reference. Original lemma 002L: sources/stacks-chow/tags.txt, source lines 102–102. Cross-chapter pointer whose target carries the canonical tag 002L in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 002K: sources/stacks-chow/tags.txt, source lines 101–101. Cross-chapter pointer whose target carries the canonical tag 002K in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Date: stacks-project:simplicial.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

Lemma 0.1. *Let $\mathcal{A}$ be an abelian category Let U be an m-truncated simplicial object of $\mathcal{A}$. For $n > m$ we have $N(i_{m!}U)_n = 0$.*

Proof. Write $V = i_{m!}U$. Let $V' \subset V$ be the simplicial subobject of V with $V'_i = V_i$ for $i \leq m$ and $N(V'_i) = 0$ for $i > m$, see Lemma 0.3. By the adjunction formula, since $\mathrm{sk}_m V' = U$, there is an inverse to the injection $V' \to V$. Hence $V' = V$. □

Definition 0.2. Let $\mathcal{A}$ be an abelian category. Let A be an object of $\mathcal{A}$ and let k be an integer ≥ 0. The *Eilenberg-Maclane object* $K(A, k)$ is given by the object $K(A, k) = i_{k!}U$ which is described in Lemma 0.4 above.

Lemma 0.3. *Let $\mathcal{A}$ be an abelian category. Let U be a simplicial object of $\mathcal{A}$. Let $n \geq 0$ be an integer. The rule*

$$U'_m = \sum\nolimits_{\varphi:[m]\to[i],\ i\leq n} \mathrm{Im}(U(\varphi))$$

defines a sub simplicial object $U' \subset U$ with $U'_i = U_i$ for $i \leq n$. Moreover, $N(U'_m) = 0$ for all $m > n$.

Proof. Pick m, $i \leq n$ and some $\varphi : [m] \to [i]$. The image under $U(\psi)$ of $\mathrm{Im}(U(\varphi))$ for any $\psi : [m'] \to [m]$ is equal to the image of $U(\varphi \circ \psi)$ and $\varphi \circ \psi : [m'] \to [i]$. Hence U' is a simplicial object. Pick $m > n$. We have to show $N(U'_m) = 0$. By definition of $N(U_m)$ and $N(U'_m)$ we have $N(U'_m) = U'_m \cap N(U_m)$ (intersection of subobjects). Since U is split by Lemma 0.5, it suffices to show that U'_m is contained in the sum

$$\sum\nolimits_{\varphi:[m]\to[m'] \text{ surjective},\ m'<m} \mathrm{Im}(U(\varphi)|_{N(U_{m'})}).$$

By the splitting each $U_{m'}$ is the sum of images of $N(U_{m''})$ via $U(\psi)$ for surjective maps $\psi : [m'] \to [m'']$. Hence the displayed sum above is the same as

$$\sum\nolimits_{\varphi:[m]\to[m'] \text{ surjective},\ m'<m} \mathrm{Im}(U(\varphi)).$$

Clearly U'_m is contained in this by the simple fact that any $\varphi : [m] \to [i]$, $i \leq n$ occurring in the definition of U'_m may be factored as $[m] \to [m'] \to [i]$ with $[m] \to [m']$ surjective and $m' < m$ as in the last displayed sum above. □

Lemma 0.4. *With A, k and U as above, so $U_i = 0$, $i < k$ and $U_k = A$.*

(1) *Given a k-truncated simplicial object V we have*

$$\mathrm{Mor}(U, V) = \{f : A \to V_k \mid d_i^k \circ f = 0,\ i = 0, \ldots, k\}$$

and

$$\mathrm{Mor}(V, U) = \{f : V_k \to A \mid f \circ s_i^{k-1} = 0,\ i = 0, \ldots, k-1\}.$$

(2) *The object $i_{k!}U$ has nth term equal to $\bigoplus_\alpha A$ where α runs over all surjective morphisms $\alpha : [n] \to [k]$.*
(3) *For any $\varphi : [m] \to [n]$ the map $i_{k!}U(\varphi)$ is described as the mapping $\bigoplus_\alpha A \to \bigoplus_{\alpha'} A$ which maps to component corresponding to $\alpha : [n] \to [k]$ to zero if $\alpha \circ \varphi$ is not surjective and by the identity to the component corresponding to $\alpha \circ \varphi$ if it is surjective.*
(4) *The object $\mathrm{cosk}_k U$ has nth term equal to $\bigoplus_\beta A$, where β runs over all injective morphisms $\beta : [k] \to [n]$.*

(5) *For any $\varphi : [m] \to [n]$ the map $cosk_k U(\varphi)$ is described as the mapping $\bigoplus_\beta A \to \bigoplus_{\beta'} A$ which maps to component corresponding to $\beta : [k] \to [n]$ to zero if β does not factor through φ and by the identity to each of the components corresponding to β' such that $\beta = \varphi \circ \beta'$ if it does.*
(6) *The canonical map $c : i_{k!}U \to cosk_k U$ in degree n has (α, β) coefficient $A \to A$ equal to zero if $\alpha \circ \beta$ is not the identity and equal to id_A if it is.*
(7) *The canonical map $c : i_{k!}U \to cosk_k U$ is injective.*

Proof. The proof of (1) is left to the reader.

Let us take the rules of (2) and (3) as the definition of a simplicial object, call it $\tilde{U}$. We will show that it is an incarnation of $i_{k!}U$. This will prove (2), (3) at the same time. We have to show that given a morphism $f : U \to \mathrm{sk}_k V$ there exists a unique morphism $\tilde{f} : \tilde{U} \to V$ which recovers f upon taking the k-skeleton. From (1) we see that f corresponds with a morphism $f_k : A \to V_k$ which maps into the kernel of d_i^k for all i. For any surjective $\alpha : [n] \to [k]$ we set $\tilde{f}_\alpha : A \to V_n$ equal to the composition $\tilde{f}_\alpha = V(\alpha) \circ f_k : A \to V_n$. We define $\tilde{f}_n : \tilde{U}_n \to V_n$ as the sum of the $\tilde{f}_\alpha$ over $\alpha : [n] \to [k]$ surjective. Such a collection of $\tilde{f}_\alpha$ defines a morphism of simplicial objects if and only if for any $\varphi : [m] \to [n]$ the diagram

$$\begin{array}{ccc} \bigoplus_{\alpha:[n]\to[k] \text{ surjective}} A & \xrightarrow[\tilde{f}_n]{} & V_n \\ \downarrow{\scriptstyle (3)} & & \downarrow{\scriptstyle V(\varphi)} \\ \bigoplus_{\alpha':[m]\to[k] \text{ surjective}} A & \xrightarrow{\tilde{f}_m} & V_m \end{array}$$

is commutative. Choosing $\varphi = \alpha$ shows our choice of $\tilde{f}_\alpha$ is uniquely determined by f_k. The commutativity in general may be checked for each summand of the left upper corner separately. It is clear for the summands corresponding to α where $\alpha \circ \varphi$ is surjective, because those get mapped by id_A to the summand with $\alpha' = \alpha \circ \varphi$, and we have $\tilde{f}_{\alpha'} = V(\alpha') \circ f_k = V(\alpha \circ \varphi) \circ f_k = V(\varphi) \circ \tilde{f}_\alpha$. For those where $\alpha \circ \varphi$ is not surjective, we have to show that $V(\varphi) \circ \tilde{f}_\alpha = 0$. By definition this is equal to $V(\varphi) \circ V(\alpha) \circ f_k = V(\alpha \circ \varphi) \circ f_k$. Since $\alpha \circ \varphi$ is not surjective we can write it as $\delta_i^k \circ \psi$, and we deduce that $V(\varphi) \circ V(\alpha) \circ f_k = V(\psi) \circ d_i^k \circ f_k = 0$ see above.

Let us take the rules of (4) and (5) as the definition of a simplicial object, call it $\tilde{U}$. We will show that it is an incarnation of $\mathrm{cosk}_k U$. This will prove (4), (5) at the same time. The argument is completely dual to the proof of (2), (3) above, but we give it anyway. We have to show that given a morphism $f : \mathrm{sk}_k V \to U$ there exists a unique morphism $\tilde{f} : V \to \tilde{U}$ which recovers f upon taking the k-skeleton. From (1) we see that f corresponds with a morphism $f_k : V_k \to A$ which is zero on the image of s_i^{k-1} for all i. For any injective $\beta : [k] \to [n]$ we set $\tilde{f}_\beta : V_n \to A$ equal to the composition $\tilde{f}_\beta = f_k \circ V(\beta) : V_n \to A$. We define $\tilde{f}_n : V_n \to \tilde{U}_n$ as the sum of the $\tilde{f}_\beta$ over $\beta : [k] \to [n]$ injective. Such a collection of $\tilde{f}_\beta$ defines a morphism of simplicial objects if and only if for any $\varphi : [m] \to [n]$ the diagram

$$\begin{array}{ccc} V_n & \xrightarrow[\tilde{f}_n]{} & \bigoplus_{\beta:[k]\to[n] \text{ injective}} A \\ \downarrow{\scriptstyle V(\varphi)} & & \downarrow{\scriptstyle (5)} \\ V_m & \xrightarrow{\tilde{f}_m} & \bigoplus_{\beta':[k]\to[m] \text{ injective}} A \end{array}$$

is commutative. Choosing $\varphi = \beta$ shows our choice of $\tilde{f}_\beta$ is uniquely determined by f_k. The commutativity in general may be checked for each summand of the right lower corner separately. It is clear for the summands corresponding to β' where $\varphi \circ \beta'$ is injective, because these summands get mapped into by exactly the summand with $\beta = \varphi \circ \beta'$ and we have in that case $\tilde{f}_{\beta'} \circ V(\varphi) = f_k \circ V(\beta') \circ V(\varphi) = f_k \circ V(\beta) = \tilde{f}_\beta$. For those where $\varphi \circ \beta'$ is not injective, we have to show that $\tilde{f}_{\beta'} \circ V(\varphi) = 0$. By definition this is equal to $f_k \circ V(\beta') \circ V(\varphi) = f_k \circ V(\varphi \circ \beta')$. Since $\varphi \circ \beta'$ is not injective we can write it as $\psi \circ \sigma_i^{k-1}$, and we deduce that $f_k \circ V(\beta') \circ V(\varphi) = f_k \circ s_i^{k-1} \circ V(\psi) = 0$ see above.

The composition $i_{k!}U \to \mathrm{cosk}_k U$ is the unique map of simplicial objects which is the identity on $A = U_k = (i_{k!}U)_k = (\mathrm{cosk}_k U)_k$. Hence it suffices to check that the proposed rule defines a morphism of simplicial objects. To see this we have to show that for any $\varphi : [m] \to [n]$ the diagram

$$\begin{array}{ccc} \bigoplus_{\alpha:[n]\to[k] \text{ surjective}} A & \xrightarrow[(6)]{} & \bigoplus_{\beta:[k]\to[n] \text{ injective}} A \\ \downarrow{\scriptstyle (3)} & & \downarrow{\scriptstyle (5)} \\ \bigoplus_{\alpha':[m]\to[k] \text{ surjective}} A & \xrightarrow{(6)} & \bigoplus_{\beta':[k]\to[m] \text{ injective}} A \end{array}$$

is commutative. Now we can think of this in terms of matrices filled with only 0's and 1's as follows: The matrix of (3) has a nonzero (α', α) entry if and only if $\alpha' = \alpha \circ \varphi$. Likewise the matrix of (5) has a nonzero (β', β) entry if and only if $\beta = \varphi \circ \beta'$. The upper matrix of (6) has a nonzero (α, β) entry if and only if $\alpha \circ \beta = \mathrm{id}_{[k]}$. Similarly for the lower matrix of (6). The commutativity of the diagram then comes down to computing the (α, β') entry for both compositions and seeing they are equal. This comes down to the following equality

$$\#\left\{\beta \mid \beta = \varphi \circ \beta' \text{ and } \alpha \circ \beta = \mathrm{id}_{[k]}\right\} = \#\left\{\alpha' \mid \alpha' = \alpha \circ \varphi \text{ and } \alpha' \circ \beta' = \mathrm{id}_{[k]}\right\}$$

whose proof may safely be left to the reader.

Finally, we prove (7). This follows directly from Lemmas 0.6, 0.7, 0.8 and 0.1. □

Lemma 0.5. *Let $\mathcal{A}$ be an abelian category. Let U be a simplicial object in $\mathcal{A}$. Then U has a splitting obtained by taking $N(U_0) = U_0$ and for $m \geq 1$ taking*

$$N(U_m) = \bigcap\nolimits_{i=0}^{m-1} \mathrm{Ker}(d_i^m).$$

Moreover, this splitting is functorial on the category of simplicial objects of $\mathcal{A}$.

Proof. For any object A of $\mathcal{A}$ we obtain a simplicial abelian group $\mathrm{Mor}_{\mathcal{A}}(A, U)$. Each of these are canonically split by Lemma 0.9. Moreover,

$$N(\mathrm{Mor}_{\mathcal{A}}(A, U_m)) = \bigcap\nolimits_{i=0}^{m-1} \mathrm{Ker}(d_i^m) = \mathrm{Mor}_{\mathcal{A}}(A, N(U_m)).$$

Hence we see that the morphism (0.8.1) becomes an isomorphism after applying the functor $\mathrm{Mor}_{\mathcal{A}}(A, -)$ for any object of $\mathcal{A}$. Hence it is an isomorphism by the Yoneda lemma. □

Lemma 0.6. *Let $\mathcal{A}$ be an abelian category. Let $f : U \to V$ be a morphism of simplicial objects of $\mathcal{A}$. If the induced morphisms $N(f)_i : N(U)_i \to N(V)_i$ are injective for all i, then f_i is injective for all i. Same holds with "injective" replaced with "surjective", or "isomorphism".*

Proof. This is clear from Lemma 0.5 and the definition of a splitting. □

Lemma 0.7. *Let $\mathcal{C}$ be a category with finite limits. Let U be an n-truncated simplicial object of $\mathcal{C}$. The morphism $sk_n cosk_n U \to U$ is an isomorphism.*

Proof. Combine Lemmas 0.10 and 0.11. □

Lemma 0.8. *Let $\mathcal{C}$ be a category which has finite colimits. Let U be an m-truncated simplicial object of $\mathcal{C}$. The map $U \to sk_m i_{m!} U$ is an isomorphism.*

Proof. Combine Lemmas 0.12 and 0.13. □

$$\coprod_{\varphi:[n]\to[m] \text{ surjective}} N(U_m) \longrightarrow U_n \tag{0.8.1}$$

Lemma 0.9. *Let U be a simplicial abelian group. Then U has a splitting obtained by taking $N(U_0) = U_0$ and for $m \geq 1$ taking*

$$N(U_m) = \bigcap\nolimits_{i=0}^{m-1} \mathrm{Ker}(d_i^m).$$

Moreover, this splitting is functorial on the category of simplicial abelian groups.

Proof. By induction on n we will show that the choice of $N(U_m)$ in the lemma guarantees that (0.8.1) is an isomorphism for $m \leq n$. This is clear for $n = 0$. In the rest of this proof we are going to drop the superscripts from the maps d_i and s_i in order to improve readability. We will also repeatedly use the relations from Remark 0.14.

First we make a general remark. For $0 \leq i \leq m$ and $z \in U_m$ we have $d_i(s_i(z)) = z$. Hence we can write any $x \in U_{m+1}$ uniquely as $x = x' + x''$ with $d_i(x') = 0$ and $x'' \in \mathrm{Im}(s_i)$ by taking $x' = (x - s_i(d_i(x)))$ and $x'' = s_i(d_i(x))$. Moreover, the element $z \in U_m$ such that $x'' = s_i(z)$ is unique because s_i is injective.

Here is a procedure for decomposing any $x \in U_{n+1}$. First, write $x = x_0 + s_0(z_0)$ with $d_0(x_0) = 0$. Next, write $x_0 = x_1 + s_1(z_1)$ with $d_1(x_1) = 0$. Continue like this to get

$$\begin{aligned} x &= x_0 + s_0(z_0), \\ x_0 &= x_1 + s_1(z_1), \\ x_1 &= x_2 + s_2(z_2), \\ \ldots &\quad \ldots \quad \ldots \\ x_{n-1} &= x_n + s_n(z_n) \end{aligned}$$

where $d_i(x_i) = 0$ for all $i = n, \ldots, 0$. By our general remark above all of the x_i and z_i are determined uniquely by x. We claim that $x_i \in \mathrm{Ker}(d_0) \cap \mathrm{Ker}(d_1) \cap \ldots \cap \mathrm{Ker}(d_i)$ and $z_i \in \mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})$ for $i = n, \ldots, 0$. Here and in the following an empty intersection of kernels indicates the whole space; i.e., the notation $z_0 \in \mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})$ when $i = 0$ means $z_0 \in U_n$ with no restriction.

We prove this by ascending induction on i. It is clear for $i = 0$ by construction of x_0 and z_0. Let us prove it for $0 < i \leq n$ assuming the result for $i - 1$. First of all we have $d_i(x_i) = 0$ by construction. So pick a j with $0 \leq j < i$. We have $d_j(x_{i-1}) = 0$ by induction. Hence

$$0 = d_j(x_{i-1}) = d_j(x_i) + d_j(s_i(z_i)) = d_j(x_i) + s_{i-1}(d_j(z_i)).$$

The last equality by the relations of Remark 0.14. These relations also imply that $d_{i-1}(d_j(x_i)) = d_j(d_i(x_i)) = 0$ because $d_i(x_i) = 0$ by construction. Then the uniqueness in the general remark above shows the equality $0 = x' + x'' = d_j(x_i) + s_{i-1}(d_j(z_i))$ can only hold if both terms are zero. We conclude that $d_j(x_i) = 0$ and by injectivity of s_{i-1} we also conclude that $d_j(z_i) = 0$. This proves the claim.

The claim implies we can uniquely write

$$x = s_0(z_0) + s_1(z_1) + \ldots + s_n(z_n) + x_n$$

with $x_n \in N(U_{n+1})$ and $z_i \in \mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})$. We can reformulate this as saying that we have found a direct sum decomposition

$$U_{n+1} = N(U_{n+1}) \oplus \bigoplus\nolimits_{i=0}^{i=n} s_i\Big(\mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})\Big)$$

with the property that

$$\mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_j) = N(U_{n+1}) \oplus \bigoplus\nolimits_{i=j+1}^{i=n} s_i\Big(\mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})\Big)$$

for $j = 0, \ldots, n$. The result follows from this statement as follows. Each of the z_i in the expression for x can be written uniquely as

$$z_i = s_i(z'_{i,i}) + \ldots + s_{n-1}(z'_{i,n-1}) + z_{i,0}$$

with $z_{i,0} \in N(U_n)$ and $z'_{i,j} \in \mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{j-1})$. The first few steps in the decomposition of z_i are zero because z_i already is in the kernel of $d_0, \ldots, d_i$. This in turn uniquely gives

$$x = x_n + s_0(z_{0,0}) + s_1(z_{1,0}) + \ldots + s_n(z_{n,0}) + \sum\nolimits_{0 \leq i \leq j \leq n-1} s_i(s_j(z'_{i,j})).$$

Continuing in this fashion we see that we in the end obtain a decomposition of x as a sum of terms of the form

$$s_{i_1} s_{i_2} \ldots s_{i_k}(z)$$

with $0 \leq i_1 \leq i_2 \leq \ldots \leq i_k \leq n - k + 1$ and $z \in N(U_{n+1-k})$. This is exactly the required decomposition, because any surjective map $[n+1] \to [n+1-k]$ can be uniquely expressed in the form

$$\sigma_{i_k}^{n-k} \ldots \sigma_{i_2}^{n-1} \sigma_{i_1}^{n}$$

with $0 \leq i_1 \leq i_2 \leq \ldots \leq i_k \leq n - k + 1$. □

Lemma 0.10. *If the category $\mathcal{C}$ has finite limits, then $cosk_m$ functors exist for all m. Moreover, for any m-truncated simplicial object U the simplicial object $cosk_m U$ is described by the formula*

$$(cosk_m U)_n = \lim\nolimits_{(\Delta/[n])^{opp}_{\leq m}} U(n)$$

and for $\varphi : [n] \to [n']$ the map $cosk_m U(\varphi)$ comes from the identification $U(n') \circ \overline{\varphi} = U(n)$ above via Categories, Lemma 002L.

Proof. During the proof of this lemma we denote $\mathrm{cosk}_m U$ the simplicial object with $(\mathrm{cosk}_m U)_n$ equal to $\lim_{(\Delta/[n])^{opp}_{\leq m}} U(n)$. We will conclude at the end of the proof that it does satisfy the required mapping property.

Suppose that V is a simplicial object. A morphism $\gamma : V \to \mathrm{cosk}_m U$ is given by a sequence of morphisms $\gamma_n : V_n \to (\mathrm{cosk}_m U)_n$. By definition of a limit, this is given

by a collection of morphisms $\gamma(\alpha) : V_n \to U_k$ where α ranges over all $\alpha : [k] \to [n]$ with $k \leq m$. These morphisms then also satisfy the rules that

$$\begin{array}{ccc} V_n & \xrightarrow[\gamma(\alpha)]{} & U_k \\ {\scriptstyle V(\varphi)}\uparrow & & \uparrow{\scriptstyle U(\psi)} \\ V_{n'} & \xrightarrow{\gamma(\alpha')} & U_{k'} \end{array}$$

are commutative, given any $0 \leq k, k' \leq m$, $0 \leq n, n'$ and any $\psi : [k] \to [k']$, $\varphi : [n] \to [n']$, $\alpha : [k] \to [n]$ and $\alpha' : [k'] \to [n']$ in Δ such that $\varphi \circ \alpha = \alpha' \circ \psi$. Taking $n = k = k'$, $\varphi = \alpha'$, and $\alpha = \psi = \mathrm{id}_{[k]}$ we deduce that $\gamma(\alpha') = \gamma(\mathrm{id}_{[k]}) \circ V(\alpha')$. In other words, the morphisms $\gamma(\mathrm{id}_{[k]})$, $k \leq m$ determine the morphism γ. And it is easy to see that these morphisms form a morphism $\mathrm{sk}_m V \to U$.

Conversely, given a morphism $\gamma : \mathrm{sk}_m V \to U$, we obtain a family of morphisms $\gamma(\alpha)$ where α ranges over all $\alpha : [k] \to [n]$ with $k \leq m$ by setting $\gamma(\alpha) = \gamma(\mathrm{id}_{[k]}) \circ V(\alpha)$. These morphisms satisfy all the displayed commutativity restraints pictured above, and hence give rise to a morphism $V \to \mathrm{cosk}_m U$. □

Lemma 0.11. *Let $\mathcal{C}$ be a category. Let U be an m-truncated simplicial object of $\mathcal{C}$. For $n \leq m$ the limit $\lim_{(\Delta/[n])^{opp}_{\leq m}} U(n)$ exists and is canonically isomorphic to U_n.*

Proof. This is true because the category $(\Delta/[n])_{\leq m}$ has an final object in this case, namely the identity map $[n] \to [n]$. □

Lemma 0.12. *Let $\mathcal{C}$ be a category which has finite colimits. The functors $i_{m!}$ exist for all m. Let U be an m-truncated simplicial object of $\mathcal{C}$. The simplicial object $i_{m!}U$ is described by the formula*

$$(i_{m!}U)_n = \mathrm{colim}_{([n]/\Delta)^{opp}_{\leq m}} U(n)$$

and for $\varphi : [n] \to [n']$ the map $i_{m!}U(\varphi)$ comes from the identification $U(n) \circ \underline{\varphi} = U(n')$ above via Categories, Lemma 002K.

Proof. In this proof we denote $i_{m!}U$ the simplicial object whose nth term is given by the displayed formula of the lemma. We will show it satisfies the adjointness property.

Let V be a simplicial object of $\mathcal{C}$. Let $\gamma : U \to \mathrm{sk}_m V$ be given. A morphism

$$\mathrm{colim}_{([n]/\Delta)^{opp}_{\leq m}} U(n) \to T$$

is given by a compatible system of morphisms $f_\alpha : U_k \to T$ where $\alpha : [n] \to [k]$ with $k \leq m$. Certainly, we have such a system of morphisms by taking the compositions

$$U_k \xrightarrow{\gamma_k} V_k \xrightarrow{V(\alpha)} V_n.$$

Hence we get an induced morphism $(i_{m!}U)_n \to V_n$. We leave it to the reader to see that these form a morphism of simplicial objects $\gamma' : i_{m!}U \to V$.

Conversely, given a morphism $\gamma' : i_{m!}U \to V$ we obtain a morphism $\gamma : U \to \mathrm{sk}_m V$ by setting $\gamma_i : U_i \to V_i$ equal to the composition

$$U_i \xrightarrow{\mathrm{id}_{[i]}} \mathrm{colim}_{([i]/\Delta)^{opp}_{\leq m}} U(i) \xrightarrow{\gamma'_i} V_i$$

for $0 \leq i \leq n$. We leave it to the reader to see that this is the inverse of the construction above. □

Lemma 0.13. *Let $\mathcal{C}$ be a category. Let U be an m-truncated simplicial object of $\mathcal{C}$. For any $n \leq m$ the colimit*

$$\mathrm{colim}_{([n]/\Delta)^{opp}_{\leq m}} U(n)$$

exists and is equal to U_n.

Proof. This is so because the category $([n]/\Delta)_{\leq m}$ has an initial object, namely $\mathrm{id} : [n] \to [n]$. □

Remark 0.14. By abuse of notation we sometimes write $d_i : U_n \to U_{n-1}$ instead of d_i^n, and similarly for $s_i : U_n \to U_{n+1}$. The relations among the morphisms d_i^n and s_i^n may be expressed as follows:

(1) If $i < j$, then $d_i \circ d_j = d_{j-1} \circ d_i$.
(2) If $i < j$, then $d_i \circ s_j = s_{j-1} \circ d_i$.
(3) We have $\mathrm{id} = d_j \circ s_j = d_{j+1} \circ s_j$.
(4) If $i > j + 1$, then $d_i \circ s_j = s_j \circ d_{i-1}$.
(5) If $i \leq j$, then $s_i \circ s_j = s_{j+1} \circ s_i$.

This means that whenever the compositions on both the left and the right are defined then the corresponding equality should hold.

Lemma 0.15. *Let $\mathcal{A}$ be an abelian category. Let A be an object of $\mathcal{A}$ and let k be an integer. We have $N(K(A,k))_i = A$ if $i = k$ and 0 else.*

Proof. It is clear that $N(K(A,k))_i = 0$ when $i < k$ because $K(A,k)_i = 0$ in that case. It is clear that $N(K(A,k))_k = A$ since $K(A,k)_{k-1} = 0$ and $K(A,k)_k = A$. For $i > k$ we have $N(K(A,k))_i = 0$ by Lemma 0.1 and the definition of $K(A,k)$, see Definition 0.2. □