A TERMWISE SURJECTIVE QUASI-ISOMORPHISM OF SIMPLICIAL ABELIAN GROUPS IS A TRIVIAL KAN FIBRATION

THE STACKS PROJECT AUTHORS

Editorial note. The counted statement is the author's own lemma environment labelled lemma-qis-simplicial-abelian-groups (source lines 6053-6059); the proof environment begins on the second line after it and is its own proof, closing at source line 6099.

Editorial note. The statement and the proof invoke two author results of the same chapter by label; both are retained as context spans and the generator expands the context transitively (budget 800 lines for this card), while reference targets in other chapters that carry canonical project tags are auto-declared as non-load-bearing source pointers.

Editorial note. The retained preamble is the chapter's own minimal preamble (sources/stacks-simplicial/preamble.tex, taken with its real line count), a verbatim excerpt of the native Xy-pic setup, the native theorem declarations with the mathematical macros, and the project's own no-op reference environments; no macro was invented or redefined.

Editorial scope. One original result of the Stacks Project's Simplicial Methods chapter, counted once: a termwise surjective homomorphism of simplicial abelian groups inducing a quasi-isomorphism on associated complexes is a trivial Kan fibration.

Allowed standard prerequisites (editorial).

- Simplicial abelian groups and associated chain complexes. The associated chain complex s(X) of a simplicial abelian group, its normalised complex N(X), and the definition of a quasi-isomorphism between them.. It is the setting and the acyclicity input (source lines 6055-6057, 6092-6095).
- Trivial Kan fibrations. The lifting property defining a trivial Kan fibration, expressed as the datum of the horn maps a and b (the author's definition-trivial-kan, retained as context).. It is what the proof has to establish (source lines 6062-6073).
- Simplicial groups are Kan complexes. Every simplicial group, in particular every simplicial abelian group, is a Kan complex (the author's lemma-simplicial-group-kan, retained as context).. It supplies the horn filler x in degree n (source lines 6087-6088).
- The long exact sequence of homology. The long exact homology sequence of a short exact sequence of complexes (Homology, Lemma homology-lemma-long-exact-sequence-chain, declared as a tagged source pointer).. It shows that the kernel of f has vanishing homology, so that the reduction to $Y = 0$ stays a quasi-isomorphism (source lines 6080-6084).

Date: stacks-project:simplicial.tex:local-snapshot-20261012; GNU FDL 1.2 or later.

Source reference. Original lemma 0111: sources/stacks-chow/tags.txt, source lines 1347–1347. Cross-chapter pointer whose target carries the canonical tag 0111 in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 0110: sources/stacks-chow/tags.txt, source lines 1346–1346. Cross-chapter pointer whose target carries the canonical tag 0110 in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 002L: sources/stacks-chow/tags.txt, source lines 102–102. Cross-chapter pointer whose target carries the canonical tag 002L in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Source reference. Original lemma 002K: sources/stacks-chow/tags.txt, source lines 101–101. Cross-chapter pointer whose target carries the canonical tag 002K in the project tag list; declared so the reference resolves inside the assembled package, while every step the counted argument uses remains inside the retained spans.

Definition 0.1. A map $X \to Y$ of simplicial sets is called a *trivial Kan fibration* if $X_0 \to Y_0$ is surjective and for all $n \geq 1$ and any commutative solid diagram

$$\begin{array}{ccc} \partial\Delta[n] & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow \\ \Delta[n] & \longrightarrow & Y \end{array}$$

a dotted arrow exists making the diagram commute.

Lemma 0.2. *Let X be a simplicial group. Then X is a Kan complex.*

Proof. The following proof is basically just a translation into English of the proof in the reference mentioned above. Using the terminology as explained in the introduction to this section, suppose $f : \Lambda_k[n] \to X$ is a morphism from a horn. Set $x_i = f(\sigma_i) \in X_{n-1}$ for $i = 0, \ldots, \hat{k}, \ldots, n$. This means that for $i < j$ we have $d_i x_j = d_{j-1} x_i$ whenever $i, j \neq k$. We have to find an $x \in X_n$ such that $x_i = d_i x$ for $i = 0, \ldots, \hat{k}, \ldots, n$.

We first prove there exists a $u \in X_n$ such that $d_i u = x_i$ for $i < k$. This is trivial for $k = 0$. If $k > 0$, one defines by induction an element $u^r \in X_n$ such that $d_i u^r = x_i$ for $0 \leq i \leq r$. Start with $u^0 = s_0 x_0$. If $r < k - 1$, we set

$$y^r = s_{r+1}((d_{r+1}u^r)^{-1} x_{r+1}), \quad u^{r+1} = u^r y^r.$$

An easy calculation shows that $d_i y^r = 1$ (unit element of the group X_{n-1}) for $i \leq r$ and $d_{r+1} y^r = (d_{r+1} u^r)^{-1} x_{r+1}$. It follows that $d_i u^{r+1} = x_i$ for $i \leq r + 1$. Finally, take $u = u^{k-1}$ to get u as promised.

Next we prove, by induction on the integer r, $0 \leq r \leq n - k$, there exists a $x^r \in X_n$ such that

$$d_i x^r = x_i \quad \text{for } i < k \text{ and } i > n - r.$$

Start with $x^0 = u$ for $r = 0$. Having defined x^r for $r \leq n - k - 1$ we set

$$z^r = s_{n-r-1}((d_{n-r} x^r)^{-1} x_{n-r}), \quad x^{r+1} = x^r z^r$$

A simple calculation, using the given relations, shows that $d_i z^r = 1$ for $i < k$ and $i > n - r$ and that $d_{n-r}(z^r) = (d_{n-r}x^r)^{-1}x_{n-r}$. It follows that $d_i x^{r+1} = x_i$ for $i < k$ and $i > n - r - 1$. Finally, we take $x = x^{n-k}$ which finishes the proof. □

Lemma 0.3. *Let $\mathcal{A}$ be an abelian category. Let V be a simplicial object of $\mathcal{A}$. The canonical morphism of chain complexes $N(V) \to s(V)$ is a quasi-isomorphism. In other words, the complex $D(V)$ of Lemma 1.4 is acyclic.*

Proof. Note that the result holds for $K(A, k)$ for any object A and any $k \geq 0$, by Lemmas 1.6 and 1.2. Consider the hypothesis $IH_{n,m}$: for all V such that $V_j = 0$ for $j \leq m$ and all $i \leq n$ the map $N(V) \to s(V)$ induces an isomorphism $H_i(N(V)) \to H_i(s(V))$.

To start of the induction, note that $IH_{n,n}$ is trivially true, because in that case $N(V)_n = 0$ and $s(V)_n = 0$.

Assume $IH_{n,m}$, with $m \leq n$. Pick a simplicial object V such that $V_j = 0$ for $j < m$. By Lemma 1.5 and Definition 1.1 we have $K(V_m, m) = i_{m!}\mathrm{sk}_m V$. By Lemma 1.7 the natural morphism

$$K(V_m, m) = i_{m!}\mathrm{sk}_m V \to V$$

is injective. Thus we get a short exact sequence

$$0 \to K(V_m, m) \to V \to W \to 0$$

for some W with $W_i = 0$ for $i = 0, \ldots, m$. This short exact sequence induces a morphism of short exact sequence of associated complexes

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N(K(V_m, m)) & \longrightarrow & N(V) & \longrightarrow & N(W) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & s(K(V_m, m)) & \longrightarrow & s(V) & \longrightarrow & s(W) & \longrightarrow & 0 \end{array}$$

see Lemmas 1.8 and 1.3. Hence we deduce the result for V from the result on the ends. □

1. Simplicial objects and chain complexes

Definition 1.1. Let $\mathcal{A}$ be an abelian category. Let A be an object of $\mathcal{A}$ and let k be an integer ≥ 0. The *Eilenberg-Maclane object* $K(A, k)$ is given by the object $K(A, k) = i_{k!}U$ which is described in Lemma 1.5 above.

Lemma 1.2. *Let $\mathcal{A}$ be an abelian category. Let A be an object of $\mathcal{A}$ and let k be an integer. We have $N(K(A, k))_i = A$ if $i = k$ and 0 else.*

Proof. It is clear that $N(K(A, k))_i = 0$ when $i < k$ because $K(A, k)_i = 0$ in that case. It is clear that $N(K(A, k))_k = A$ since $K(A, k)_{k-1} = 0$ and $K(A, k)_k = A$. For $i > k$ we have $N(K(A, k))_i = 0$ by Lemma 1.9 and the definition of $K(A, k)$, see Definition 1.1. □

Lemma 1.3. *The functor N is exact.*

Proof. By Lemma 1.8 and the functorial decomposition of Lemma 1.4. □

Lemma 1.4. *Let $\mathcal{A}$ be an abelian category. Let U be a simplicial object of $\mathcal{A}$. The canonical morphism of chain complexes $N(U) \to s(U)$ is split. In fact,*

$$s(U) = N(U) \oplus D(U)$$

for some complex $D(U)$. The construction $U \mapsto D(U)$ is functorial.

Proof. Define $D(U)_n$ to be the image of

$$\bigoplus\nolimits_{\varphi:[n]\to[m] \text{ surjective, } m<n} N(U_m) \xrightarrow{\bigoplus U(\varphi)} U_n$$

which is a subobject of U_n complementary to $N(U_n)$ according to Lemma 1.11 and Definition 1.10. We show that $D(U)$ is a subcomplex. Pick a surjective map $\varphi : [n] \to [m]$ with $m < n$ and consider the composition

$$N(U_m) \xrightarrow{U(\varphi)} U_n \xrightarrow{d_n} U_{n-1}.$$

This composition is the sum of the maps

$$N(U_m) \xrightarrow{U(\varphi\circ\delta_i^n)} U_{n-1}$$

with sign $(-1)^i$, $i = 0, \ldots, n$.

First we will prove by ascending induction on m, $0 \leq m < n-1$ that all the maps $U(\varphi \circ \delta_i^n)$ map $N(U_m)$ into $D(U)_{n-1}$. (The case $m = n-1$ is treated below.) Whenever the map $\varphi \circ \delta_i^n : [n-1] \to [m]$ is surjective then the image of $N(U_m)$ under $U(\varphi \circ \delta_i^n)$ is contained in $D(U)_{n-1}$ by definition. If $\varphi \circ \delta_i^n : [n-1] \to [m]$ is not surjective, set $j = \varphi(i)$ and observe that i is the unique index whose image under φ is j. We may write $\varphi \circ \delta_i^n = \delta_j^m \circ \psi \circ \delta_i^n$ for some $\psi : [n-1] \to [m-1]$. Hence $U(\varphi \circ \delta_i^n) = U(\psi \circ \delta_i^n) \circ d_j^m$ which is zero on $N(U_m)$ unless $j = m$. If $j = m$, then $d_m^m(N(U_m)) \subset N(U_{m-1})$ and hence $U(\varphi \circ \delta_i^n)(N(U_m)) \subset U(\psi \circ \delta_i^n)(N(U_{m-1}))$ and we win by induction hypothesis.

To finish proving that $D(U)$ is a subcomplex we still have to deal with the composition

$$N(U_m) \xrightarrow{U(\varphi)} U_n \xrightarrow{d_n} U_{n-1}.$$

in case $m = n-1$. In this case $\varphi = \sigma_j^{n-1}$ for some $0 \leq j \leq n-1$ and $U(\varphi) = s_j^{n-1}$. Thus the composition is given by the sum

$$\sum (-1)^i d_i^n \circ s_j^{n-1}$$

Recall from Remark 1.12 that $d_j^n \circ s_j^{n-1} = d_{j+1}^n \circ s_j^{n-1} = \mathrm{id}$ and these drop out because the corresponding terms have opposite signs. The map $d_n^n \circ s_j^{n-1}$, if $j < n-1$, is equal to $s_j^{n-2} \circ d_{n-1}^{n-1}$. Since d_{n-1}^{n-1} maps $N(U_{n-1})$ into $N(U_{n-2})$, we see that the image $d_n^n(s_j^{n-1}(N(U_{n-1}))$ is contained in $s_j^{n-2}(N(U_{n-2}))$ which is contained in $D(U_{n-1})$ by definition. For all other combinations of (i, j) we have either $d_i^n \circ s_j^{n-1} = s_{j-1}^{n-2} \circ d_i^{n-1}$ (if $i < j$), or $d_i^n \circ s_j^{n-1} = s_j^{n-2} \circ d_{i-1}^{n-1}$ (if $n > i > j+1$) and in these cases the map is zero because of the definition of $N(U_{n-1})$. □

Lemma 1.5. *With A, k and U as above, so $U_i = 0$, $i < k$ and $U_k = A$.*

(1) *Given a k-truncated simplicial object V we have*

$$\mathrm{Mor}(U,V) = \{f : A \to V_k \mid d_i^k \circ f = 0,\ i = 0,\ldots,k\}$$

and

$$\mathrm{Mor}(V,U) = \{f : V_k \to A \mid f \circ s_i^{k-1} = 0,\ i = 0,\ldots,k-1\}.$$

(2) *The object $i_{k!}U$ has nth term equal to $\bigoplus_\alpha A$ where α runs over all surjective morphisms $\alpha : [n] \to [k]$.*

(3) *For any $\varphi : [m] \to [n]$ the map $i_{k!}U(\varphi)$ is described as the mapping $\bigoplus_\alpha A \to \bigoplus_{\alpha'} A$ which maps to component corresponding to $\alpha : [n] \to [k]$ to zero if $\alpha \circ \varphi$ is not surjective and by the identity to the component corresponding to $\alpha \circ \varphi$ if it is surjective.*

(4) *The object $\mathrm{cosk}_k U$ has nth term equal to $\bigoplus_\beta A$, where β runs over all injective morphisms $\beta : [k] \to [n]$.*

(5) *For any $\varphi : [m] \to [n]$ the map $\mathrm{cosk}_k U(\varphi)$ is described as the mapping $\bigoplus_\beta A \to \bigoplus_{\beta'} A$ which maps to component corresponding to $\beta : [k] \to [n]$ to zero if β does not factor through φ and by the identity to each of the components corresponding to β' such that $\beta = \varphi \circ \beta'$ if it does.*

(6) *The canonical map $c : i_{k!}U \to \mathrm{cosk}_k U$ in degree n has (α,β) coefficient $A \to A$ equal to zero if $\alpha \circ \beta$ is not the identity and equal to id_A if it is.*

(7) *The canonical map $c : i_{k!}U \to \mathrm{cosk}_k U$ is injective.*

Proof. The proof of (1) is left to the reader.

Let us take the rules of (2) and (3) as the definition of a simplicial object, call it $\tilde{U}$. We will show that it is an incarnation of $i_{k!}U$. This will prove (2), (3) at the same time. We have to show that given a morphism $f : U \to \mathrm{sk}_k V$ there exists a unique morphism $\tilde{f} : \tilde{U} \to V$ which recovers f upon taking the k-skeleton. From (1) we see that f corresponds with a morphism $f_k : A \to V_k$ which maps into the kernel of d_i^k for all i. For any surjective $\alpha : [n] \to [k]$ we set $\tilde{f}_\alpha : A \to V_n$ equal to the composition $\tilde{f}_\alpha = V(\alpha) \circ f_k : A \to V_n$. We define $\tilde{f}_n : \tilde{U}_n \to V_n$ as the sum of the $\tilde{f}_\alpha$ over $\alpha : [n] \to [k]$ surjective. Such a collection of $\tilde{f}_\alpha$ defines a morphism of simplicial objects if and only if for any $\varphi : [m] \to [n]$ the diagram

$$\begin{array}{ccc} \bigoplus_{\alpha:[n]\to[k] \text{ surjective}} A & \xrightarrow[\tilde{f}_n]{} & V_n \\ {\scriptstyle (3)}\downarrow & & \downarrow{\scriptstyle V(\varphi)} \\ \bigoplus_{\alpha':[m]\to[k] \text{ surjective}} A & \xrightarrow{\tilde{f}_m} & V_m \end{array}$$

is commutative. Choosing $\varphi = \alpha$ shows our choice of $\tilde{f}_\alpha$ is uniquely determined by f_k. The commutativity in general may be checked for each summand of the left upper corner separately. It is clear for the summands corresponding to α where $\alpha \circ \varphi$ is surjective, because those get mapped by id_A to the summand with $\alpha' = \alpha \circ \varphi$, and we have $\tilde{f}_{\alpha'} = V(\alpha') \circ f_k = V(\alpha \circ \varphi) \circ f_k = V(\varphi) \circ \tilde{f}_\alpha$. For those where $\alpha \circ \varphi$ is not surjective, we have to show that $V(\varphi) \circ \tilde{f}_\alpha = 0$. By definition this is equal to $V(\varphi) \circ V(\alpha) \circ f_k = V(\alpha \circ \varphi) \circ f_k$. Since $\alpha \circ \varphi$ is not surjective we can write it as $\delta_i^k \circ \psi$, and we deduce that $V(\varphi) \circ V(\alpha) \circ f_k = V(\psi) \circ d_i^k \circ f_k = 0$ see above.

Let us take the rules of (4) and (5) as the definition of a simplicial object, call it $\tilde{U}$. We will show that it is an incarnation of $\mathrm{cosk}_k U$. This will prove (4), (5) at the same time. The argument is completely dual to the proof of (2), (3) above, but we

give it anyway. We have to show that given a morphism $f : \mathrm{sk}_k V \to U$ there exists a unique morphism $\tilde{f} : V \to \tilde{U}$ which recovers f upon taking the k-skeleton. From (1) we see that f corresponds with a morphism $f_k : V_k \to A$ which is zero on the image of s_i^{k-1} for all i. For any injective $\beta : [k] \to [n]$ we set $\tilde{f}_\beta : V_n \to A$ equal to the composition $\tilde{f}_\beta = f_k \circ V(\beta) : V_n \to A$. We define $\tilde{f}_n : V_n \to \tilde{U}_n$ as the sum of the $\tilde{f}_\beta$ over $\beta : [k] \to [n]$ injective. Such a collection of $\tilde{f}_\beta$ defines a morphism of simplicial objects if and only if for any $\varphi : [m] \to [n]$ the diagram

$$\begin{array}{ccc} V_n & \xrightarrow[\tilde{f}_n]{} & \bigoplus_{\beta:[k]\to[n] \text{ injective}} A \\ {\scriptstyle V(\varphi)}\downarrow & & \downarrow{\scriptstyle (5)} \\ V_m & \xrightarrow{\tilde{f}_m} & \bigoplus_{\beta':[k]\to[m] \text{ injective}} A \end{array}$$

is commutative. Choosing $\varphi = \beta$ shows our choice of $\tilde{f}_\beta$ is uniquely determined by f_k. The commutativity in general may be checked for each summand of the right lower corner separately. It is clear for the summands corresponding to β' where $\varphi \circ \beta'$ is injective, because these summands get mapped into by exactly the summand with $\beta = \varphi \circ \beta'$ and we have in that case $\tilde{f}_{\beta'} \circ V(\varphi) = f_k \circ V(\beta') \circ V(\varphi) = f_k \circ V(\beta) = \tilde{f}_\beta$. For those where $\varphi \circ \beta'$ is not injective, we have to show that $\tilde{f}_{\beta'} \circ V(\varphi) = 0$. By definition this is equal to $f_k \circ V(\beta') \circ V(\varphi) = f_k \circ V(\varphi \circ \beta')$. Since $\varphi \circ \beta'$ is not injective we can write it as $\psi \circ \sigma_i^{k-1}$, and we deduce that $f_k \circ V(\beta') \circ V(\varphi) = f_k \circ s_i^{k-1} \circ V(\psi) = 0$ see above.

The composition $i_{k!}U \to \mathrm{cosk}_k U$ is the unique map of simplicial objects which is the identity on $A = U_k = (i_{k!}U)_k = (\mathrm{cosk}_k U)_k$. Hence it suffices to check that the proposed rule defines a morphism of simplicial objects. To see this we have to show that for any $\varphi : [m] \to [n]$ the diagram

$$\begin{array}{ccc} \bigoplus_{\alpha:[n]\to[k] \text{ surjective}} A & \xrightarrow[(6)]{} & \bigoplus_{\beta:[k]\to[n] \text{ injective}} A \\ {\scriptstyle (3)}\downarrow & & \downarrow{\scriptstyle (5)} \\ \bigoplus_{\alpha':[m]\to[k] \text{ surjective}} A & \xrightarrow{(6)} & \bigoplus_{\beta':[k]\to[m] \text{ injective}} A \end{array}$$

is commutative. Now we can think of this in terms of matrices filled with only 0's and 1's as follows: The matrix of (3) has a nonzero (α', α) entry if and only if $\alpha' = \alpha \circ \varphi$. Likewise the matrix of (5) has a nonzero (β', β) entry if and only if $\beta = \varphi \circ \beta'$. The upper matrix of (6) has a nonzero (α, β) entry if and only if $\alpha \circ \beta = \mathrm{id}_{[k]}$. Similarly for the lower matrix of (6). The commutativity of the diagram then comes down to computing the (α, β') entry for both compositions and seeing they are equal. This comes down to the following equality

$$\#\left\{\beta \mid \beta = \varphi \circ \beta' \text{ and } \alpha \circ \beta = \mathrm{id}_{[k]}\right\} = \#\left\{\alpha' \mid \alpha' = \alpha \circ \varphi \text{ and } \alpha' \circ \beta' = \mathrm{id}_{[k]}\right\}$$

whose proof may safely be left to the reader.

Finally, we prove (7). This follows directly from Lemmas 1.13, 1.14, 1.15 and 1.9. □

Lemma 1.6. *Let $\mathcal{A}$ be an abelian category. Let A be an object of $\mathcal{A}$ and let k be an integer. We have $H_i(s(K(A, k))) = A$ if $i = k$ and 0 else.*

Proof. First, let us prove this if $k = 0$. In this case we have $K(A,0)_n = A$ for all n. Furthermore, all the maps in this simplicial abelian group are id_A, in other words $K(A,0)$ is the constant simplicial object with value A. The boundary maps $d_n = \sum_{i=0}^{n}(-1)^i \mathrm{id}_A = 0$ if n odd and $= \mathrm{id}_A$ if n is even. Thus $s(K(A,0))$ looks like this

$$\ldots \to A \xrightarrow{0} A \xrightarrow{1} A \xrightarrow{0} A \to 0$$

and the result is clear.

Next, we prove the result for all k by induction. Given the result for k consider the short exact sequence

$$0 \to K(A,k) \to E \to K(A,k+1) \to 0$$

from Lemma 1.17. By Lemma 1.16 the associated sequence of chain complexes is exact. By Lemma 1.18 we see that $s(E)$ is acyclic. Hence the result for $k+1$ follows from the long exact sequence of homology, see Homology, Lemma 0111. □

Lemma 1.7. *Let $\mathcal{A}$ be an abelian category. Let U be a simplicial object of $\mathcal{A}$. Let $n \geq 0$ be an integer. The morphism $i_{n!}sk_nU \to U$ identifies $i_{n!}sk_nU$ with the simplicial subobject $U' \subset U$ defined in Lemma 1.19.*

Proof. By Lemma 1.9 we have $N(i_{n!}\mathrm{sk}_n U)_i = 0$ for $i > n$. The map $i_{n!}\mathrm{sk}_n U \to U$ is an isomorphism in degrees $\leq n$, see Lemma 1.15. Combined we conclude that the map $i_{n!}\mathrm{sk}_n U \to U$ induces injective maps $N(i_{n!}\mathrm{sk}_n U)_i \to N(U)_i$ for all i. Hence Lemma 1.13 applies. Thus $i_{n!}\mathrm{sk}_n U \to U$ is injective. The result follows easily from this. □

Lemma 1.8. *The functor s is exact.*

Proof. Clear from Lemma 1.16. □

Lemma 1.9. *Let $\mathcal{A}$ be an abelian category Let U be an m-truncated simplicial object of $\mathcal{A}$. For $n > m$ we have $N(i_{m!}U)_n = 0$.*

Proof. Write $V = i_{m!}U$. Let $V' \subset V$ be the simplicial subobject of V with $V'_i = V_i$ for $i \leq m$ and $N(V'_i) = 0$ for $i > m$, see Lemma 1.19. By the adjunction formula, since $\mathrm{sk}_m V' = U$, there is an inverse to the injection $V' \to V$. Hence $V' = V$. □

Definition 1.10. Let $\mathcal{C}$ be a category which admits finite nonempty coproducts. We say a simplicial object U of $\mathcal{C}$ is *split* if there exist subobjects $N(U_m)$ of U_m, $m \geq 0$ with the property that

$$\coprod_{\varphi:[n]\to[m] \text{ surjective}} N(U_m) \longrightarrow U_n \tag{1.10.1}$$

is an isomorphism for all $n \geq 0$. If U is an r-truncated simplicial object of $\mathcal{C}$ then we say U is *split* if there exist subobjects $N(U_m)$ of U_m, $r \geq m \geq 0$ with the property that (1.10.1) is an isomorphism for $r \geq n \geq 0$.

Lemma 1.11. *Let $\mathcal{A}$ be an abelian category. Let U be a simplicial object in $\mathcal{A}$. Then U has a splitting obtained by taking $N(U_0) = U_0$ and for $m \geq 1$ taking*

$$N(U_m) = \bigcap\nolimits_{i=0}^{m-1} \mathrm{Ker}(d_i^m).$$

Moreover, this splitting is functorial on the category of simplicial objects of $\mathcal{A}$.

Proof. For any object A of $\mathcal{A}$ we obtain a simplicial abelian group $\mathrm{Mor}_{\mathcal{A}}(A, U)$. Each of these are canonically split by Lemma 1.20. Moreover,

$$N(\mathrm{Mor}_{\mathcal{A}}(A, U_m)) = \bigcap\nolimits_{i=0}^{m-1} \mathrm{Ker}(d_i^m) = \mathrm{Mor}_{\mathcal{A}}(A, N(U_m)).$$

Hence we see that the morphism (1.10.1) becomes an isomorphism after applying the functor $\mathrm{Mor}_{\mathcal{A}}(A, -)$ for any object of $\mathcal{A}$. Hence it is an isomorphism by the Yoneda lemma. □

Remark 1.12. By abuse of notation we sometimes write $d_i : U_n \to U_{n-1}$ instead of d_i^n, and similarly for $s_i : U_n \to U_{n+1}$. The relations among the morphisms d_i^n and s_i^n may be expressed as follows:

(1) If $i < j$, then $d_i \circ d_j = d_{j-1} \circ d_i$.
(2) If $i < j$, then $d_i \circ s_j = s_{j-1} \circ d_i$.
(3) We have $\mathrm{id} = d_j \circ s_j = d_{j+1} \circ s_j$.
(4) If $i > j + 1$, then $d_i \circ s_j = s_j \circ d_{i-1}$.
(5) If $i \leq j$, then $s_i \circ s_j = s_{j+1} \circ s_i$.

This means that whenever the compositions on both the left and the right are defined then the corresponding equality should hold.

Lemma 1.13. *Let $\mathcal{A}$ be an abelian category. Let $f : U \to V$ be a morphism of simplicial objects of $\mathcal{A}$. If the induced morphisms $N(f)_i : N(U)_i \to N(V)_i$ are injective for all i, then f_i is injective for all i. Same holds with "injective" replaced with "surjective", or "isomorphism".*

Proof. This is clear from Lemma 1.11 and the definition of a splitting. □

Lemma 1.14. *Let $\mathcal{C}$ be a category with finite limits. Let U be an n-truncated simplicial object of $\mathcal{C}$. The morphism $sk_n cosk_n U \to U$ is an isomorphism.*

Proof. Combine Lemmas 1.21 and 1.22. □

Lemma 1.15. *Let $\mathcal{C}$ be a category which has finite colimits. Let U be an m-truncated simplicial object of $\mathcal{C}$. The map $U \to sk_m i_{m!} U$ is an isomorphism.*

Proof. Combine Lemmas 1.23 and 1.24. □

Lemma 1.16. *Let $\mathcal{A}$ be an abelian category.*

(1) *The categories $Simp(\mathcal{A})$ and $CoSimp(\mathcal{A})$ are abelian.*
(2) *A morphism of (co)simplicial objects $f : A \to B$ is injective if and only if each $f_n : A_n \to B_n$ is injective.*
(3) *A morphism of (co)simplicial objects $f : A \to B$ is surjective if and only if each $f_n : A_n \to B_n$ is surjective.*
(4) *A sequence of (co)simplicial objects*

$$A \xrightarrow{f} B \xrightarrow{g} C$$

is exact at B if and only if each sequence

$$A_i \xrightarrow{f_i} B_i \xrightarrow{g_i} C_i$$

is exact at B_i.

Proof. Pre-additivity is easy. A final object is given by $U_n = 0$ in all degrees. Existence of direct products we saw in Lemmas 1.25 and 1.26. Kernels and cokernels are obtained by taking termwise kernels and cokernels. □

Lemma 1.17. *Let $\mathcal{A}$ be an abelian category. Let A be an object of $\mathcal{A}$ and let k be an integer ≥ 0. Consider the simplicial object E defined by the following rules*

1. *$E_n = \bigoplus_\alpha A$, where the sum is over $\alpha : [n] \to [k+1]$ whose image is either $[k]$ or $[k+1]$.*
2. *Given $\varphi : [m] \to [n]$ the map $E_n \to E_m$ maps the summand corresponding to α via id_A to the summand corresponding to $\alpha \circ \varphi$, provided $\mathrm{Im}(\alpha \circ \varphi)$ is equal to $[k]$ or $[k+1]$.*

Then there exists a short exact sequence

$$0 \to K(A, k) \to E \to K(A, k+1) \to 0$$

which is term by term split exact.

Proof. The maps $K(A,k)_n \to E_n$ resp. $E_n \to K(A,k+1)_n$ are given by the inclusion of direct sums, resp. projection of direct sums which is obvious from the inclusions of index sets. It is clear that these are maps of simplicial objects. □

Lemma 1.18. *Let $\mathcal{A}$ be an abelian category. Let A be an object of $\mathcal{A}$ and let k be an integer. Let E be the object described in Lemma 1.17. Then the complex $s(E)$ is acyclic.*

Proof. For a morphism $\alpha : [n] \to [k+1]$ we define $\alpha' : [n+1] \to [k+1]$ to be the map such that $\alpha'|_{[n]} = \alpha$ and $\alpha'(n+1) = k+1$. Note that if the image of α is $[k]$ or $[k+1]$, then the image of α' is $[k+1]$. Consider the family of maps $h_n : E_n \to E_{n+1}$ which maps the summand corresponding to α to the summand corresponding to α' via the identity on A. Let us compute $d_{n+1} \circ h_n - h_{n-1} \circ d_n$. We will first do this in case the category $\mathcal{A}$ is the category of abelian groups. Let us use the notation x_α to indicate the element $x \in A$ in the summand of E_n corresponding to the map α occurring in the index set. Let us also adopt the convention that x_α designates the zero element of E_n whenever $\mathrm{Im}(\alpha)$ is not $[k]$ or $[k+1]$. With these conventions we see that

$$d_{n+1}(h_n(x_\alpha)) = \sum\nolimits_{i=0}^{n+1} (-1)^i x_{\alpha' \circ \delta_i^{n+1}}$$

and

$$h_{n-1}(d_n(x_\alpha)) = \sum\nolimits_{i=0}^{n} (-1)^i x_{(\alpha \circ \delta_i^n)'}$$

It is easy to see that $\alpha' \circ \delta_i^{n+1} = (\alpha \circ \delta_i^n)'$ for $i = 0, \ldots, n$. It is also easy to see that $\alpha' \circ \delta_{n+1}^{n+1} = \alpha$. Thus we see that

$$(d_{n+1} \circ h_n - h_{n-1} \circ d_n)(x_\alpha) = (-1)^{n+1} x_\alpha$$

These identities continue to hold if $\mathcal{A}$ is any abelian category because they hold in the simplicial abelian group $[n] \mapsto \mathrm{Hom}(A, E_n)$; details left to the reader. We conclude that the identity map on E is homotopic to zero, with homotopy given by the system of maps $h'_n = (-1)^{n+1} h_n : E_n \to E_{n+1}$. Hence we see that E is acyclic, for example by Homology, Lemma 0110. □

Lemma 1.19. *Let $\mathcal{A}$ be an abelian category. Let U be a simplicial object of $\mathcal{A}$. Let $n \geq 0$ be an integer. The rule*

$$U'_m = \sum\nolimits_{\varphi : [m] \to [i],\ i \leq n} \mathrm{Im}(U(\varphi))$$

defines a sub simplicial object $U' \subset U$ with $U'_i = U_i$ for $i \leq n$. Moreover, $N(U'_m) = 0$ for all $m > n$.

Proof. Pick m, $i \leq n$ and some $\varphi : [m] \to [i]$. The image under $U(\psi)$ of $\mathrm{Im}(U(\varphi))$ for any $\psi : [m'] \to [m]$ is equal to the image of $U(\varphi \circ \psi)$ and $\varphi \circ \psi : [m'] \to [i]$. Hence U' is a simplicial object. Pick $m > n$. We have to show $N(U'_m) = 0$. By definition of $N(U_m)$ and $N(U'_m)$ we have $N(U'_m) = U'_m \cap N(U_m)$ (intersection of subobjects). Since U is split by Lemma 1.11, it suffices to show that U'_m is contained in the sum

$$\sum\nolimits_{\varphi:[m]\to[m'] \text{ surjective, } m'<m} \mathrm{Im}(U(\varphi)|_{N(U_{m'})}).$$

By the splitting each $U_{m'}$ is the sum of images of $N(U_{m''})$ via $U(\psi)$ for surjective maps $\psi : [m'] \to [m'']$. Hence the displayed sum above is the same as

$$\sum\nolimits_{\varphi:[m]\to[m'] \text{ surjective, } m'<m} \mathrm{Im}(U(\varphi)).$$

Clearly U'_m is contained in this by the simple fact that any $\varphi : [m] \to [i]$, $i \leq n$ occurring in the definition of U'_m may be factored as $[m] \to [m'] \to [i]$ with $[m] \to [m']$ surjective and $m' < m$ as in the last displayed sum above. □

Lemma 1.20. *Let U be a simplicial abelian group. Then U has a splitting obtained by taking $N(U_0) = U_0$ and for $m \geq 1$ taking*

$$N(U_m) = \bigcap\nolimits_{i=0}^{m-1} \mathrm{Ker}(d_i^m).$$

Moreover, this splitting is functorial on the category of simplicial abelian groups.

Proof. By induction on n we will show that the choice of $N(U_m)$ in the lemma guarantees that (1.10.1) is an isomorphism for $m \leq n$. This is clear for $n = 0$. In the rest of this proof we are going to drop the superscripts from the maps d_i and s_i in order to improve readability. We will also repeatedly use the relations from Remark 1.12.

First we make a general remark. For $0 \leq i \leq m$ and $z \in U_m$ we have $d_i(s_i(z)) = z$. Hence we can write any $x \in U_{m+1}$ uniquely as $x = x' + x''$ with $d_i(x') = 0$ and $x'' \in \mathrm{Im}(s_i)$ by taking $x' = (x - s_i(d_i(x)))$ and $x'' = s_i(d_i(x))$. Moreover, the element $z \in U_m$ such that $x'' = s_i(z)$ is unique because s_i is injective.

Here is a procedure for decomposing any $x \in U_{n+1}$. First, write $x = x_0 + s_0(z_0)$ with $d_0(x_0) = 0$. Next, write $x_0 = x_1 + s_1(z_1)$ with $d_1(x_1) = 0$. Continue like this to get

$$\begin{aligned} x &= x_0 + s_0(z_0), \\ x_0 &= x_1 + s_1(z_1), \\ x_1 &= x_2 + s_2(z_2), \\ \ldots &\quad \ldots \quad \ldots \\ x_{n-1} &= x_n + s_n(z_n) \end{aligned}$$

where $d_i(x_i) = 0$ for all $i = n, \ldots, 0$. By our general remark above all of the x_i and z_i are determined uniquely by x. We claim that $x_i \in \mathrm{Ker}(d_0) \cap \mathrm{Ker}(d_1) \cap \ldots \cap \mathrm{Ker}(d_i)$ and $z_i \in \mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})$ for $i = n, \ldots, 0$. Here and in the following an empty intersection of kernels indicates the whole space; i.e., the notation $z_0 \in \mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})$ when $i = 0$ means $z_0 \in U_n$ with no restriction.

We prove this by ascending induction on i. It is clear for $i = 0$ by construction of x_0 and z_0. Let us prove it for $0 < i \leq n$ assuming the result for $i - 1$. First of all we

have $d_i(x_i) = 0$ by construction. So pick a j with $0 \leq j < i$. We have $d_j(x_{i-1}) = 0$ by induction. Hence

$$0 = d_j(x_{i-1}) = d_j(x_i) + d_j(s_i(z_i)) = d_j(x_i) + s_{i-1}(d_j(z_i)).$$

The last equality by the relations of Remark 1.12. These relations also imply that $d_{i-1}(d_j(x_i)) = d_j(d_i(x_i)) = 0$ because $d_i(x_i) = 0$ by construction. Then the uniqueness in the general remark above shows the equality $0 = x' + x'' = d_j(x_i) + s_{i-1}(d_j(z_i))$ can only hold if both terms are zero. We conclude that $d_j(x_i) = 0$ and by injectivity of s_{i-1} we also conclude that $d_j(z_i) = 0$. This proves the claim.

The claim implies we can uniquely write

$$x = s_0(z_0) + s_1(z_1) + \ldots + s_n(z_n) + x_n$$

with $x_n \in N(U_{n+1})$ and $z_i \in \mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})$. We can reformulate this as saying that we have found a direct sum decomposition

$$U_{n+1} = N(U_{n+1}) \oplus \bigoplus\nolimits_{i=0}^{i=n} s_i\Big(\mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})\Big)$$

with the property that

$$\mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_j) = N(U_{n+1}) \oplus \bigoplus\nolimits_{i=j+1}^{i=n} s_i\Big(\mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{i-1})\Big)$$

for $j = 0, \ldots, n$. The result follows from this statement as follows. Each of the z_i in the expression for x can be written uniquely as

$$z_i = s_i(z'_{i,i}) + \ldots + s_{n-1}(z'_{i,n-1}) + z_{i,0}$$

with $z_{i,0} \in N(U_n)$ and $z'_{i,j} \in \mathrm{Ker}(d_0) \cap \ldots \cap \mathrm{Ker}(d_{j-1})$. The first few steps in the decomposition of z_i are zero because z_i already is in the kernel of $d_0, \ldots, d_i$. This in turn uniquely gives

$$x = x_n + s_0(z_{0,0}) + s_1(z_{1,0}) + \ldots + s_n(z_{n,0}) + \sum\nolimits_{0 \leq i \leq j \leq n-1} s_i(s_j(z'_{i,j})).$$

Continuing in this fashion we see that we in the end obtain a decomposition of x as a sum of terms of the form

$$s_{i_1} s_{i_2} \ldots s_{i_k}(z)$$

with $0 \leq i_1 \leq i_2 \leq \ldots \leq i_k \leq n - k + 1$ and $z \in N(U_{n+1-k})$. This is exactly the required decomposition, because any surjective map $[n+1] \to [n+1-k]$ can be uniquely expressed in the form

$$\sigma^{n-k}_{i_k} \ldots \sigma^{n-1}_{i_2} \sigma^n_{i_1}$$

with $0 \leq i_1 \leq i_2 \leq \ldots \leq i_k \leq n - k + 1$. □

Lemma 1.21. *If the category $\mathcal{C}$ has finite limits, then $cosk_m$ functors exist for all m. Moreover, for any m-truncated simplicial object U the simplicial object $cosk_m U$ is described by the formula*

$$(cosk_m U)_n = \lim\nolimits_{(\Delta/[n])^{opp}_{\leq m}} U(n)$$

and for $\varphi : [n] \to [n']$ the map $cosk_m U(\varphi)$ comes from the identification $U(n') \circ \overline{\varphi} = U(n)$ above via Categories, Lemma 002L.

Proof. During the proof of this lemma we denote $\mathrm{cosk}_m U$ the simplicial object with $(\mathrm{cosk}_m U)_n$ equal to $\lim_{(\Delta/[n])^{opp}_{\leq m}} U(n)$. We will conclude at the end of the proof that it does satisfy the required mapping property.

Suppose that V is a simplicial object. A morphism $\gamma : V \to \mathrm{cosk}_m U$ is given by a sequence of morphisms $\gamma_n : V_n \to (\mathrm{cosk}_m U)_n$. By definition of a limit, this is given by a collection of morphisms $\gamma(\alpha) : V_n \to U_k$ where α ranges over all $\alpha : [k] \to [n]$ with $k \leq m$. These morphisms then also satisfy the rules that

$$\begin{array}{ccc} V_n & \xrightarrow{\gamma(\alpha)} & U_k \\ {\scriptstyle V(\varphi)}\uparrow & & \uparrow{\scriptstyle U(\psi)} \\ V_{n'} & \xrightarrow{\gamma(\alpha')} & U_{k'} \end{array}$$

are commutative, given any $0 \leq k, k' \leq m$, $0 \leq n, n'$ and any $\psi : [k] \to [k']$, $\varphi : [n] \to [n']$, $\alpha : [k] \to [n]$ and $\alpha' : [k'] \to [n']$ in Δ such that $\varphi \circ \alpha = \alpha' \circ \psi$. Taking $n = k = k'$, $\varphi = \alpha'$, and $\alpha = \psi = \mathrm{id}_{[k]}$ we deduce that $\gamma(\alpha') = \gamma(\mathrm{id}_{[k]}) \circ V(\alpha')$. In other words, the morphisms $\gamma(\mathrm{id}_{[k]})$, $k \leq m$ determine the morphism γ. And it is easy to see that these morphisms form a morphism $\mathrm{sk}_m V \to U$.

Conversely, given a morphism $\gamma : \mathrm{sk}_m V \to U$, we obtain a family of morphisms $\gamma(\alpha)$ where α ranges over all $\alpha : [k] \to [n]$ with $k \leq m$ by setting $\gamma(\alpha) = \gamma(\mathrm{id}_{[k]}) \circ V(\alpha)$. These morphisms satisfy all the displayed commutativity restraints pictured above, and hence give rise to a morphism $V \to \mathrm{cosk}_m U$. □

Lemma 1.22. *Let $\mathcal{C}$ be a category. Let U be an m-truncated simplicial object of $\mathcal{C}$. For $n \leq m$ the limit $\lim_{(\Delta/[n])^{opp}_{\leq m}} U(n)$ exists and is canonically isomorphic to U_n.*

Proof. This is true because the category $(\Delta/[n])_{\leq m}$ has an final object in this case, namely the identity map $[n] \to [n]$. □

Lemma 1.23. *Let $\mathcal{C}$ be a category which has finite colimits. The functors $i_{m!}$ exist for all m. Let U be an m-truncated simplicial object of $\mathcal{C}$. The simplicial object $i_{m!}U$ is described by the formula*

$$(i_{m!}U)_n = \mathrm{colim}_{([n]/\Delta)^{opp}_{\leq m}} U(n)$$

and for $\varphi : [n] \to [n']$ the map $i_{m!}U(\varphi)$ comes from the identification $U(n) \circ \underline{\varphi} = U(n')$ above via Categories, Lemma 002K.

Proof. In this proof we denote $i_{m!}U$ the simplicial object whose nth term is given by the displayed formula of the lemma. We will show it satisfies the adjointness property.

Let V be a simplicial object of $\mathcal{C}$. Let $\gamma : U \to \mathrm{sk}_m V$ be given. A morphism

$$\mathrm{colim}_{([n]/\Delta)^{opp}_{\leq m}} U(n) \to T$$

is given by a compatible system of morphisms $f_\alpha : U_k \to T$ where $\alpha : [n] \to [k]$ with $k \leq m$. Certainly, we have such a system of morphisms by taking the compositions

$$U_k \xrightarrow{\gamma_k} V_k \xrightarrow{V(\alpha)} V_n.$$

Hence we get an induced morphism $(i_{m!}U)_n \to V_n$. We leave it to the reader to see that these form a morphism of simplicial objects $\gamma' : i_{m!}U \to V$.

Conversely, given a morphism $\gamma' : i_{m!}U \to V$ we obtain a morphism $\gamma : U \to \mathrm{sk}_m V$ by setting $\gamma_i : U_i \to V_i$ equal to the composition

$$U_i \xrightarrow{\mathrm{id}_{[i]}} \mathrm{colim}_{([i]/\Delta)^{opp}_{\leq m}} U(i) \xrightarrow{\gamma'_i} V_i$$

for $0 \leq i \leq n$. We leave it to the reader to see that this is the inverse of the construction above. □

Lemma 1.24. *Let $\mathcal{C}$ be a category. Let U be an m-truncated simplicial object of $\mathcal{C}$. For any $n \leq m$ the colimit*

$$\mathrm{colim}_{([n]/\Delta)^{opp}_{\leq m}} U(n)$$

exists and is equal to U_n.

Proof. This is so because the category $([n]/\Delta)_{\leq m}$ has an initial object, namely $\mathrm{id} : [n] \to [n]$. □

Lemma 1.25. *If U and V are simplicial objects in the category $\mathcal{C}$, and if $U \times V$ exists, then we have*

$$\mathrm{Mor}(W, U \times V) = \mathrm{Mor}(W, U) \times \mathrm{Mor}(W, V)$$

for any third simplicial object W of $\mathcal{C}$.

Proof. Omitted. □

Lemma 1.26. *If U and V are cosimplicial objects in the category $\mathcal{C}$, and if $U \times V$ exists, then we have*

$$\mathrm{Mor}(W, U \times V) = \mathrm{Mor}(W, U) \times \mathrm{Mor}(W, V)$$

for any third cosimplicial object W of $\mathcal{C}$.

Proof. Omitted. □

Lemma 1.27. *Let $f : X \to Y$ be a homomorphism of simplicial abelian groups which is termwise surjective and induces a quasi-isomorphism on associated chain complexes. Then f is a trivial Kan fibration of simplicial sets.*

Proof. Consider a commutative solid diagram

$$\begin{array}{ccc} \partial\Delta[n] & \xrightarrow{a} & X \\ \downarrow & \nearrow & \downarrow \\ \Delta[n] & \xrightarrow{b} & Y \end{array}$$

as in Definition 0.1. The map a corresponds to $x_0, \ldots, x_n \in X_{n-1}$ satisfying $d_i x_j = d_{j-1} x_i$ for $i < j$. The map b corresponds to an element $y \in Y_n$ such that $d_i y = f(x_i)$. Our task is to produce an $x \in X_n$ such that $d_i x = x_i$ and $f(x) = y$.

Since f is termwise surjective we can find $x \in X_n$ with $f(x) = y$. Replace y by $0 = y - f(x)$ and x_i by $x_i - d_i x$. Then we see that we may assume $y = 0$. In particular $f(x_i) = 0$. In other words, we can replace X by $\mathrm{Ker}(f) \subset X$ and Y by 0. This works, because by Homology, Lemma 0111 the homology of the chain complex associated to $\mathrm{Ker}(f)$ is zero and hence $\mathrm{Ker}(f) \to 0$ induces a quasi-isomorphism on associated chain complexes.

Since X is a Kan complex (Lemma 0.2) we can find $x \in X_n$ with $d_i x = x_i$ for $i = 0, \ldots, n-1$. After replacing x_i by $x_i - d_i x$ for $i = 0, \ldots, n$ we may assume that $x_0 = x_1 = \ldots = x_{n-1} = 0$. In this case we see that $d_i x_n = 0$ for $i = 0, \ldots, n-1$. Thus $x_n \in N(X)_{n-1}$ and lies in the kernel of the differential $N(X)_{n-1} \to N(X)_{n-2}$. Here $N(X)$ is the normalized chain complex associated to X, see Section 1. Since $N(X)$ is quasi-isomorphic to $s(X)$ (Lemma 0.3) and thus acyclic we find $x \in N(X_n)$ whose differential is x_n. This x answers the question posed by the lemma and we are done. $\square$