MITTAG-LEFFLER CONDITION FROM GRADED ACC ON BOUNDARY IMAGES

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied verbatim from the pinned Stacks Project chapter file cohomology.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. The counted span cites the lemmas recorded in assembly.external_references with canonical tags as navigation pointers.

Lemma 0.1. *Let I be an ideal of a ring A. Let X be a topological space. Let*

$$\ldots \to \mathcal{F}_3 \to \mathcal{F}_2 \to \mathcal{F}_1$$

be an inverse system of sheaves of A-modules on X such that $\mathcal{F}_n = \mathcal{F}_{n+1}/I^n\mathcal{F}_{n+1}$. Let $p \geq 0$. Assume

$$\bigoplus\nolimits_{n\geq 0} H^{p+1}(X, I^n\mathcal{F}_{n+1})$$

satisfies the ascending chain condition as a graded $\bigoplus_{n\geq 0} I^n/I^{n+1}$-module. Then the inverse system $M_n = H^p(X, \mathcal{F}_n)$ satisfies the Mittag-Leffler condition[1]*.*

Proof. Set $N_n = H^{p+1}(X, I^n\mathcal{F}_{n+1})$ and let $\delta_n : M_n \to N_n$ be the boundary map on cohomology coming from the short exact sequence $0 \to I^n\mathcal{F}_{n+1} \to \mathcal{F}_{n+1} \to \mathcal{F}_n \to 0$. Then $\bigoplus \mathrm{Im}(\delta_n) \subset \bigoplus N_n$ is a graded submodule. Namely, if $s \in M_n$ and $f \in I^m$, then we have a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & I^n\mathcal{F}_{n+1} & \longrightarrow & \mathcal{F}_{n+1} & \longrightarrow & \mathcal{F}_n & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle f} & & \downarrow{\scriptstyle f} & & \downarrow{\scriptstyle f} & & \\ 0 & \longrightarrow & I^{n+m}\mathcal{F}_{n+m+1} & \longrightarrow & \mathcal{F}_{n+m+1} & \longrightarrow & \mathcal{F}_{n+m} & \longrightarrow & 0 \end{array}$$

The middle vertical map is given by lifting a local section of $\mathcal{F}_{n+1}$ to a section of $\mathcal{F}_{n+m+1}$ and then multiplying by f; similarly for the other vertical arrows. We conclude that $\delta_{n+m}(fs) = f\delta_n(s)$. By assumption we can find $s_j \in M_{n_j}$, $j = 1, \ldots, N$ such that $\delta_{n_j}(s_j)$ generate $\bigoplus \mathrm{Im}(\delta_n)$ as a graded module. Let $n > c = \max(n_j)$. Let $s \in M_n$. Then we can find $f_j \in I^{n-n_j}$ such that $\delta_n(s) = \sum f_j\delta_{n_j}(s_j)$. We conclude that $\delta(s - \sum f_j s_j) = 0$, i.e., we can find $s' \in M_{n+1}$ mapping to $s - \sum f_j s_j$ in M_n. It follows that

$$\mathrm{Im}(M_{n+1} \to M_{n-c}) = \mathrm{Im}(M_n \to M_{n-c})$$

Namely, the elements $f_j s_j$ map to zero in M_{n-c}. This proves the lemma. □

Date: stacks-project:cohomology.tex:local-snapshot-20261011; GNU FDL 1.2 or later.

[1] In fact, there exists a $c \geq 0$ such that $\mathrm{Im}(M_n \to M_{n-c})$ is the stable image for all $n \geq c$.