DIAGONAL MAP K -> R SHEAFHOM(L, K OTIMES ^L L)

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied verbatim from the pinned Stacks Project chapter file cohomology.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. The counted span cites the lemmas recorded in assembly.external_references with canonical tags as navigation pointers.

Allowed standard prerequisites (editorial).

- Diagonal map of hom complexes. Cohomology of Sheaves, Lemma cohomology-lemma-diagonal. K^* -> SheafHom^*(L^*, Tot(K^* otimes L^*)).

Source reference. Original lemma 0A8Q: sources/stacks-chow/tags.txt, source lines 12923–12923. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Lemma 0.1. *Let $(X, \mathcal{O}_X)$ be a ringed space. Given K, L in $D(\mathcal{O}_X)$ there is a canonical morphism*

$$K \longrightarrow R\mathcal{H}om(L, K \otimes^{\mathbf{L}}_{\mathcal{O}_X} L)$$

in $D(\mathcal{O}_X)$ functorial in both K and L.

Proof. Choose a K-flat complex $\mathcal{K}^\bullet$ representing K and any complex of $\mathcal{O}_X$-modules $\mathcal{L}^\bullet$ representing L. Choose a K-injective complex $\mathcal{J}^\bullet$ and a quasi-isomorphism $\mathrm{Tot}(\mathcal{K}^\bullet \otimes_{\mathcal{O}_X} \mathcal{L}^\bullet) \to \mathcal{J}^\bullet$. Then we use

$$\mathcal{K}^\bullet \to \mathcal{H}om^\bullet(\mathcal{L}^\bullet, \mathrm{Tot}(\mathcal{K}^\bullet \otimes_{\mathcal{O}_X} \mathcal{L}^\bullet)) \to \mathcal{H}om^\bullet(\mathcal{L}^\bullet, \mathcal{J}^\bullet)$$

where the first map comes from Lemma 0A8Q. □

Date: stacks-project:cohomology.tex:local-snapshot-20261011; GNU FDL 1.2 or later.