LARGE-INTERSECTION AND HAUSDORFF LOWER BOUNDS FOR INHOMOGENEOUS IID RANDOM LIMSUP SETS

PERSSON, TOMAS

Editorial note. Only selected author theorem counted; complete original mathematical body and bibliography preserved. Native vector figures retained; publisher date/editor frontmatter omitted from rendering preamble and proofc class environment supplied where needed.

Editorial note. Web-conversion-only DefTralics block omitted, retaining the native PDF math-operator definition.

Editorial note. One native mathscr{D_n} group is rendered as mathscr{D}_n so the unchanged index n uses an available math font. Bibliography literal beta letters use ensuremath beta. Original source.tex and all mathematical symbols retained.

Allowed standard prerequisites (editorial).

- Standard functional analysis, measure theory and the author-cited classical theorems. Standard prior results stay cited; complete source-local proofs and definitions retained.

1. Introduction

There has recently been some attention given to shrinking targets and randomly generated limsup-sets. If $T\colon M \to M$ is a dynamical system and M is a metric space, then the sequence of balls $B(y, r_n)$, where $r_n \searrow 0$, is called a shrinking target, and one is interested in whether, given an $x \in M$, the orbit of x hits the target infinitely many times or not, that is whether $T^n(x) \in B(y, r_n)$ holds for infinitely many n or not. It is usually not possible to say anything interesting about this for general points x and y, but there are several results for "typical" x or y.

For instance, if μ is a T-invariant measure, then one can consider the sets

$$E(x, r_n) = \{\, y : T^n(x) \in B(y, r_n) \text{ for infinitely many } n \,\},$$
$$F(y, r_n) = \{\, x : T^n(x) \in B(y, r_n) \text{ for infinitely many } n \,\},$$

and try to say something about the sizes of these sets. Hill and Velani [15] studied sets of the form $F(y, r_n)$ when T is a rational map of the Riemann sphere and M is its Julia set on which T is expanding. They estimated the Hausdorff dimension of $F(y, r_n)$ when $r_n = e^{-\tau n}$ and calculated it when $r_n = |(T^n)'(x)|^{-\tau}$. Similar results have been proved for the Gauß-map by Li, Wang, Wu and Xu [19]. For β-transformations such results were obtained by Bugeaud and Wang [4], and by Bugeaud and Liao [3]. Aspenberg and Persson [1] obtained some results for piecewise expanding maps that are not necessarily Markov maps.

The Hausdorff dimension of sets of the form $E(x, r_n)$ when $T\colon x \mapsto 2x \mod 1$ was calculated by Fan, Schmeling and Troubetzkoy [13]. Liao and Seuret [17] considered the case when T is an expanding Markov map. Persson and Rams [23] considered more general piecewise expanding maps that are not necessarily Markov maps.

In this paper we shall study sets of the type $E(x, r_n)$. Since $T^n(x) \in B(y, r_n)$ if and only if $y \in B(T^n(x), r_n)$, we have

$$E(x, r_n) = \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} B(T^n(x), r_n) = \limsup_n B(T^n(x), r_n).$$

Hence, we are dealing with a set which is the limit superior of a sequence of balls $B(T^n(x), r_n)$. By Birkhoff's ergodic theorem, the centres of the balls are distributed according to the measure μ for μ-almost every x, and if the system (X, T, μ) is sufficiently fast mixing, then one might expect that for μ-almost every x, the sequence of balls behaves in a random way. It is therefore reasonable to expect that the Hausdorff dimension of $E(x, r_n)$ is typically the same as that of the set $\limsup B(x_n, r_n)$, where x_n are random points that are independent and distributed according to the measure μ.

This brings us to the randomly generated limsup-sets or random covers, studied in [14, 5, 16, 21, 11, 25, 6]. In this paper we will build on ideas from [21] to develop a new method for analysing the Hausdorff dimension

Date: doi:10.5802/ahl.15; CC BY 4.0.

of limsup-sets. In some sense, the idea is a development of the following (new) proof of the following classical lemma.

Lemma 1.1 (Frostman [12, Théorème 47.2]). *Suppose $E \subset \mathbb{R}^d$ and that μ is a measure with $\emptyset \neq \operatorname{supp}\mu \subset E$. If*

$$\iint |x-y|^{-s}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y) < \infty$$

then $\dim_{\mathrm{H}} E \geq s$.

Proof. We may assume that $\mu(E)$ is finite, since we may replace μ by a restriction to a set of finite measure. Since $\iint |x-y|^{-s}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y) < \infty$, the function $x \mapsto \int |x-y|^{-s}\,\mathrm{d}\mu(y)$ is finite μ-almost everywhere, and it is positive on a set of positive measure. Hence

$$\nu(A) = \int_A \left(\int |x-y|^{-s}\,\mathrm{d}\mu(y) \right)^{-1} \mathrm{d}\mu(x)$$

defines a measure ν with $\nu(E) > 0$ and $\operatorname{supp}\nu \subset E$. The measure ν satisfies $\nu(U) \leq |U|^s$ for any set U, where $|U|$ denotes the diameter of U. This is proved using Jensen's inequality in the following way.

$$\begin{aligned}
\nu(U) &= \int_U \left(\int |x-y|^{-s}\,\mathrm{d}\mu(y) \right)^{-1} \mathrm{d}\mu(x) \\
&\leq \int_U \left(\int_U |x-y|^{-s}\,\frac{\mathrm{d}\mu(y)}{\mu(U)} \right)^{-1} \frac{\mathrm{d}\mu(x)}{\mu(U)} \\
&\leq \int_U \int_U |x-y|^{s}\,\frac{\mathrm{d}\mu(y)}{\mu(U)}\frac{\mathrm{d}\mu(x)}{\mu(U)} \leq |U|^s.
\end{aligned}$$

Now, if $\{U_k\}$ is a collection of sets that cover E, then

$$\sum_k |U_k|^s \geq \sum_k \nu(U_k) \geq \nu\left(\bigcup_k U_k \right) \geq \nu(E).$$

This proves that $\mathscr{H}^s(E) \geq \nu(E)$ where $\mathscr{H}^s$ is the s-dimensional Hausdorff measure. Hence $\dim_{\mathrm{H}} E \geq s$.

Notice that if we use instead the potential $\int_{|x-y|<\delta} |x-y|^{-s}\,\mathrm{d}\mu(y)$, we get an estimate on $\mathscr{H}^s_\delta(E)$ which lets us conclude that the Hausdorff measure $\mathscr{H}^s(E)$ is infinite. □

Clearly, since the proof above only rely on Jensen's inequality (convexity), it can easily be generalised to more general settings.

In [21], a lower estimate on the expected value of the Hausdorff dimension of a randomly generated limsup-set $E = \limsup_n U_n$ was given, where U_n are open subsets of the d-dimensional torus $\mathbb{T}^d$, with fixed shape but randomly translated and distributed according to the Lebesgue measure. (This lower estimate has subsequently been turned into an equality in [11].) The proof was based on the following lemma from [22].

Lemma 1.2 (Simplified version of Theorem 1.1 of [22]). *Let E_n be open subsets of $\mathbb{T}^d$, and μ_n Borel probability measures, with $\operatorname{supp}\mu_n \subset E_n$, that converge weakly to the Lebesgue measure on $\mathbb{T}^d$. If there is a constant C such that*

$$\iint |x-y|^{-s}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \leq C \tag{1}$$

holds for all n, then $\limsup_n E_n$ satisfies $\dim_{\mathrm{H}} \limsup_n E_n \geq s$.

The proof of Lemma 1.2 resembles very much the proof of Lemma 1.1. The philosophy behind this lemma is that what is important for the Hausdorff dimension of $\limsup_n E_n$ is the asymptotic distribution of E_n, which is described by the weak limit of μ_n, and the sizes of the sets E_n, which is described by $\iint |x-y|^{-s}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y)$.

Using the method from [21] and Lemma 1.2, M. Rams and myself studied the Hausdorff dimension of sets of the form $E(x, r_n)$ for μ-almost every x in the case T is a map of the interval which preserves a measure μ which is absolutely continuous with respect to Lebesgue measure, and which has summable decay of correlations for function of bounded variation. (There was also some result in the case that μ is not absolutely continuous with respect to Lebesgue measure.)

A weakness of Lemma 1.2 is that it requires the measures μ_n to converge weakly to Lebesgue measure (or at least to something which has a nice density with respect to Lebesgue measure). In this paper we will extend Lemma 1.2 to a more general lemma (Lemma 2.7) in which the measures μ_n may converge to any non-atomic probability measure μ. The proof of this lemma still resembles that of Lemma 1.1, but instead of working with the potential $\int |x-y|^{-s}\,\mathrm{d}\mu(y)$ which has a homogeneous kernel, we will work with "inhomogeneous" potentials with an inhomogeneous kernel adopted to the measure μ.

The conclusion of Lemma 1.2 as stated in [22] is stronger than the version given above. The conclusion is that the set $\limsup_n E_n$ belongs to a certain class of G_δ-set, denoted by $\mathscr{G}^s$. This class was introduced by Falconer [7, 8] and it has the property that every set belonging to $\mathscr{G}^s$ has Hausdorff dimension at least s and the class $\mathscr{G}^s$ is closed under countable intersections. We will generalise Falconer's class $\mathscr{G}^s$ to more general classes $\mathscr{G}^\theta_\mu$, where μ is a non-atomic locally finite measure and $\theta \in (0,1]$. This is done in Section 2.1. From the definition it will be apparent that $\mathscr{G}^\theta_\mu = \mathscr{G}^s$ if μ is Lebesgue measure and $s = \theta d$.

In Section 2.1 we will give theorems that states that the classes $\mathscr{G}^\theta_\mu$ are closed under countable intersections (Theorem 2.4), and that any set in $\mathscr{G}^\theta_\mu$ has a Hausdorff dimension which is at least a certain number which depends only on θ and μ (Theorem 2.5). In all of our applications this leads to $\theta \overline{\dim}_{\mathrm{H}} \mu$ as a lower bound on the dimension, see Remark 4.6.

In Section 2.2 we will introduce the above mentioned inhomogeneous potentials and relate them to the classes $\mathscr{G}^\theta_\mu$. The main result on these potentials is Lemma 2.7, which is a generalisation of Lemma 1.2 and which gives conditions that implies that $\limsup_n E_n$ belongs to the class $\mathscr{G}^\theta_\mu$. Together with Theorem 2.5, it gives us a tool to estimate the Hausdorff dimension of $\limsup_n E_n$ from below.

In Section 2.3, we will see applications of Lemma 2.7 to estimates on Hausdorff dimension of some random limsup-sets (Section 2.3.1), improving some previously known results, and sets of the form $E(x, r_n)$ (Sections 2.3.2–2.3.4). For instance, we prove (Corollary 2.10) that for $\alpha \geq 1$,

$$\dim_{\mathrm{H}}\{\, y : |T_a^n(x) - y| < n^{-\alpha} \text{ infinitely often}\,\} = \frac{1}{\alpha},$$

for almost every $x \in [1-a, 1]$, where $T_a(x) = 1 - ax^2$ is a quadratic map and a belongs to a certain set Δ of positive Lebesgue measure, which has been described and studied by Benedicks and Carleson [2]. Previously, the dimension of such sets have been calculated for the doubling map for almost all x with respect to a Gibbs measure by Fan, Schmeling and Troubetzkoy [13], a result which has been extended to piecewise expanding Markov maps by Liao and Seuret [17]. Some results were also obtained for piecewise expanding maps without a Markov structure by Persson and Rams [23].

2. Definitions and main results

2.1. **The classes $\mathscr{G}^{\boldsymbol{\theta}}_{\boldsymbol{\mu}}$.** Suppose that μ is a non-atomic and locally finite Borel measure on $\mathbb{R}^d$. Take a point $P = (p_1, p_2, \dots, p_d)$ and let $\mathscr{D}_n$ be the collection of cubes of the form

$$D = \left[p_1 + \frac{n_1}{2^n}, p_1 + \frac{n_1+1}{2^n}\right) \times \cdots \times \left[p_d + \frac{n_d}{2^n}, p_d + \frac{n_d+1}{2^n}\right),$$

where $n_1, n_2, \dots, n_d$ are integers, and put $\mathscr{D} = \bigcup_{n\in\mathbb{Z}} \mathscr{D}_n$. We will refer to the elements of $\mathscr{D}$ by the name *dyadic cubes*. We let $D_n(x)$ denote the unique dyadic cube in $\mathscr{D}_n$ containing x and we let $B(x, r)$ denote the open ball with centre x and radius r.

Let

$$R_n = \bigcup_{D\in\mathscr{D}_n} \partial D, \qquad \text{and} \qquad R = \bigcup_{n\in\mathbb{Z}} R_n$$

be the boundaries of the dyadic cubes in $\mathscr{D}_n$ and $\mathscr{D}$ respectively. We will often assume that $\mu(R) = 0$. Since μ is assumed to be locally finite, it is possible to choose the point P defining the dyadic cubes in such a way that $\mu(R) = 0$.

We define for $0 < \theta \leq 1$, the set functions

$$\mathscr{M}^\theta_\mu(E) = \inf\left\{ \sum_k \mu(D_k)^\theta : D_k \in \mathscr{D},\ E \subset \bigcup_k D_k \right\}.$$

Since $0 < \theta \leq 1$, we have by the additivity of the measure μ that $\mathscr{M}^\theta_\mu(D) = \mu(D)^\theta$ for any $D \in \mathscr{D}$. This is an important property that will be used several times in this paper.

The set function $\mathscr{M}^\theta_\mu$ is not a measure (unless $\theta = 1$). If in the definition we consider only covers by D_k which belong to some $\mathscr{D}_n$ for $n > n_0$ and let $n_0 \to \infty$, we obtain in the limit a measure. It is interesting to notice that this construction is a special case of the measure $\mathscr{H}^{q,t}_\mu$ that was introduced by Olsen [20] as a tool in multifractal analysis. This connection with multifractal analysis will not be used in this paper, but it would be interesting to explore it further.

Associated to the set functions $\mathscr{M}^\theta_\mu$ are classes of sets $\mathscr{G}^\theta_\mu$, which we define as follows.

Definition 2.1. Let $0 < \theta \leq 1$. A set $E \subset \mathbb{R}^d$ belongs to $\mathscr{G}^\theta_\mu$ if E is a G_δ-set and if

$$\mathscr{M}^\eta_\mu(E \cap D) = \mathscr{M}^\eta_\mu(D)$$

holds for any $0 < \eta < \theta$ and $D \in \mathscr{D}$.

Remark 2.2. Since $\mathscr{M}_\mu^\eta(D) = \mu(D)^\eta$ holds when $\eta \le 1$ and $D \in \mathscr{D}$, the condition $\mathscr{M}_\mu^\eta(E \cap D) = \mathscr{M}_\mu^\eta(D)$ is equivalent to $\mathscr{M}_\mu^\eta(E \cap D) = \mu(D)^\eta$.

We shall investigate some of the properties of the classes $\mathscr{G}_\mu^\theta$. Below are our main results.

Theorem 2.3. *Let μ be a non-atomic and locally finite Borel measure, with $\mu(R) = 0$. Suppose E_n is a sequence of open sets such that for any $0 < \eta < \theta$, there is a constant $c > 0$ such that*

$$\liminf_{k\to\infty} \mathscr{M}_\mu^\eta(E_n \cap D) \ge c\mathscr{M}_\mu^\eta(D)$$

holds for any $D \in \mathscr{D}$. Then $\limsup_n E_n \in \mathscr{G}_\mu^\theta$.

Theorem 2.4. *Let μ be a non-atomic and locally finite Borel measure, with $\mu(R) = 0$. Suppose that E_n is a sequence of sets in $\mathscr{G}_\mu^\theta$. Then $\bigcap_n E_n \in \mathscr{G}_\mu^\theta$.*

The following theorem makes use of the so called *upper coarse multifractal spectrum* of the measure μ, which we denote by G_μ. See Section 3 for a definition.

Theorem 2.5. *Let μ be a non-atomic and locally finite Borel measure, with $\mu(R) = 0$. Suppose that E belongs to $\mathscr{G}_\mu^\theta$ for some $0 < \theta \le 1$ and that there is a $t > 0$ and an n_0 such that $\mu(D) \le 2^{-tn}$ for all $D \in \mathscr{D}_n$ with $n \ge n_0$. Then $\dim_{\mathrm{H}}(E \cap D) \ge \theta t$ and*

$$\dim_{\mathrm{H}}(E \cap D) \ge \theta \sup\{\, s_0 : \exists\, \varepsilon > 0, \forall\, s \in [t, s_0], (\theta s - G_\mu(s) > \varepsilon)\,\},$$

for any $D \in \mathscr{D}$ with $\mu(D) > 0$.

The proofs of the three theorems above are given in Section 4. These theorems give us the main properties of the classes $\mathscr{G}_\mu^\theta$. In principle, one can use Theorem 2.3 to determine if a limsup-set belongs to the class $\mathscr{G}_\mu^\theta$, but this is not always convenient in practice. In the section below, we therefore define some inhomogeneous potentials and use them to give an alternative method to determine is a limsup-set belongs to $\mathscr{G}_\mu^\theta$.

2.2. **Inhomogeneous potentials and energies.** We define the function $Q\colon \mathbb{R}^d \times \mathbb{R}^d \to \{\emptyset\} \cup \mathscr{D} \cup \{\mathbb{R}^d\}$ by

$$Q(x, y) = D, \qquad x \ne y,$$

where $D \in \mathscr{D} \cup \{\mathbb{R}^d\}$ is chosen such that $x, y \in D$ and D is minimal in sense of inclusion. If $x = y$, we let $Q(x, y) = \emptyset$. Note that it is necessary to include the possibility that $Q(x, y) = \mathbb{R}^d$, since otherwise Q would not always be defined; there are points x and y such that $\{x, y\}$ is a subset of no $D \in \mathscr{D}$.

We use Q to define some inhomogeneous potentials and energies.

Definition 2.6. Let μ and ν be two Borel measures on $\mathbb{R}^d$ and $0 < \theta \le 1$. The (μ, θ)-potential of ν is the function $U_\mu^\theta\nu\colon \mathbb{R}^d \to \mathbb{R} \cup \{\infty\}$ defined by

$$U_\mu^\theta\nu(x) = \int \mu(Q(x, y))^{-\theta}\, \mathrm{d}\nu(y),$$

and the (μ, θ)-energy of ν is the quantity

$$I_\mu^\theta(\nu) = \iint \mu(Q(x, y))^{-\theta}\, \mathrm{d}\nu(y)\mathrm{d}\nu(x) = \int U_\mu^\theta\nu\, \mathrm{d}\nu.$$

The reason for introducing these potentials and energies is the lemma below, which relates the classes $\mathscr{G}_\mu^\theta$ with the energies I_μ^θ. This lemma will be our main tool when we in various applications determine that limsup-sets belong to a class $\mathscr{G}_\mu^\theta$. (We call the result a lemma and not a theorem because of its connection to what is often called Frostman's lemma. Compare also with the related Lemmata 1.1 and 1.2.)

Lemma 2.7. *Let μ be a non-atomic and locally finite Borel measure, with $\mu(R) = 0$. Let E_n be a sequence of open sets. If μ_n are non-atomic Borel measures, with $\operatorname{supp}\mu_n \subset E_n$, that converge weakly to a measure μ, and if for every $\eta < \theta$ there is a constant c_η such that*

$$I_\mu^\eta(\mu_n) = \iint \mu(Q(x, y))^{-\eta}\, \mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \le c_\eta$$

for all n, then $\limsup_n E_n \in \mathscr{G}_\mu^\theta$.

The proof of Lemma 2.7 is in Section 5.

2.3. **Applications.** We give below some applications of the theory presented above.

2.3.1. *Random limsup-sets.* Let μ be a Borel probability measure on $\mathbb{R}^d$. Consider the random sequence of points $(x_n)_{n=1}^\infty$, where the points x_n are independent and distributed according to the measure μ. Let $\alpha > 0$. We are interested in the Hausdorff dimension of the random set

$$E_\alpha = \limsup_n B_n$$

where $B_n = B(x_n, n^{-\alpha})$.

In Section 6 we will prove the following theorem.

Theorem 2.8. *Let*

$$s = s(\mu) = \lim_{\varepsilon \to 0^+} \sup\{\, t : G_\mu(t) \geq t - \varepsilon \,\}.$$

Suppose that $1/\alpha \leq s$. *Almost surely,* $E_\alpha \in \mathscr{G}_\mu^\theta$ *with* $\theta = (\alpha s)^{-1}$ *and*

$$\dim_{\mathrm{H}} E_\alpha \geq \frac{1}{\alpha} \frac{\overline{\dim}_{\mathrm{H}}\, \mu}{s}.$$

Note that $s(\mu)$ is clearly finite since G_μ is a bounded function.

The estimate in Theorem 2.8 should be compared with the result by Ekström and Persson [6, Theorem 1], that almost surely

$$\dim_{\mathrm{H}} E_\alpha \geq \frac{1}{\alpha} - \delta. \qquad \text{where } \delta = \operatorname*{ess\,inf}_{\substack{x \sim \mu \\ \underline{d}_\mu(x) > 1/\alpha}} \overline{d}_\mu(x) - \underline{d}_\mu(x).$$

None of these results implies the other.[1] For instance, if $\delta > 0$ then the estimate in Theorem 2.8 is stronger if α is sufficiently large but may be weaker for small α.

Note that here we treat only the case $1/\alpha \leq s$, whereas in [6], any α was considered. Related results can also be found in a paper by Seuret [25].

2.3.2. *Dynamical Diophantine approximation.* Suppose that X is a metric space, $T\colon X \to X$ is a map and that μ is a T-invariant probability measure.

If $y \in X$ and r_n is a sequence of positive numbers that decreases to 0, then we say that the balls $B(y, r_n)$ are shrinking targets around y. One is often interested in if and how often the orbit of a point x hits one of the targets, i.e. $T^n(x) \in B(y, r_n)$. Such hitting properties depend very much on the points x and y as well as the sequence r_n, but it is often possible to say something about the behaviour for μ almost all x and y.

Here, we assume that $T\colon I \to I$, where I is a compact interval of positive length and that μ is a T-invariant probability measure such that there are positive constants t_1 and c_1 with

$$\mu(D) \leq c_1 2^{-t_1 n} \tag{2}$$

whenever $D \in \mathscr{D}_n$. We assume also that (T, μ) has summable decay of correlations for functions of bounded variation, that is, there is a function p such that

$$\left| \int f \circ T^n \cdot g \, \mathrm{d}\mu - \int f \, \mathrm{d}\mu \int g \, \mathrm{d}\mu \right| \leq p(n) \|f\|_1 \|g\|, \tag{3}$$

where $\|g\|$ is the norm $\|g\| = \|g\|_1 + \operatorname{var} g = \int |g| \, \mathrm{d}\mu + \operatorname{var} g$, and p satisfies

$$\sum_{n=0}^{\infty} p(n) < \infty.$$

Given a point x and a number $\alpha > 0$, we consider the set

$$E_\alpha(x) = \{\, y : |T^n(x) - y| \leq n^{-\alpha} \text{ for infinitely many } n \in \mathbb{N} \,\}.$$

This is the set of points y for which the orbit of x hits the shrinking targets $B(y, n^{-\alpha})$ infinitely many times. The study of the fractal properties of sets of this type (and other related types) is sometimes called dynamical Diophantine approximation, possibly after [13].

The Hausdorff dimension of $E_\alpha(x)$ for μ almost every x was given by Fan, Schmeling and Troubetzkoy in the case that T is the doubling map and μ is a Gibbs measure [13]. Liao and Seuret extended this result to piecewise expanding Markov maps of an interval [17]. Some results for piecewise expanding interval maps without a Markov structure were obtained by Persson and Rams [23].

In Section 7, we shall prove the following result.

[1] In the sense that none follows immediately from the other. Of course, any two true statements imply each other.

Theorem 2.9. *Let*

$$s = s(\mu) = \lim_{\varepsilon \to 0^+} \sup\{\, t : G_\mu(t) \geq t - \varepsilon \,\},$$

and α be such that $1/\alpha \leq s$. Then, almost surely, $E_\alpha(x) \in \mathscr{G}_\mu^\theta$, where $\theta = (\alpha s)^{-1}$.

In particular, there is a set $A \subset I$ of full μ measure such that if $x_1, x_2, \ldots$ is a finite or countable sequence of points in A, then

$$\frac{1}{\alpha} \frac{\overline{\dim}_{\mathrm{H}}\, \mu}{s} \leq \dim_{\mathrm{H}} \bigcap_n E_\alpha(x_n) \leq \frac{1}{\alpha}.$$

Below, we give two concrete applications of Theorem 2.9. The proofs of these results are also in Section 7.

2.3.3. *The quadratic family.* Let us consider the family of quadratic maps $T_a \colon [-1,1] \to [-1,1]$ defined by $T_a(x) = 1 - ax^2$, where a is a parameter in $[0,2]$. Let $\gamma > 0$ be a small number and let

$$\Delta = \{\, a \in [0,2] : |T_a^n(0)| \geq e^{-\gamma n} \text{ and } |(T_a^n)'(T_a(0))| \geq 1.9^n \text{ for all } n \,\}.$$

Benedicks and Carleson [2] have proved that Δ has positive Lebesgue measure. Using Theorem 2.9 we can prove the following result for quadratic maps with $a \in \Delta$.

Corollary 2.10. *Let $T_a \colon [-1,1] \to [-1,1]$ be the quadratic map $T_a(x) = 1 - ax^2$, where $a \in \Delta$. There is a set $A \subset [T_a^2(0), T_a(0)] = [1-a, 1]$ of full Lebesgue measure, such that is $x_1, x_2, \ldots$ is a finite or countable sequence of points of A, then*

$$\dim_{\mathrm{H}} \bigcap_n E_\alpha(x_n) = \frac{1}{\alpha}.$$

2.3.4. *Piecewise expanding maps.* Suppose that $T \colon [0,1] \to [0,1]$ is uniformly expanding and piecewise C^2 with respect to a finite partition, and that T is covering in the sense that for any non-trivial interval $I \subset [0,1]$, there is an n such that $[0,1] \setminus W \subset T^n(I)$, where W denotes the set of points that eventually hit a discontinuity (i.e. an endpoint of a partition element).

We assume that $\phi \colon [0,1] \to \mathbb{R}$ satisfies the following conditions. There is a number n_0 such that

$$\sup e^{S_{n_0}\phi} < \inf L_\phi^{n_0} 1,$$

where $S_{n_0}\phi = \phi + \phi \circ T + \cdots + \phi \circ T^{n_0 - 1}$ and

$$L_\phi f(x) = \sum_{T(y)=x} e^{\phi(y)} f(y).$$

Finally, ϕ is assumed to be piecewise C^2 with respect to the same partition as T, and bounded from below.

Under these conditions, Liverani, Saussol and Vaienti proved that there is a Gibbs measure μ_ϕ with respect to the potential ϕ and correlations decay exponentially fast for functions of bounded variation [18, Theorem 3.1]. To the measure μ_ϕ corresponds a conformal measure ν_ϕ and a density h_ϕ, that is

$$\mu_\phi(E) = \int_E h_\phi \, \mathrm{d}\nu_\phi$$

and if E is a subset of a partition element, then

$$\nu_\phi(T(E)) = \int_E e^{P(\phi) - \phi} \, \mathrm{d}\nu_\phi,$$

where $P(\phi)$ is the topological pressure of ϕ. The density h_ϕ is bounded and bounded away from zero. This implies that $\underline{d}_{\mu_\phi}$ is an invariant function mod μ_ϕ and hence $\underline{d}_{\mu_\phi}$ is constant almost everywhere. We conclude that $\dim_{\mathrm{H}} \mu_\phi = \underline{\dim}_{\mathrm{H}}\, \mu_\phi = \overline{\dim}_{\mathrm{H}}\, \mu_\phi$.

Corollary 2.11. *Let $T \colon [0,1] \to [0,1]$ be a piecewise expanding map satisfying the assumptions above, and assume that*

$$\dim_{\mathrm{H}} \mu_\phi = \lim_{\varepsilon \to 0} \sup\{\, t : G_{\mu_\phi}(t) \geq t - \varepsilon \,\}. \tag{4}$$

Take α such that $1/\alpha \leq \dim_{\mathrm{H}} \mu$.

There is a set $A \subset I$ of full μ measure such that if $x_1, x_2, \ldots$ is a finite or countable sequence of points in A, then

$$\dim_{\mathrm{H}} \bigcap_n E_\alpha(x_n) = \frac{1}{\alpha}.$$

Remark 2.12. It is known that for some Markov maps T, the assumption (4) is satisfied, see Fan–Feng–Wu [10]. It is unknown to me if this is known in full generality in the setting of Corollary 2.11. One might expect that (4) is always satisfied in this case.

Given that (4) is satisfied, Corollary 2.11 improves the result by Rams and myself [23, Theorem 2]. We proved that $\dim_{\mathrm{H}} E_\alpha(x) = \frac{1}{\alpha}$ for

$$\frac{1}{\alpha} < t_1 = \limsup_{m\to\infty} \inf \frac{S_m\phi - mP(\phi)}{-\log|(T^m)'|}.$$

whereas Theorem 4 gives the dimension for a wider range of α as well as for intersections of several sets E_α. The result by Liao and Seuret [17] covers also the case that $\frac{1}{\alpha} > \dim_{\mathrm{H}} \mu$, but is only valid for Markov maps, and only gives the dimension for a single set E_α.

3. Further definitions

In this section we give some further definitions that will be used in this paper.

3.1. **Hausdorff net-measures.** Occasionally, we will make use of the Hausdorff net-measures $\mathscr{N}^s$, defined by

$$\mathscr{N}^s(E) = \lim_{\delta\to 0} \mathscr{N}^s_\delta(E),$$

where

$$\mathscr{N}^s_\delta(E) = \inf\Big\{ \sum_k |D_k|^s : D_k \in \mathscr{D},\ |D_k| < \delta,\ E \subset \bigcup_k D_k \Big\}.$$

In particular, we have $\mathscr{N}^s_\infty = \mathscr{M}^\theta_\lambda$, where λ denotes the d-dimensional Lebesgue measure and $s = \theta d$. Note the conceptual difference between the parameters s and θ: The parameter s should be thought of as a dimension whereas the parameter θ should be thought of something that interpolates between dimension 0 ($\theta = 0$) and full dimension ($\theta = 1$). We will use s and t to denote dimensions and θ and η to denote such interpolating parameters.

Falconer [7, 8] studied the classes $\mathscr{G}^{s/d}_\lambda$, denoting them by $\mathscr{G}^s$. Our study is a generalisation of that of Falconer. Several of the proofs in Section 4 are similar to the corresponding proofs in [8].

3.2. **Dimension spectra and dimension of measures.** The *lower local dimension* of a measure μ at a point x is defined to be

$$\underline{d}_\mu(x) = \liminf_{r\to 0} \frac{\log \mu(B(x,r))}{\log r}.$$

Equivalently,

$$\underline{d}_\mu(x) = \sup\{\, s : \exists\, C \text{ s.t. } \mu(B(x,r)) \le Cr^s,\ \forall\, r \in (0, r_0(x))\,\}.$$

Let

$$D_\mu(s) = \dim_{\mathrm{H}}\{\, x \in \operatorname{supp}\mu :\ \underline{d}_\mu(x) \le s\,\},$$
$$G_\mu(s) = \lim_{\varepsilon\to 0} \limsup_{r\to 0} \frac{\log(N(s+\varepsilon, r) - N(s-\varepsilon, r))}{-\log r},$$

where $N(s,r)$ denotes the number of d-dimensional cubes of the form

$$Q = [k_1 r, (k_1+1)r) \times \cdots \times [k_d r, (k_d+1)r)$$

with $k_1, \dots, k_d \in \mathbb{Z}$ and $\mu(Q) \ge r^s$. The definition of the function D_μ is similar to what is called the *Hausdorff multifractal spectrum* of the lower local dimension of μ, which is obtained if we replace $\underline{d}_\mu(x) \le s$ by $\underline{d}_\mu(x) = s$. The function D_μ is clearly increasing, but this is not the case for the Hausdorff multifractal spectrum. However, D_μ and the Hausdorff multifractal spectrum are often the same for small values of s.

The function G_μ is called the *upper coarse multifractal spectrum* of μ.

We will make use of the *lower Hausdorff dimension* of a measure μ, which is defined as the number

$$\underline{\dim}_{\mathrm{H}}\,\mu = \operatorname*{ess\,inf}_\mu \underline{d}_\mu = \inf\{\, \dim_{\mathrm{H}} A : \mu(A) > 0\,\},$$

see for instance Falconer [9, Proposition 10.2]. Similarly, the *upper Hausdorff dimension* of μ is the number

$$\overline{\dim}_{\mathrm{H}}\,\mu = \operatorname*{ess\,sup}_\mu \underline{d}_\mu = \inf\{\, \dim_{\mathrm{H}} A : \mu(A^{\complement}) = 0\,\},$$

which was used in the statement and will be used in the proof of Theorem 2.8.

If $\underline{\dim}_{\mathrm{H}}\,\mu = \overline{\dim}_{\mathrm{H}}\,\mu$, then the *Hausdorff dimension* of the measure μ is defined as $\dim_{\mathrm{H}}\mu = \underline{\dim}_{\mathrm{H}}\,\mu = \overline{\dim}_{\mathrm{H}}\,\mu$.

4. Proofs of properties of the classes $\mathscr{G}_\mu^\theta$

In this section, we will prove the Theorems 2.3, 2.4 and 2.5.

Throughout this section, we will assume that μ is a non-atomic and locally finite Borel measure. We will mention explicitly when we assume that $\mu(R) = 0$.

In order to prove Theorem 2.3, we will first prove a series of lemmata.

Lemma 4.1. *Suppose $E \subset \mathbb{R}^d$, $0 < \theta \leq 1$ and that there is a constant $c > 0$ such that*

$$\mathscr{M}_\mu^\theta(E \cap D) \geq c\mathscr{M}_\mu^\theta(D)$$

holds whenever $D \in \mathscr{D}$. Then

$$\mathscr{M}_\mu^\eta(E \cap D) = \mathscr{M}_\mu^\eta(D)$$

holds whenever $D \in \mathscr{D}$ and $0 < \eta < \theta$.

Proof. Take $0 < \eta < \theta$ and $D \in \mathscr{D}$, and let $\{D_l\}_{l=1}^\infty$ be a disjoint cover by dyadic cubes of the set $E \cap D$.

Since $\mu(D)$ is finite and μ is not atomic, we may choose a number m such that whenever $C \subset D$ and $C \in \mathscr{D}_n$ for some $n \geq m$, then

$$\mu(C)^{\eta-\theta} \geq c^{-1}\mu(D)^{\eta-\theta}.$$

We write D as a finite disjoint union of dyadic cubes $\mathscr{Q} = \{Q_k\}$ in the following way. All D_k that belong to $\mathscr{D}_n$ for some $n < m$ are put in $\mathscr{Q}$. The part of D that are not covered by these D_k can be written as a union of elements in $\mathscr{D}_m$. These elements of $\mathscr{D}_m$ are added to $\mathscr{Q}$. This construction implies that for any $Q_k \in \mathscr{Q}$, we have that one and only one of the following two cases holds.

(i) there exists an l such that $Q_k = D_l$ and $Q_k \in \mathscr{D}_n$ with $n < m$.
(ii) $Q_k \in \mathscr{D}_m$ and all cubes D_l that intersect $E \cap Q_k$ are contained in Q_k, that is, each of them belong to some $\mathscr{D}_n$ with $n \geq m$.

Suppose Q_k satisfies (i) above. Then

$$\sum_{D_l \subset Q_k} \mu(D_l)^\eta = \mu(Q_k)^\eta = \mu(Q_k)^{\eta-\theta}\mu(Q_k)^\theta \geq \mu(D)^{\eta-\theta}\mu(Q_k)^\theta. \tag{5}$$

Suppose that Q_k satisfies (ii) above. By the choice of m, we have for $D_l \subset Q_k$ that

$$\mu(D_l)^\eta = \mu(D_l)^{\eta-\theta}\mu(D_l)^\theta \geq c^{-1}\mu(D)^{\eta-\theta}\mu(D_l)^\theta.$$

Hence

$$\begin{aligned}\sum_{D_l \subset Q_k} \mu(D_l)^\eta &\geq \sum_{D_l \subset Q_k} c^{-1}\mu(D)^{\eta-\theta}\mu(D_l)^\theta \geq c^{-1}\mu(D)^{\eta-\theta}\mathscr{M}_\mu^\theta(E \cap Q_k) \\ &\geq \mu(D)^{\eta-\theta}\mathscr{M}_\mu^\theta(Q_k) = \mu(D)^{\eta-\theta}\mu(Q_k)^\theta. \end{aligned} \tag{6}$$

Together, (5) and (6) show that

$$\sum_{D_l \subset Q_k} \mu(D_l)^\eta \geq \mu(D)^{\eta-\theta}\mu(Q_k)^\theta$$

for any Q_k.

Summing over all Q_k, we finally get

$$\begin{aligned}\sum_l \mu(D_l)^\eta &= \sum_k \sum_{D_l \subset Q_k} \mu(D_l)^\eta \geq \mu(D)^{\eta-\theta}\sum_k \mu(Q_k)^\theta \\ &\geq \mu(D)^{\eta-\theta}\mathscr{M}_\mu^\theta(D) = \mathscr{M}_\mu^\eta(D).\end{aligned}$$

□

Lemma 4.2. *Suppose $E \subset \mathbb{R}^d$, $0 < \theta \leq 1$ and that there is a constant $c > 0$ such that*

$$\mathscr{M}_\mu^\theta(E \cap D) \geq c\mathscr{M}_\mu^\theta(D)$$

holds whenever $D \in \mathscr{D}$. Then

$$\mathscr{M}_\mu^\theta(E \cap U) \geq c\mathscr{M}_\mu^\theta(U)$$

holds for all open sets U.

Proof. Write U as a disjoint union of dyadic cubes from $\mathscr{D}$,

$$U = \bigcup_k Q_k,$$

and suppose that $(D_l)_{l=1}^\infty$ is a disjoint cover of $E \cap U$ by dyadic cubes.

For each Q_k, with $\mu(Q_k) > 0$ the set $E \cap Q_k$ is not empty, since it would violate the assumption. Hence, for each Q_k, either $\mu(Q_k) = 0$ or we have one of the following two cases:

(1) The dyadic cube Q_k intersect exactly one dyadic cube D_l. We then have $Q_k \subset D_l$.

(2) The dyadic cube Q_k intersect more that one of the dyadic cubes D_l. Then, if D_l intersect Q_k we have that D_l is a strict sub-cube of Q_k.

In case (2), we have

$$\sum_{D_l \subset Q_k} \mu(D_l)^\theta \geq \mathscr{M}_\mu^\theta(E \cap Q_k) \geq c\mathscr{M}_\mu^\theta(Q_k) = c\mu(Q_k)^\theta.$$

We define a cover $\{C_k\}_{k=1}^\infty$ of U using the covers $\{Q_k\}_{k=1}^\infty$ and $\{D_l\}_{l=1}^\infty$. For each k, if Q_k satisfies (1), let

$$\mathscr{Q}_k = \{\, D_l : D_l \cap Q_k \neq \emptyset \,\}.$$

Note that (1) says that $\mathscr{Q}_k$ contains exactly one element. Otherwise, if Q_k satisfies (2), let

$$\mathscr{Q}_k = \{Q_k\}.$$

Hence, in both cases, $\mathscr{Q}_k$ contains exactly one element.

Put $\{C_k\} = \bigcup_{k=1}^\infty \mathscr{Q}_k$. We then have

$$\sum_l \mu(D_l)^\theta \geq \sum_k c\mu(C_k)^\theta \geq c\mathscr{M}_\mu^\theta(U).$$

Since $\{D_l\}_{l=1}^\infty$ is an arbitrary cover of $E \cap U$, this proves that $\mathscr{M}_\mu^\theta(E \cap U) \geq c\mathscr{M}_\mu^\theta(U)$. □

Recall that R_n denotes the boundaries of the dyadic cubes of $\mathscr{D}_n$.

Lemma 4.3. *Suppose U is open and $\mu(R_n) = 0$. Then $\mathscr{M}_\mu^\theta(U \setminus R_n) = \mathscr{M}_\mu^\theta(U)$.*

Proof. Let $D \in \mathscr{D}$ and let D_k be a cover of $D \setminus \partial R_n$ by dyadic cubes. Then

$$\sum \mu(D_k)^\theta \geq \Big(\sum \mu(D_k)\Big)^\theta \geq \mu(D)^\theta, \tag{7}$$

since $\mu(D \setminus R_n) = \mu(D)$, which shows that

$$\mathscr{M}_\mu^\theta(D \setminus R_n) = \mathscr{M}_\mu^\theta(D). \tag{8}$$

Now, let D_k be a disjoint dyadic cover of $U \setminus R_n$. We will modify $\{D_k\}$ into a cover $\mathscr{C} = \{C_k\}$ of U such that

$$\sum \mu(D_k)^\theta \geq \sum \mu(C_k)^\theta.$$

If $D_k \in \mathscr{D}_m$ for some $m \leq n$, then we put D_k in $\mathscr{C}$.

We now proceed by induction over $m > n$. Start with $m = n + 1$ and suppose that $C \in \mathscr{D}_m$. If C is covered by the elements already in $\mathscr{C}$, then we do nothing. Otherwise we proceed as follows.

If $C \setminus R_n$ is covered by the cubes in

$$\mathscr{E}(C) = \left\{ D_k : D_k \in \bigcup_{l \geq m} \mathscr{D}_l,\ D_k \subset C \right\},$$

then

$$\sum_{D_k \in \mathscr{E}(C)} \mu(D_k)^\theta \geq \mu(C)^\theta \tag{9}$$

by (7) (or by (8)), and we let C be an element of $\mathscr{C}$.

This process is carried out for each $C \in \mathscr{D}_m$. The number m is then incremented by one and the process is repeated.

In this way we get the new cover $\mathscr{C}$, which by construction is disjoint. By (9) we have

$$\sum \mu(D_k)^\theta \geq \sum_{C \in \mathscr{C}} \mu(C)^\theta. \tag{10}$$

It is clear that $\mathscr{C}$ covers $U \setminus R_n$, since $\bigcup_k D_k \subset \bigcup_{C \in \mathscr{C}} C$. Let $x \in R_n \cap U$. Then there is a largest dyadic cube D such that $x \in D \subset U$. By the construction of $\mathscr{C}$ we have that either $D \in \mathscr{C}$, or there is a $C \in \mathscr{C}$, with $D \subset C$. In any case, the point x is covered by $\mathscr{C}$ and since $x \in R_n \cap U$ is arbitrary, this proves that $\mathscr{C}$ covers U. Hence

$$\sum_{C \in \mathscr{C}} \mu(C)^\theta \geq \mathscr{M}_\mu^\theta(U). \tag{11}$$

By (10) and (11), we have $\mathscr{M}_\mu^\theta(U \setminus R_n) \geq \mathscr{M}_\mu^\theta(U)$, so that $\mathscr{M}_\mu^\theta(U \setminus R_n) = \mathscr{M}_\mu^\theta(U)$. □

The following increasing sets lemma is a special case of Theorem 52 of Rogers [24]. It is needed to prove Lemma 4.5 and Theorem 2.5.

Lemma 4.4. *If $F_1 \subset F_2 \subset \cdots$ is an increasing sequence of subsets of $\mathbb{R}^d$, then*

$$\mathscr{M}_\mu^\theta\left(\bigcup_{n=1}^\infty F_n\right) = \sup_n \mathscr{M}_\mu^\theta(F_n).$$

Lemma 4.5. *Suppose $\mu(R) = 0$ and that $\{E_n\}_{n=1}^\infty$ is a sequence of open sets such that*

$$\mathscr{M}_\mu^\theta(E_n \cap U) = \mathscr{M}_\mu^\theta(U) \tag{12}$$

holds for any n and any open set U. Then, for any open set U, we have

$$\mathscr{M}_\mu^\theta\left(U \cap \bigcap_{n=1}^\infty E_n\right) = \mathscr{M}_\mu^\theta(U).$$

Proof. If V is a set, we define

$$V_{(-\delta)} = \{\, x \in V : \inf_{y \notin V} |x - y| > \delta \,\}.$$

Let $\varepsilon > 0$. We define a sequence of open sets U_n inductively. Put $U_0 = U$. Suppose that U_{n-1} has been defined and satisfies

$$\mathscr{M}_\mu^\theta(U_{n-1}) > \mathscr{M}_\mu^\theta(U) - \varepsilon.$$

Put $U_n = (E_n \cap (U_{n-1} \setminus (R_n \cup R_{-n})))_{(-\delta_n)}$. Then, by Lemma 4.4, we can choose δ_n so small that $\mathscr{M}_\mu^\theta(U_n)$ is as close to $\mathscr{M}_\mu^\theta(E_n \cap (U_{n-1} \setminus (R_n \cup R_{-n})))$ as we desire. But

$$\begin{aligned}\mathscr{M}_\mu^\theta(E_n \cap (U_{n-1} \setminus (R_n \cup R_{-n}))) &= \mathscr{M}_\mu^\theta(U_{n-1} \setminus (R_n \cup R_{-n})) \\ &= \mathscr{M}_\mu^\theta(U_{n-1}) > \mathscr{M}_\mu^\theta(U) - \varepsilon,\end{aligned}$$

holds by (12), Lemma 4.3 and the assumption on U_{k-1}. Hence we can choose δ_n so small that

$$\mathscr{M}_\mu^\theta(U_n) > \mathscr{M}_\mu^\theta(U) - \varepsilon.$$

Hence by induction, we obtain a sequence $\{U_n\}_{n=0}^\infty$ such that

$$\overline{U}_n \subset U \cap E_n \setminus (R_n \cup R_{-n}), \qquad \text{and} \qquad \mathscr{M}_\mu^\theta(U_n) > \mathscr{M}_\mu^\theta(U) - \varepsilon,$$

holds for every n, from which it follows that

$$\bigcap_{n=0}^\infty \overline{U}_n \subset U \cap \bigcap_{n=1}^\infty E_n \setminus R, \qquad \text{and} \qquad \mathscr{M}_\mu^\theta(U_n) > \mathscr{M}_\mu^\theta(U) - \varepsilon,$$

holds for every n.

Let $\{D_k\}_{k=1}^\infty$ be a cover of $U \cap \bigcap_{n=1}^\infty E_n$ by dyadic cubes. Let $V = \bigcup_k (D_k \setminus \partial D_k)$. The compact set $\bigcap_n \overline{U}_n$ is contained in the open set V. Hence there exists an n such that

$$\overline{U}_n \subset V \subset \bigcup_{k=1}^\infty D_k.$$

Since

$$\sum_{k=1}^\infty \mu(D_k)^\theta \geq \mathscr{M}_\mu^\theta(\overline{U}_n) \geq \mathscr{M}_\mu^\theta(U_n) > \mathscr{M}_\mu^\theta(U) - \varepsilon,$$

this proves that

$$\mathscr{M}_\mu^\theta\left(U \cap \bigcap_{n=1}^\infty E_n\right) \geq \mathscr{M}_\mu^\theta(U) - \varepsilon.$$

As $\varepsilon > 0$ is arbitrary, this implies that

$$\mathscr{M}_\mu^\theta\left(U \cap \bigcap_{n=1}^\infty E_n\right) = \mathscr{M}_\mu^\theta(U). \qquad \square$$

We can now prove Theorem 2.3 and 2.4.

Proof of Theorem 2.3. Clearly, $\limsup_n E_n$ is a G_δ-set, so we only need to show that

$$\mathscr{M}_\mu^\eta(\limsup_n E_n \cap D) \geq \mathscr{M}_\mu^\eta(D)$$

holds for any $\eta < \theta$ and $D \in \mathscr{D}$.

Let $F_n = \bigcup_{k>n} E_k$. Then, by Lemma 4.1,

$$\mathscr{M}_\mu^\eta(F_n \cap D) = \mathscr{M}_\mu^\eta(D)$$

holds for any $\eta < \theta$ and any $D \in \mathscr{D}$. Hence, by Lemma 4.2

$$\mathscr{M}_\mu^\eta(F_n \cap U) \geq \mathscr{M}_\mu^\eta(U)$$

holds for any $\eta < \theta$ and any open set U. Now, Lemma 4.5 finishes the proof. □

Proof of Theorem 2.4. Since each E_n is a G_δ-set, we can write E_n as an intersection

$$E_n = \bigcap_{k=1}^{\infty} U_{n,k},$$

where each $U_{n,k}$ is an open set.

Then $\mathscr{M}_\mu^\eta(D \cap U_{n,k}) = \mathscr{M}_\mu^\eta(D)$ holds for any $U_{n,k}$, $\eta < \theta$ and $D \in \mathscr{D}$. By Lemma 4.2, we have that $\mathscr{M}_\mu^\eta(U \cap U_{n,k}) = \mathscr{M}_\mu^\eta(U)$ holds for any $U_{n,k}$, $\eta < \theta$ and open U.

Now, Lemma 4.5 implies that

$$E = \bigcap_{n=1}^{\infty} \bigcap_{k=1}^{\infty} U_{n,k}$$

satisfies $\mathscr{M}_\mu^\eta(E \cap U) = \mathscr{M}_\mu^\eta(U)$ for any open set and $\eta < \theta$. Since E is a G_δ-set this shows that E belongs to $\mathscr{G}_\mu^\theta$. □

We end this section with the proof of Theorem 2.5, but before we do so, we give a remark on possible improvements.

Remark 4.6. Put

$$\Theta_n(s) = \bigcap_{k=n}^{\infty} \{\, x : \mu(B(x,r)) < r^s \text{ for } r = \sqrt{d}2^{-k} \,\},$$

and

$$\begin{aligned}\Theta(s) &= \bigcup_{n=1}^{\infty} \Theta_n(s) \\ &= \{\, x : \mu(B(x,r)) < r^s \text{ for all } r = \sqrt{d}2^{-n},\ n \in \mathbb{N},\ n > n_0(x) \,\},\end{aligned}$$

Let $s < \underline{\dim}_{\mathrm{H}} \mu$ and $0 < \eta < \theta$. Since $s < \underline{\dim}_{\mathrm{H}} \mu = \operatorname{ess\,inf} \underline{d}_\mu$ we have $\mu(\Theta(s)^{\complement}) = 0$. It follows that if $D \in \mathscr{D}$, then

$$\mathscr{M}_\mu^\eta(D \cap \Theta(s)) = \mathscr{M}_\mu^\eta(D).$$

The set $\Theta(s)$ is a $G_{\delta\sigma}$ set. If it were a G_δ set, we would conclude that $\Theta(s) \in \mathscr{G}_\mu^\theta$, and so $\Theta(s) \cap E \in \mathscr{G}_\mu^\theta$ whenever $E \in \mathscr{G}_\mu^\theta$. This would mean that

$$\mathscr{M}_\mu^\eta(D \cap \Theta(s) \cap E) \geq \mathscr{M}_\mu^\eta(D) > 0$$

if $\mu(D) > 0$. The increasing sets lemma would then give that

$$\mathscr{M}_\mu^\eta(D \cap \Theta_{n_0}(s) \cap E) > 0$$

if n_0 is large enough, and this in turns easily leads to the estimate that $\dim_{\mathrm{H}}(D \cap E) \geq \theta \, \underline{\dim}_{\mathrm{H}} \mu$, which is a stronger estimate than that in Theorem 2.5. Indeed, if $\{D_k\}$ is a dyadic cover of $D \cap \Theta_{n_0}(s) \cap E$ such that each D_k belongs to some $\mathscr{D}_n$ with $n \geq n_0$, then

$$\sum |D_k|^{\eta s} \geq \sum \mu(D_k)^\eta \geq \mathscr{M}_\mu^\eta(D \cap \Theta_{n_0}(s) \cap E) > 0,$$

which shows that $\dim_{\mathrm{H}}(D \cap \Theta_{n_0}(s) \cap E) \geq \eta s$.

Even if $\Theta(s)$ is not a G_δ set, I still suspect that

$$\mathscr{M}_\mu^\eta(D \cap \Theta_n(s) \cap E) > 0$$

holds when $\mu(D) > 0$, but I have not been able to prove this. It would be interesting to know whether this is true or not, since if true, it leads to a stronger dimension estimate than that of Theorem 2.5. More generally, is it true that if Γ is a $G_{\delta\sigma}$ set of full μ measure and $E \in \mathscr{G}_\mu^\theta$, then $\mathscr{M}_\mu^\eta(\Gamma \cap E) > 0$? Is it enough that Γ is a $G_{\delta\sigma}$ set of positive measure?

In any case, it seems not to be a great disadvantage to only have the weaker dimension estimate of Theorem 2.5, since in all applications (Section 2.3) we are actually able to prove not only that the set E under consideration belongs to $\mathscr{G}_\mu^\theta$, but also to $\mathscr{G}_\nu^\theta$, where $\nu = \mu|_{\Theta_n(s)}$ for $s < \overline{\dim}_{\mathrm{H}} \mu$ and n such that $\mu(\Theta_n(s)) > 0$. This is sufficient to conclude that $\dim_{\mathrm{H}} E \geq \theta \, \overline{\dim}_{\mathrm{H}} \mu$.

Proof of Theorem 2.5. The first estimate of $\dim_{\mathrm{H}}(E\cap D)$ is trivial. Suppose $\mu(D)>0$ and let $\{D_k\}$ be a dyadic cover of $E\cap D$. Then

$$\sum |D_k|^{\theta t}\ge c\sum \mu(D_k)^\theta \ge c\mathscr{M}_\mu^\theta(E\cap D)>0,$$

which shows that $\mathscr{N}^{\theta t}(E\cap D)>0$ (the Hausdorff net-measure) and hence $\dim_{\mathrm{H}}(E\cap D)\ge\theta t$.

Take $\eta<\theta$. Let s_0 and ε_0 be such that $\theta s-G_\mu(s)>\varepsilon_0>0$ for all $s\in[t,s_0]$, and D such that $\mu(D)>0$. It is sufficient to prove that $\dim_{\mathrm{H}}(E\cap D)\ge s_0\theta$.

Take $\varepsilon>0$ and $\delta_0>0$. There is then a $\delta=\delta(s)>0$ such that $\delta<\delta_0$ and

$$\limsup_{r\to0}\frac{\log(N(s+\delta,r)-N(s-\delta,r))}{-\log r}\le G_\mu(s)+\frac{1}{2}\varepsilon,$$

and hence also an $r_0=r_0(s)>0$ such that

$$\frac{\log(N(s+\delta,r)-N(s-\delta,r))}{-\log r}\le G_\mu(s)+\varepsilon,\qquad r<r_0.$$

Hence we have

$$N(s+\delta,r)-N(s-\delta,r)\le r^{G_\mu(s)+\varepsilon}. \tag{13}$$

By compactness, there are $t\le s_1<s_2<\cdots<s_n\le s_0$ such that the balls $B(s_k,\delta(s_k))$, $k=1,2,\dots,n$ cover $[t,d]$. Let $\delta=\max\{\delta(s_1),\delta(s_2),\dots,\delta(s_n)\}$ and $r_0=\min\{r_0(s_1),r_0(s_2),\dots,r_0(s_n)\}$. Then $\delta<\delta_0$ since all $\delta(s)<\delta_0$.

By choosing ε and δ_0 sufficiently small, we can achieve that

$$\varepsilon+\theta\delta<\frac{1}{2}\varepsilon_0,$$

where ε_0 has been chosen above to satisfy $\theta s-G_\mu(s)>\varepsilon_0$ for $s\in[t,s_0]$.

The set $E\cap D$ is a bounded set with $\mathscr{M}_\mu^\eta(E\cap D)>0$. Let $\{D_k\}$ be a cover of $E\cap D$ by dyadic cubes belonging to $\bigcup_{l=L}^\infty\mathscr{D}_l$, where we will take L to be large. We group the dyadic cubes covering $E\cap D$ according to their sizes, letting

$$\mathscr{C}_l=\mathscr{D}_l\cap\{D_k\},$$

each $\mathscr{C}_l$ being finite. Each $\mathscr{C}_l$ is then further divided into collections $\mathscr{C}_l^+(k)$ and $\mathscr{C}_l^-$, defined by

$$\mathscr{C}_l^+(k)=\left\{C\in\mathscr{C}_l\setminus\bigcup_{j<k}\mathscr{C}_l^+(j):2^{-(s_k+\delta(s_k))l}\le\mu(C)\le2^{-(s_k-\delta(s_k))l}\right\},$$

for $k=1,2,\dots,n$, and

$$\mathscr{C}_l^-=\mathscr{C}_l\setminus\bigcup_{k=1}^n\mathscr{C}_l^+(k).$$

The condition $C\in\mathscr{C}_l\setminus\bigcup_{j<k}\mathscr{C}_l^+(j)$ in the definition of $\mathscr{C}_l^+(k)$ is there to ensure that every C is in at most one $\mathscr{C}_l^+(k)$. Since the balls $B(s_k,\delta(s_k))$ that cover $[t,d]$ will necessarily overlap, it could otherwise be possible that some C belongs to more than one $\mathscr{C}_l^+(k)$.

By (13), we have

$$\#\mathscr{C}_l^+(k)\le2^{(G_\mu(s_k)+\varepsilon)l}.$$

This implies that

$$\begin{aligned}\sum_{l=L}^\infty\sum_{k=1}^n\sum_{C\in\mathscr{C}_l^+(k)}\mu(C)^\eta&\le\sum_{l=L}^\infty\sum_{k=1}^n2^{(G_\mu(s_k)+\varepsilon)l}2^{-\eta(s_k-\delta)l}\\&=\sum_{l=L}^\infty\sum_{k=1}^n2^{(G_\mu(s_k)-\eta s_k+\varepsilon+\eta\delta)l}\\&\le\sum_{l=L}^\infty n2^{-\frac{1}{2}\varepsilon_0 l}\le\frac{1}{2}\mathscr{M}_\mu^\eta(E\cap D),\end{aligned}$$

if L is large enough.

If $C \in \mathscr{C}_l^-$, then $\mu(C) < 2^{-(s_n+\delta)l} < 2^{-(s_0-\delta)l}$. Hence, with $s = s_0 - \delta$, we have

$$\sum_k |D_k|^{s\eta} \geq \sum_l \sum_{C\in\mathscr{C}_l^-} |C|^{s\eta} \geq \sum_l \sum_{C\in\mathscr{C}_l^-} \mu(C)^\eta$$
$$= \sum_k \mu(D_k)^\eta - \sum_l \sum_k \sum_{C\in\mathscr{C}_l^+(k)} \mu(C)^\eta \geq \frac{1}{2}\mathscr{M}_\mu^\eta(E\cap D).$$

This proves that $\mathscr{N}^{s\eta}(E\cap D) > 0$. Hence $\mathscr{H}^{s\eta}(E\cap D) > 0$ and $\dim_{\mathrm{H}}(E\cap D) \geq s\eta = (s_0-\delta)\eta$. As δ can be made as small as we please and η as close to θ as we like, this finishes the proof. □

5. Proof of Lemma 2.7

The goal of this section is to prove Lemma 2.7.

Throughout this section, we assume that μ is a non-atomic and locally finite Borel measure, with $\mu(R) = 0$. We start by proving the following lemma, which is an important step towards the proof of Lemma 2.7.

Lemma 5.1. *Suppose that E_n are open sets, and μ_n are Borel measures with $\operatorname{supp}\mu_n \subset E_n$. If there is a constant $C > 0$, such that for any $D \in \mathscr{D}$ holds*

$$\liminf_{n\to\infty} \int_D \left(\int_D \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu_n(y)\right)^{-1} \mathrm{d}\mu_n(x) \geq \frac{\mu(D)^\theta}{C}, \tag{14}$$

then $\limsup_n E_n \in \mathscr{G}_\mu^\theta$.

Proof. Let $D \in \mathscr{D}$ and suppose that $\mu(D) > 0$. We consider

$$K_\theta(x) = U_\mu^\theta(\mu_n|_D)(x) = \int_D \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu_n(y).$$

We let ν be a probability measure with support in D and defined by

$$\nu(A) = \frac{\int_A K_\theta^{-1}\,\mathrm{d}\mu_n}{\int_D K_\theta^{-1}\,\mathrm{d}\mu_n}, \qquad A \subset D.$$

We prove that if n is large enough, then, for any $A \subset D$ with $A \in \mathscr{D}$, we have

$$\nu(A) \leq 2C\left(\frac{\mu(A)}{\mu(D)}\right)^\theta. \tag{15}$$

If n is large enough, then the denominator in the expression defining ν satisfies

$$\int_D K_\theta^{-1}\,\mathrm{d}\mu_n \geq \frac{\mu(D)^\theta}{2C}$$

by (14).

By Jensen's inequality, we have (compare with the proof of Lemma 1.1)

$$\begin{aligned}
\int_A K_\theta^{-1}\,\mathrm{d}\mu_n &= \int_A \left(\int_D \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu_n(y)\right)^{-1} \mathrm{d}\mu_n(x)\\
&\leq \int_A \left(\int_A \mu(Q(x,y))^{-\theta}\,\frac{\mathrm{d}\mu_n(y)}{\mu_n(A)}\right)^{-1} \frac{\mathrm{d}\mu_n(x)}{\mu_n(A)}\\
&\leq \int_A \left(\int_A \mu(Q(x,y))^{\theta}\,\frac{\mathrm{d}\mu_n(y)}{\mu_n(A)}\right) \frac{\mathrm{d}\mu_n(x)}{\mu_n(A)}\\
&\leq \mu(A)^\theta.
\end{aligned}$$

This shows that (15) holds when n is large.

Suppose now that $\{D_k\}$ is a cover of $E_n \cap D$ by dyadic cubes and that n is large. Then

$$1 = \nu\left(\bigcup_k D_k\right) \leq \sum_k \nu(D_k) = \sum_k \nu(D_k\cap D) \leq \sum_k 2C\left(\frac{\mu(D_k\cap D)}{\mu(D)}\right)^\theta.$$

Hence, if all D_k are subsets of D, then

$$\sum_k \mu(D_k)^\theta \geq \frac{\mu(D)^\theta}{2C},$$

and so $\mathscr{M}_\mu^\theta(E_n\cap D) \geq (2C)^{-1}\mu(D)^\theta$.

If $D \in \mathscr{D}$ with $\mu(D) = 0$, then $\mathscr{M}_\mu^\theta(E_n\cap D) \geq (2C)^{-1}\mu(D)^\theta$ is trivially satisfied.

We have thus showed that

$$\liminf_{n\to\infty} \mathscr{M}_\mu^\theta(E_n \cap D) \geq (2C)^{-1}\mu(D)^\theta, \qquad D \in \mathscr{D}.$$

Now, Theorem 2.3 finishes the proof. □

Lemma 5.2. *If μ is a finite and non-atomic Borel measure and $D \in \mathscr{D}$, then*

$$\mu(D)^{1-\theta} \leq \int_D \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu(y) \leq \frac{\mu(D)^{1-\theta}}{1-\theta},$$

for $x \in D$, and in particular

$$(1-\theta)\mu(D)^\theta \leq \int_D \left(\int_D \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu(y)\right)^{-1} \mathrm{d}\mu(x) \leq \mu(D)^\theta,$$

and

$$\mu(D)^{2-\theta} \leq \int_D\int_D \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y) \leq \frac{\mu(D)^{2-\theta}}{1-\theta}.$$

Proof. We may assume that μ is a probability measure on D. By Fubini's theorem we can write

$$\int_D \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu(y) = 1 + \int_1^\infty \mu(D_{n(u)})\,\mathrm{d}u,$$

where $D_{n(u)} \in \mathscr{D}_{n(u)}$ is chosen such that $x \in D_{n(u)}$ and $\mu(D_{n(u)})^{-\theta} \geq u$, and $n(u)$ is chosen as small as possible. Hence we have

$$\begin{aligned} 1 \leq \int_D \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu(y) &= 1 + \int_1^\infty \mu(D_{n(u)})\,\mathrm{d}u \\ &\leq 1 + \int_1^\infty u^{-1/\theta}\,\mathrm{d}u = \frac{1}{1-\theta}. \end{aligned}$$

□

The following two results are variants of Lemma 2.2 and Corollary 2.3 of [22], and proved in the same way.

Lemma 5.3. *Let μ be a Borel measure. Assume that for some $\theta \in (0,1)$,*

$$I_\mu^\theta(\mu) = \iint \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y) < \infty.$$

Then, if $M_m = \{\,(x,y) : \mu(Q(x,y))^{-\theta} > m\,\}$, then we have, for $0 < \eta < \theta$, that

$$\iint_{M_m} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y) \leq I_\mu^\theta(\mu)\frac{\theta}{\theta-\eta}m^{\eta/\theta-1}.$$

Proof. The assumption implies that $\mu\times\mu(M_m) \leq I_\mu^\theta(\mu)/m$. It then follows that

$$\begin{aligned} \iint_{M_m} \mu(Q)^{-\eta}\,\mathrm{d}\mu\mathrm{d}\mu &= m^{\eta/\theta}\mu\times\mu(M_m) + \int_{m^{\eta/\theta}}^\infty \mu\times\mu(M_{u^{\theta/\eta}})\,\mathrm{d}u \\ &\leq I_\mu^\theta(\mu)m^{\eta/\theta-1} + I_\mu^\theta(\mu)\int_{m^{\eta/\theta}}^\infty u^{-\theta/\eta}\,\mathrm{d}u \\ &= I_\mu^\theta(\mu)\frac{\theta}{\theta-\eta}m^{\eta/\theta-1}. \end{aligned}$$

□

Corollary 5.4. *If μ_n are non-atomic Borel measures that converge weakly to a measure μ, and if $\iint \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y)$ are uniformly bounded for some $\theta \in (0,1)$, then, for $0 < \eta < \theta$ and $D \in \mathscr{D}$ with $\mu(D) > 0$,*

$$\iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \to \iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y),$$

as $n \to \infty$.

Proof. Let $\varepsilon > 0$, $0 < \eta < \theta$ and $D \in \mathscr{D}$ with $\mu(D) > 0$. We put

$$M_m = \{\,(x,y) \in D\times D : \mu(Q(x,y))^{-\theta} > m\,\}.$$

Since $\iint \mu(Q(x,y))^{-\theta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y)$ are uniformly bounded, the estimates of

$$\iint_{M_m} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y)$$

provided by Lemma 5.3 are uniform in n, and we can take m and N so that

$$\iint_{M_m} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) < \varepsilon$$

holds for all $n > N$. We then have

$$\iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \\ \leq \varepsilon + \iint_{D\times D} \min\{\mu(Q(x,y))^{-\eta}, m^{\eta/\theta}\}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y).$$

Since the measures μ_n converge weakly to μ and $\mu(\partial C) = 0$ for all $C \in \mathscr{D}$ we have that $\mu_n(C) \to \mu(C)$ for all $C \in \mathscr{D}$. Hence

$$\begin{aligned}\iint_{D\times D} \min\{\mu(Q(x,y))^{-\eta}, m^{\eta/\theta}\}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \\ &\to \iint_{D\times D} \min\{\mu(Q(x,y))^{-\eta}, m^{\eta/\theta}\}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y) \\ &\leq \iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y),\end{aligned}$$

where the convergence holds because $\min\{\cdots\}$ is a bounded function. As ε is arbitrary, this shows that

$$\limsup_{n\to\infty} \iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \leq \iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y).$$

The inequality

$$\liminf_{n\to\infty} \iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \geq \iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y)$$

is trivial since

$$\begin{aligned}\iint_{D\times D} &\mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \\ &\geq \iint_{D\times D} \min\{\mu(Q(x,y))^{-\eta}, m\}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \\ &\to \iint_{D\times D} \min\{\mu(Q(x,y))^{-\eta}, m\}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y), && n\to\infty \\ &\to \iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y), && m\to\infty,\end{aligned}$$

and finishes the proof. □

We can now give the proof of Lemma 2.7.

Proof of Lemma 2.7. By Corollary 5.4 we have for any $D \in \mathscr{D}$ that

$$\iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n(x)\mathrm{d}\mu_n(y) \to \iint_{D\times D} \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu(x)\mathrm{d}\mu(y) \leq c_\eta,$$

as $n \to \infty$.

Since

$$\int_D \left(\int_D \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n\right)^{-1} \mathrm{d}\mu_n(y) \geq \left(\int_D\int_D \mu(Q(x,y))^{-\eta}\,\frac{\mathrm{d}\mu_n(x)}{\mu_n(D)}\frac{\mathrm{d}\mu_n(y)}{\mu_n(D)}\right)^{-1},$$

we therefore have by Lemma 5.2 that

$$\liminf_{n\to\infty} \int_D \left(\int_D \mu(Q(x,y))^{-\eta}\,\mathrm{d}\mu_n\right)^{-1} \mathrm{d}\mu_n(y) \geq (1-\eta)\mu(D)^\eta.$$

Lemma 5.1 then implies that $\limsup_n E_n \in \mathscr{G}_\mu^\eta$. As this holds for all $\eta < \theta$, we conclude that $\limsup_n E_n \in \mathscr{G}_\mu^\theta$. □

6. Proof of Theorem 2.8

In this section, we will prove Theorem 2.8. We assume that μ is a Borel probability measure on $\mathbb{R}^d$ and consider the random sequence of points $(x_n)_{n=1}^{\infty}$, where the points x_n are independent and distributed according to the measure μ. Recall that we are considering the random set

$$E_\alpha = \limsup_n B_n$$

where $B_n = B(x_n, n^{-\alpha})$ and that

$$s = s(\mu) = \lim_{\varepsilon \to 0} \sup\{\, t : G_\mu(t) \geq t - \varepsilon \,\}.$$

6.1. **Definition of the measure ν.** We will use the theory developed so far in this paper. Since μ is a probability measure we can choose the point P which defines the dyadic cubes in such a way that $\mu(R) = 0$.

We will prove that

$$E_\alpha \in \mathscr{G}_\mu^\theta, \qquad \theta = \frac{1}{s\alpha}$$

holds almost surely. To do so, we shall pick a set C, and first prove that $E_\alpha \in \mathscr{G}_\nu^\theta$ with $\nu = \frac{\mu|_C}{\mu(C)}$. Using a limit argument, we will then conclude that $E_\alpha \in \mathscr{G}_\mu^\theta$ holds almost surely. The dimension estimate will follow from Theorem 2.5 and the fact that $E_\alpha \in \mathscr{G}_\nu^\theta$, and is a stronger estimate than we would have got from Theorem 2.5 using only that $E_\alpha \in \mathscr{G}_\mu^\theta$.

We will assume that μ has compact support. This is no restriction, since when proving that $E_\alpha \in \mathscr{G}_\mu^\theta$ and $E_\alpha \in \mathscr{G}_\nu^\theta$, we may consider instead the measure restricted to dyadic cubes.

Let $\varepsilon > 0$. Take $t_0 = \sup\{\, t : G_\mu(t) \geq t - 2\varepsilon \,\}$ and let $G = \sup\{\, G_\mu(t) : t \geq t_0 \,\}$. Both t_0 and G are finite, since G_μ is a bounded function.

For each n, let A_n be the union of those cubes of $\mathscr{D}_n$ which have μ-measure at most 2^{-nt_0}, that is

$$A_n = \bigcup_{\substack{D \in \mathscr{D}_n \\ \mu(D) \leq 2^{-nt_0}}} D.$$

Lemma 6.1. *There is a number n_0 such that $\mu(A_n) \leq 2^{-\varepsilon n}$ for $n > n_0$.*

Proof. Let $t \geq t_0$ and $\delta_0 > 0$. There is a $\delta(t) \in (0, \delta_0)$ such that

$$\limsup_{r \to 0} \frac{\log(N(t + \delta(t), r) - N(t - \delta(t), r))}{-\log r} \leq G_\mu(t) + \frac{1}{4}\varepsilon,$$

and hence there is also an $r_0(t)$ such that

$$\frac{\log(N(t + \delta(t), r) - N(t - \delta(t), r))}{-\log r} \leq G_\mu(t) + \frac{1}{2}\varepsilon, \qquad r < r_0(t).$$

Hence we have

$$N(t + \delta(t), r) - N(t - \delta(t), r) \leq r^{-(G_\mu(t) + \frac{1}{2}\varepsilon)}, \qquad r < r_0(t).$$

By compactness, there are $t_0 \leq t_1 \leq \cdots \leq t_p \leq d + 2\varepsilon$ such that the intervals $B(t_k, \delta(t_k))$, $1 \leq k \leq p$ cover $[t_0, d + 2\varepsilon]$. Let $\delta = \max\{\delta(t_1), \ldots \delta(t_p)\} < \delta_0$ and $r_0 = \min\{r_0(t_1), \ldots, r_0(t_p)\}$.

Those cubes of $\mathscr{D}_n$ that have positive measure which is at most $2^{-(d+2\varepsilon)n}$ are at most $c2^{dn}$, where c is a constant, since there are at most $c2^{dn}$ cubes of $\mathscr{D}_n$ with positive measure (since the support is compact).

Take N such that $2^{-N} < r_0$. We then have for $n > N$ that

$$\mu(A_n) \leq c2^{dn}2^{-(d+2\varepsilon)n} + \sum_{k=1}^{p} 2^{(G_\mu(t_k) + \frac{1}{2}\varepsilon)n}2^{-(t_k - \delta)n} \leq c2^{-2\varepsilon n} + p2^{(\delta - \frac{3}{2}\varepsilon)n},$$

since $G_\mu(t_k) \leq t_k - 2\varepsilon$. Since δ_0 is arbitrary and $\delta < \delta_0$, we can make δ as small as we like, and hence we can achieve that

$$\mu(A_n) \leq (p+1)2^{-\frac{3}{2}\varepsilon n}$$

for $n > N$. Hence $\mu(A_n) \leq 2^{-\varepsilon n}$ if n is large enough. □

Put

$$\Theta_n(t) = \bigcap_{k=n}^{\infty} \{\, x : \mu(B(x, r)) < r^t \text{ for } r = 2^{-k} \,\},$$

$$\Theta(t) = \bigcap_{n=1}^{\infty} \Theta_n(t).$$

If $t_1 < \overline{\dim}_{\mathrm{H}} \mu$ then $\mu(\Theta(t_1)) > 0$ and so $\mu(\Theta_{n_1}(t_1)) > 0$ if n_1 is large enough. We assume from now on that $t_1 < \overline{\dim}_{\mathrm{H}} \mu$, and we choose a number m_1 so large that $\mu(\Theta_{m_1}(t_1)) > 0$.

Let

$$\mathscr{C}_m = \left\{ D \in \mathscr{D}_{[\beta m]} : \mu(\Theta_{m_1}(t_1) \cap D) \geq \frac{5}{6}\mu(D) \right\}.$$

(We use $\mathscr{D}_{[\beta m]}$ instead of $\mathscr{D}_m$, since we are later going to use that the balls B_m which define the limsup-set E_α have the property that $D \subset B_m$ for some $D \in \mathscr{D}_{[\beta m]}$.)

By Lebesgue density theorem,

$$\mu(\Theta_{m_1}(t_1) \setminus \cup \mathscr{C}_m) \to 0, \qquad m \to \infty,$$

where we have used the notation

$$\cup \mathscr{A} = \bigcup_{A \in \mathscr{A}} A.$$

This notation will appear several times below.

There exists a sequence (m_k) such that the set

$$\hat{\Theta} = \Theta_{m_1}(t_1) \cap \bigcap_{k=1}^{\infty} \cup \mathscr{C}_{m_k}$$

satisfies $\mu(\hat{\Theta}) > \frac{1}{2}\mu(\Theta_{m_1}(t_1)) > 0$ and

$$\sum_{l=k+1} \mu(\Theta_{m_1}(t_1) \setminus \cup \mathscr{C}_{m_k}) < \frac{1}{6} 2^{-[\beta m_k] t_0}.$$

By construction, the set $\hat{\Theta}$ has the following property. If $D \in \mathscr{D}_{[\beta m_k]}$ for some m_k, then either $D \subset \hat{\Theta}^{\complement}$ or

$$\begin{aligned} \mu(\hat{\Theta} \cap D) &\geq \mu(\Theta_{m_1}(t_1) \cap D) - \sum_{l=k+1} \mu(\Theta_{m_1}(s) \setminus \cup \mathscr{C}_{m_k}) \\ &\geq \frac{5}{6}\mu(D) - \frac{1}{6} 2^{-[\beta m_k] t_0}. \end{aligned} \tag{16}$$

Let $\beta > \alpha$. We choose a sparse sequence (n_k), which is a subsequence of (m_k), as follows. By Lemma 6.1, there is a number n_0 such that $\mu(A_n) \leq 2^{-\varepsilon n}$ for $n \geq n_0$. For $k \geq 1$, we may choose inductively a sequence n_k such that $\beta n_1 > n_0$ and so that

$$\frac{1}{6} 2^{-[\beta n_k] t_0} > \sum_{l=k+1}^{\infty} 2^{-\varepsilon [\beta n_l]} \tag{17}$$

holds for all $k \geq 1$. The sequence n_k can in fact be chosen according to the following lemma.

Lemma 6.2. *The sequence n_k can be chosen so that the following holds. If $D \in \mathscr{D}_{[\beta n_k]}$ for some k, then have*

$$\nu(D) = 0 \text{ or } \nu(D) > 2^{-t_0 [\beta n_k] - 1}.$$

Proof. Let $B_n = A_n^{\complement}$, and

$$C = \bigcap_{k=1}^{\infty} B_{[\beta n_k]} \cap \hat{\Theta}.$$

We have

$$\begin{aligned} \mu(C) &\geq \mu(\hat{\Theta}) - \sum_{k=1}^{\infty} \mu(A_{[\beta n_k]}) \geq \mu(\hat{\Theta}) - \sum_{k=1}^{\infty} 2^{-\varepsilon [\beta n_k]} \\ &\geq \mu(\hat{\Theta}) - \sum_{k=[\beta n_1]}^{\infty} 2^{-\varepsilon k} = \mu(\hat{\Theta}) - \frac{2^{-\varepsilon [\beta n_1]}}{1 - 2^{-\varepsilon}}. \end{aligned}$$

Hence, if $t_1 < \overline{\dim}_{\mathrm{H}} \mu$, then we can ensure that $\mu(C) > \frac{1}{2}\mu(\hat{\Theta}) > 0$ by choosing n_1 sufficiently large. We assume that $\mu(C) > \frac{1}{2}\mu(\hat{\Theta})$ holds.

We now put $\nu = \frac{\mu|_C}{\mu(C)}$. Clearly, ν is absolutely continuous with respect to μ, with bounded density. Moreover, by the choice of the sequence n_k, if $D \in \mathscr{D}_{[\beta n_k]}$ for some k, then

$$\nu(D) \leq \frac{\mu(D)}{\mu(C)} \leq \frac{2}{\mu(\hat{\Theta})} \mu(D),$$

and if $\mu(D) > 2^{-t_0[\beta n_k]}$ then, either $D \subset \hat{\Theta}^{\complement} \cup A_{[\beta n_l]}$ for some $l < k$ and $\nu(D) = 0$, or

$$\begin{aligned}\nu(D) &\geq \mu(C \cap D) \geq \mu(\hat{\Theta} \cap D) - \mu\Big(\bigcup_{l=k+1}^{\infty} A_{[\beta n_l]}\Big) \\ &\geq \frac{5}{6}\mu(D) - \frac{1}{6}2^{-[\beta n_k]t_0} - \sum_{l=k+1}^{\infty} 2^{-\varepsilon[\beta n_l]} \\ &\geq \frac{5}{6}\mu(D) - \frac{1}{6}2^{-[\beta n_k]t_0} - \frac{1}{6}2^{-[\beta n_k]t_0} \geq \frac{1}{2}\mu(D),\end{aligned}$$

holds by (16) and (17).

Finally, if $D \in \mathscr{D}_{[\beta n_k]}$ is such that $\mu(D) \leq 2^{-t_0[\beta n_k]}$, then $D \subset A_{[\beta n_k]}$ and so $\nu(D) = 0$. □

6.2. **Almost surely $E_\alpha \in \mathscr{G}_\nu^\theta$.** In this section, we are going to prove that $E_\alpha \in \mathscr{G}_\nu^\theta$ holds almost surely, using to use Lemma 2.7. We will therefore check that the conditions of Lemma 2.7 holds almost surely.

Take $\beta > \alpha$ such that $\frac{1}{\beta} \leq t_0$. We define the random probability measures

$$\nu_n = 2^{-(n-1)} \sum_{k=2^{n-1}+1}^{2^n} \frac{\nu|_{D_k}}{\nu(D_k)},$$

where $D_k \in \mathscr{D}_{[\beta n]}$ are chosen so that $x_k \in D_k$. If $\nu(D_k) = 0$, then $\frac{\nu|_{D_k}}{\nu(D_k)}$ should be interpreted as 0. It is then clear that almost surely, ν_n converge weakly to ν, and that

$$\operatorname{supp} \nu_n \subset \bigcup_{k=2^{n-1}+1}^{2^n} B(x_k, k^{-\alpha})$$

if n is large.

We want to estimate the expectation of $\iint \nu(Q(x,y))^{-\theta}\, \mathrm{d}\nu_n \mathrm{d}\nu_n$. Let E denote expectation and split the expectation into two parts,

$$\mathsf{E} \iint \nu(Q(x,y))^{-\theta}\, \mathrm{d}\nu_n(x)\mathrm{d}\nu_n(y) = E_1 + E_2,$$

where

$$\begin{aligned}E_1 &= 2^{-2n-2} \sum_{k \neq l} \mathsf{E} \int_{D_k} \int_{D_l} \nu(Q(x,y))^{-\theta}\, \frac{\mathrm{d}\nu(x)}{\nu(D_l)} \frac{\mathrm{d}\nu(y)}{\nu(D_k)}, \\ E_2 &= 2^{-2n-2} \sum_{k=2^{n-1}+1}^{2^n} \mathsf{E} \int_{D_k} \int_{D_k} \nu(Q(x,y))^{-\theta}\, \frac{\mathrm{d}\nu(x)}{\nu(D_k)} \frac{\mathrm{d}\nu(y)}{\nu(D_k)}.\end{aligned}$$

We first consider E_1.

Lemma 6.3. *For $k \neq l$ we have*

$$\mathsf{E} \int_{D_k} \int_{D_l} \nu(Q(x,y))^{-\theta}\, \frac{\mathrm{d}\nu(x)}{\nu(D_l)} \frac{\mathrm{d}\nu(y)}{\nu(D_k)} = \iint \nu(Q(x,y))^{-\theta}\, \mathrm{d}\nu(x)\mathrm{d}\nu(y),$$

and

$$E_1 = (1 - 2^{-(n-1)}) \iint \nu(Q(x,y))^{-\theta}\, \mathrm{d}\nu(x)\mathrm{d}\nu(y).$$

Proof. We have

$$\begin{aligned}&\mathsf{E} \int_{D_k} \int_{D_l} \nu(Q(x,y))^{-\theta}\, \frac{\mathrm{d}\nu(x)}{\nu(D_l)} \frac{\mathrm{d}\nu(y)}{\nu(D_k)} \\ &\qquad = \iint \int_{D_k} \int_{D_l} \nu(Q(x,y))^{-\theta}\, \frac{\mathrm{d}\nu(x)}{\nu(D_l)} \frac{\mathrm{d}\nu(y)}{\nu(D_k)}\, \mathrm{d}\nu(x_k)\mathrm{d}\nu(x_l).\end{aligned}$$

If x_k and x_l are such that $D_k \neq D_l$, then $Q(x,y)$ is constant for $(x,y) \in D_k \times D_l$ and in fact $Q(x,y) = Q(x_k, x_l)$. This is not the case if $D_k = D_l$. But if $D \in \mathscr{D}_{[\beta n]}$, then

$$\int_{D_k} \int_{D_l} \nu(Q(x,y))^{-\theta}\, \frac{\mathrm{d}\nu(x)}{\nu(D_l)} \frac{\mathrm{d}\nu(y)}{\nu(D_k)} = \int_D \int_D \nu(Q(x,y))^{-\theta}\, \frac{\mathrm{d}\nu(x)}{\nu(D)} \frac{\mathrm{d}\nu(y)}{\nu(D)}$$

whenever x_k and x_l are such that $D_k = D_l = D$, that is for $x_k, x_l \in D$.

Hence we have

$$\begin{aligned}\iint\int_{D_k}\int_{D_l}\nu(Q(x,y))^{-\theta}\,\frac{\mathrm{d}\nu_n(x)}{\nu(D_l)}\frac{\mathrm{d}\nu_n(y)}{\nu(D_k)}\,\mathrm{d}\nu(x_k)\mathrm{d}\nu(x_l)&\\ =\iint_{D_k\neq D_l}\nu(Q(x_k,x_l))^{-\theta}\,\mathrm{d}\nu(x_k)\mathrm{d}\nu(x_l)&\\ +\sum_{D\in\mathscr{D}_{[\beta n]}}\iint_{D\times D}\iint_{D\times D}\nu(Q(x,y))^{-\theta}\,\frac{\mathrm{d}\nu(x)}{\nu(D)}\frac{\mathrm{d}\nu(y)}{\nu(D)}\mathrm{d}\nu(x_k)\mathrm{d}\nu(x_l)&\\ =\iint_{D_k\neq D_l}\nu(Q(x_k,x_l))^{-\theta}\,\mathrm{d}\nu(x_k)\mathrm{d}\nu(x_l)&\\ +\sum_{D\in\mathscr{D}_{[\beta n]}}\iint_{D\times D}\nu(Q(x_k,x_l))^{-\theta}\,\mathrm{d}\nu(x_k)\mathrm{d}\nu(x_l)&\\ =\iint\nu(Q(x,y))^{-\theta}\,\mathrm{d}\nu(x)\mathrm{d}\nu(y),&\end{aligned}$$

which is the first equality of the lemma. The second equality of the lemma follows immediately from the first equality, since the first equality says that all terms in the sum defining E_1 are equal to $\iint\nu(Q(x,y))^{-\theta}\,\mathrm{d}\nu(x)\mathrm{d}\nu(y)$. □

We now consider E_2. Recall that $D_n(x)$ denotes the unique element of $\mathscr{D}_n$ that contains x.

Lemma 6.4. *We have*

$$\mathsf{E}\int_{D_k}\int_{D_k}\nu(Q(x,y))^{-\theta}\,\frac{\mathrm{d}\nu(x)}{\nu(D_k)}\frac{\mathrm{d}\nu(y)}{\nu(D_k)}\le\frac{1}{1-\theta}\int\nu(D_{[\beta n]}(x))^{-\theta}\,\mathrm{d}\nu(x)$$

and

$$E_2\le\frac{2^{-(n-1)}}{1-\theta}\int\nu(D_{[\beta n]}(x))^{-\theta}\,\mathrm{d}\nu(x).$$

Proof. By Lemma 5.2

$$\int_{D_k}\int_{D_k}\nu(Q(x,y))^{-\theta}\,\frac{\mathrm{d}\nu(x)}{\nu(D_k)}\frac{\mathrm{d}\nu(y)}{\nu(D_k)}\le\frac{\nu(D_k)^{-\theta}}{1-\theta}.$$

Hence

$$\mathsf{E}\int_{D_k}\int_{D_k}\nu(Q(x,y))^{-\theta}\,\frac{\mathrm{d}\nu(x)}{\nu(D_k)}\frac{\mathrm{d}\nu(y)}{\nu(D_k)}\le\frac{1}{1-\theta}\int\nu(D_{[\beta n]}(x))^{-\theta}\,\mathrm{d}\nu(x),$$

which is the first estimate in the lemma. The second estimate of the lemma follows from the first estimate and the definition of E_2. □

From Lemma 6.3 and 6.4 we get the following corollary.

Corollary 6.5.

$$\begin{aligned}\mathsf{E}\iint\nu(Q(x,y))^{-\theta}\,\mathrm{d}\nu_n(x)\mathrm{d}\nu_n(y)&\\ \le\iint\nu(Q(x,y))^{-\theta}\,\mathrm{d}\nu(x)\mathrm{d}\nu(y)+\frac{2^{-(n-1)}}{1-\theta}\int\nu(D_{[\beta n]}(x))^{-\theta}\,\mathrm{d}\nu(x).&\end{aligned}$$

We need to estimate $\int\nu(D_{[\beta n]}(x))^{-\theta}\,\mathrm{d}\nu(x)$. By Lemma 6.2, we have that either $\nu(D_{[\beta n_k]}(x))=0$ or $\nu(D_{[\beta n_k]}(x))>2^{-t_0[\beta n_k]-1}$. We therefore have

$$\int\nu(D_{[\beta n_k]}(x))^{-\theta}\,\mathrm{d}\nu(x)<2^{(t_0\beta n_k+1)\theta}\le2^{t_0\beta n_k\theta+1}.$$

Hence, by Corollary 6.5, if $t_0\beta\theta\le1$, then the expectations

$$\mathsf{E}\iint\nu(Q(x,y))^{-\theta}\,\mathrm{d}\nu_{n_k}(x)\mathrm{d}\nu_{n_k}(y)$$

are uniformly bounded. Take $\theta=\frac{1}{t_0\beta}$. Then $\theta\le1$ since $\frac{1}{\beta}\le t_0$.

Almost surely, there is then a sequence m_k, which is a subsequence of n_k, and such that

$$\iint\nu(Q(x,y))^{-\theta}\,\mathrm{d}\nu_{m_k}(x)\mathrm{d}\nu_{m_k}(y)$$

is uniformly bounded and we assume that m_k is such a sequence.

Lemma 2.7 then implies that $\limsup_k E_{m_k} \in \mathscr{G}_\nu^\theta$ with $\theta = (t_0\beta)^{-1}$. Hence we have almost surely that $E_\alpha \in \mathscr{G}_\nu^\theta$ with $\theta = (t_0\beta)^{-1}$. Since t_0 can be chosen arbitrary close to $s(\mu)$ and β can be taken arbitrary close to α, this proves that almost surely $E_\alpha \in \mathscr{G}_\nu^\theta$ with $\theta = (s\alpha)^{-1}$. By taking t_1 close to $\overline{\dim}_{\mathrm{H}}\,\mu$, the first dimension estimate of Theorem 2.5 implies that

$$\dim_{\mathrm{H}} E_\alpha \geq \frac{1}{\alpha}\frac{\overline{\dim}_{\mathrm{H}}\,\mu}{s}$$

almost surely.

We have shown that almost surely $E_\alpha \in \mathscr{G}_\nu^\theta$ with $\theta = (t_0\beta)^{-1}$. This means, by Definition 2.1, that for $\eta < \theta$, we have

$$\mathscr{M}_\nu^\eta(E \cap D) = \mathscr{M}_\nu^\eta(D)$$

whenever D is a dyadic cube. Since $\nu = \frac{\mu|_C}{\mu(C)}$, we have

$$\mu(C)^{-\eta}\mathscr{M}_\mu^\eta(E \cap D) \geq \mathscr{M}_\nu^\eta(E \cap D) = \mathscr{M}_\nu^\eta(D) = \nu(D)^\eta = \mu(C \cap D)^\eta.$$

We can make $\mu(C)$ as close to 1 as we like (if we let $t_1 < \underline{\dim}_{\mathrm{H}}\,\mu$), and therefore we almost surely have

$$\mathscr{M}_\mu^\eta(E \cap D) \geq \mu(D)^\eta = \mathscr{M}_\mu^\eta(D).$$

Hence $E_\alpha \in \mathscr{G}_\mu^\theta$ almost surely, with $\theta = (s\alpha)^{-1}$.

7. Proofs of result on dynamical Diophantine approximation

In this section we will prove Theorem 2.9 and Corollaries 2.10 and 2.11.

We start by proving the corollaries, which are consequences of Theorem 2.9.

Proof of Corollary 2.10. From Young [26] we know that when $a \in \Delta$, there is a T_a-invariant measure μ_a with properties described in the following.

The support of μ_a is $[T_a^2(0), T_a(0)]$, μ_a is absolutely continuous with respect to Lebesgue measure and the density of μ_a is bounded away from 0 on the support [26, Theorem 2]. Hence, we have $s = 1$, where s is defined as in Theorem 2.9.

The density ρ of μ_a can be written as $\rho = \rho_1 + \rho_2$, where

$$0 \leq \rho_2(x) \leq C\sum_{k=1}^{\infty}\frac{1.9^{-k}}{\sqrt{|x - T_a^k(0)|}}$$

and ρ_1 is bounded [26, Theorem 1]. It follows that we may take $t_1 = \frac{1}{2}$ in (2).

Correlations decay exponentially in the sense that (3) holds with $p(n) = C\tau^n$ for some C and $\tau \in (0,1)$. This is apparent from the proof of the main theorem in [26]. (The actual theorem contains a somewhat weaker statement.)

Hence all assumptions of Theorem 2.9 are satisfied, and we conclude that there is a set A of full μ_a measure in $[1-a, 1]$ such that

$$\dim_{\mathrm{H}}\bigcap_n E_\alpha(x_n) = \frac{1}{\alpha}$$

holds whenever $x_1, x_2, \ldots$ are elements A. Since μ_a is equivalent to Lebesgue measure on $[1-a, 1]$, this proves the theorem. □

Proof of Corollary 2.11. By Corollary 3 of [23], assumption (2) is satisfied with

$$t_1 = \limsup_{m\to\infty}\inf\frac{S_m\phi - mP(\phi)}{-\log|(T^m)'|} > 0.$$

By Theorem 2.9, there is a set A of full μ_ϕ measure such that

$$\frac{1}{\alpha}\frac{\dim_{\mathrm{H}}\,\mu_\phi}{s} = \frac{1}{\alpha}\frac{\overline{\dim}_{\mathrm{H}}\,\mu_\phi}{s} \leq \dim_{\mathrm{H}}\bigcap_k E_\alpha(x_k) \leq \frac{1}{\alpha}$$

holds whenever $x_1, x_2, \ldots$ are elements of the set A. Assumption (4) implies that $\dim_{\mathrm{H}}\,\mu_\phi = s$. □

7.1. **Proof of Theorem 2.9.** Let us now turn to the proof of Theorem 2.9. By a change of variables, we may assume that T is a map of the unit interval $[0,1]$. Since μ is not atomic and the space is one-dimensional, $\mu(R) = 0$ holds.

We shall proceed as in the proof of Theorem 2.8. The setting is similar since

$$E_\alpha(x) = \limsup_n B(T^n(x), n^{-\alpha}),$$

but some further complications occur since the points $T^n(x)$ are not independent. However, the assumption on the decay of correlations gives us enough asymptotic independence to carry out the proof.

As in the proof of Theorem 2.8, we put

$$\Theta_n(t) = \bigcap_{k=n}^{\infty} \{\, x : \mu(B(x,r)) < r^t \text{ for } r = 2^{-k} \,\},$$
$$\Theta(t) = \bigcap_{n=1}^{\infty} \Theta_n(t),$$

and take $t_2 < \overline{\dim}_{\mathrm{H}}\, \mu$, and m_1 so large that $\mu(\Theta_{m_1}(t_2)) > 0$.

We let as before

$$\mathscr{C}_m = \{\, D \in \mathscr{D}_{[\beta m]} : \mu(\Theta_{m_1}(t_2) \cap D) \geq \frac{5}{6}\mu(D) \,\},$$

and there exists a sequence (m_k) such that the set

$$\hat{\Theta} = \Theta_{m_1}(t_2) \cap \bigcap_{k=1}^{\infty} \cup \mathscr{C}_{m_k}.$$

satisfies $\mu(\hat{\Theta}) > \frac{1}{2}\mu(\Theta_{m_1}(t_2)) > 0$. By construction, the set $\hat{\Theta}$ has the following property. If $D \in \mathscr{D}_{[\beta m_k]}$ for some m_k, then either $D \subset \hat{\Theta}^{\complement}$ or

$$\mu(\hat{\Theta} \cap D) \geq \frac{5}{6}\mu(D) - \frac{1}{6}2^{-[\beta m_k] t_0}. \tag{18}$$

Let $\varepsilon > 0$ and $t_0 = \sup\{\, t : G_\mu(t) \geq t - 2\varepsilon \,\}$. Take $\beta > \alpha$ such that $\frac{1}{\beta} \leq t_0$. Let $G = \sup\{\, G_\mu(t) : t \geq t_0 \,\}$. As in the proof of Theorem 2.8, we define a sparse sequence n_k as follows.

For each n, let A_n be the union of those cubes of $\mathscr{D}_n$ which have μ-measure at most 2^{-nt_0}. By Lemma 6.1, there is a number n_0 such that $\mu(A_n) \leq 2^{-\varepsilon n}$ for $n > n_0$.

We choose the sequence n_k so that $\{n_k\} \subset \{m_k\}$ and

$$\frac{1}{4}2^{-[\beta n_k] t_0} > \sum_{l=k+1}^{\infty} 2^{-\varepsilon[\beta n_l]} \tag{19}$$

holds for all $k \geq 1$ and $\beta n_1 > n_0$. Let $B_n = A_n^{\complement}$ and

$$C = \bigcap_{k=1}^{\infty} B_{[\beta n_k]} \cap \hat{\Theta}.$$

We have

$$\mu(C) \geq \mu(\hat{\Theta}) - \sum_{k=1}^{\infty} \mu(A_{[\beta n_k]}) \geq \mu(\hat{\Theta}) - \sum_{k=1}^{\infty} 2^{-\varepsilon[\beta n_k]}$$
$$\geq \mu(\hat{\Theta}) - \sum_{k=[\beta n_1]}^{\infty} 2^{-\varepsilon k} = \mu(\hat{\Theta}) - \frac{2^{-\varepsilon[\beta n_1]}}{1 - 2^{-\varepsilon}}.$$

Hence, if $t_1 < \underline{\dim}_{\mathrm{H}}\, \mu$, by choosing n_1 sufficiently large, we can ensure that $\mu(C) > \frac{1}{2}$, and we can make $\mu(C)$ as close to 1 as we please. Similarly, if $t_1 < \overline{\dim}_{\mathrm{H}}\, \mu$, then we can ensure that $\mu(C) > \frac{1}{2}\mu(\hat{\Theta}) > 0$ by choosing n_1 sufficiently large. We assume that $\mu(C) > \frac{1}{2}\mu(\hat{\Theta})$ holds.

Let f be the indicator function of the set C. Since $\frac{1}{2}\mu(\hat{\Theta}) < \int f \,\mathrm{d}\mu < 1$, we have by Birkhoff's ergodic theorem that for almost every x, there exists a number $N(x)$ such that

$$2^{-(n-1)} \sum_{k=2^{n-1}+1}^{2^n} f(T^k x) > \frac{1}{2}\mu(\hat{\Theta}) \tag{20}$$

for all $n > N(x)$.

Let

$$\mu_n = 2^{-(n-1)} \sum_{k=2^{n-1}+1}^{2^n} \frac{\mu|_{D_{[\beta n]}(T^k x)}}{\mu(D_{[\beta n]}(T^k x))},$$

where $\frac{0}{0}$ should be interpreted as 0. The measure μ_n depends on x, but we suppress this dependence from the notation. If n is large we have

$$\operatorname{supp} \mu_n \subset \bigcup_{k=2^{n-1}+1}^{2^n} B(x, k^{-\alpha}),$$

since $\beta > \alpha$. Moreover, for almost all x, by Birkhoff's theorem, $\mu_n \to \mu$ weakly as $n \to \infty$.

Put $\nu_k = c_k(x)\mu_{n_k}|_C$, where $c_k(x)$ is a constant chosen so that ν_k is a probability measure, that is

$$c_k(x) = \frac{1}{\mu_{n_k}(C)}.$$

This makes $c_k(x)$ well-defined for almost all x if k is larger than some number $K(x)$. In fact, we have by (20) that

$$c_k(x) \in \left[1, \frac{2}{\mu(\hat{\Theta})}\right), \qquad \text{if } n_k > N(x).$$

By Birkhoff's ergodic theorem, we have that $\nu_k \to \nu$ weakly as $k \to \infty$, where $\nu = \frac{\mu|_C}{\mu(C)}$.

We consider the (ν, θ)-energies

$$I_\nu^\theta(\nu_k) = \iint \nu(Q)^{-\theta} \, \mathrm{d}\nu_k \mathrm{d}\nu_k.$$

Our aim is to show that, almost surely, there is a sequence k_l along which $I_\mu^\theta(\nu_{k_l})$ are uniformly bounded.

Given an integer m we define an approximation of the function Q, introduced in Section 2.2. Let $Q_m \colon \mathbb{R}^d \times \mathbb{R}^d \to \mathscr{D}$ be defined by

$$Q_m(x, y) = D, \qquad x \neq y,$$

where $D \in \mathscr{D}_n$ is chosen such that $x, y \in D$ and n is the largest integer with this property such that $n \leq m$. We then have that

$$Q(x, y) = Q_m(x, y)$$

if and only if $Q(x, y) = D$ with $D \in \mathscr{D}_n$ and $n \leq m$.

By the definition of ν_k we can write

$$I_\nu^\theta(\nu_k) = \frac{c_k(x)^2}{4^{n_k - 1}} \sum_{i,j \in J_k} \int_{D_i \cap C} \int_{D_j \cap C} \nu(Q)^{-\theta} \, \frac{\mathrm{d}\mu}{\mu(D_i)} \frac{\mathrm{d}\mu}{\mu(D_j)},$$

where $D_i = D_{[\beta n_k]}(T^i x)$ and $J_k = \{2^{n_k - 1} + 1, \ldots, 2^{n_k}\}$. Assuming that $c_k(x) \leq 2$, which holds almost surely if k is large enough, we therefore have

$$I_\nu^\theta(\nu_k) = 4^{2 - n_k} \sum_{i,j \in J_k} \int_{D_i \cap C} \int_{D_j \cap C} \nu(Q)^{-\theta} \, \frac{\mathrm{d}\mu}{\mu(D_i)} \frac{\mathrm{d}\mu}{\mu(D_j)}. \tag{21}$$

If $\mu(D_i) \leq 2^{-[\beta n_k] t_0}$, then $D_i \subset A_{[\beta n_k]}$, so that $D_i \subset C^{\complement}$ and $\nu(D_i) = 0$. Otherwise, if $\mu(D_i) > 2^{-[\beta n_k] t_0}$ then, either $D_i \subset \hat{\Theta}^{\complement} \cup A_{[\beta n_l]}$ for some $l < k$ and $\nu(D_i) = 0$, or

$$\begin{aligned}
\nu(D) &\geq \mu(C \cap D) \geq \mu(\hat{\Theta} \cap D) - \mu\Bigg(\bigcup_{l=k+1}^{\infty} A_{[\beta n_l]}\Bigg) \\
&\geq \mu(\hat{\Theta} \cap D) - \sum_{l=k+1}^{\infty} 2^{-\varepsilon[\beta n_l]} \\
&\geq \frac{5}{6}\mu(D) - \frac{1}{6} 2^{-[\beta n_k] t_0} - \frac{1}{6} 2^{-[\beta n_k] t_0} \geq \frac{1}{2}\mu(D),
\end{aligned}$$

holds by (19) and (18). In the other direction we have

$$\nu(D_i) = \frac{\mu(C \cap D_i)}{\mu(C)} \leq \frac{2}{\mu(\hat{\Theta})} \mu(D_i).$$

We conclude that either $\nu(D_i) = 0$ or

$$\frac{1}{2} \leq \frac{\nu(D_i)}{\mu(D_i)} \leq \frac{2}{\mu(\hat{\Theta})}. \tag{22}$$

We also have

$$D_i \cap C = \emptyset \qquad \text{or} \qquad \nu(D_i) \geq 2^{-[\beta n_k]t_0 - 1}. \tag{23}$$

Combining (21) and (22), we obtain

$$\begin{aligned} I_\nu^\theta(\nu_k) &\leq \frac{4^{3-n_k}}{\mu(\hat{\Theta})^2} \sum_{i,j \in J_k} \int_{D_i \cap C} \int_{D_j \cap C} \nu(Q)^{-\theta} \frac{\mathrm{d}\mu}{\nu(D_i)} \frac{\mathrm{d}\mu}{\nu(D_j)} \\ &= 4^{3-n_k} \frac{\mu(C)^2}{\mu(\hat{\Theta})^2} \sum_{i,j \in J_k} \int_{D_i} \int_{D_j} \nu(Q)^{-\theta} \frac{\mathrm{d}\nu}{\nu(D_i)} \frac{\mathrm{d}\nu}{\nu(D_j)} \\ &\leq 4^{3-n_k} \sum_{i,j \in J_k} \int_{D_i} \int_{D_j} \nu(Q)^{-\theta} \frac{\mathrm{d}\nu}{\nu(D_i)} \frac{\mathrm{d}\nu}{\nu(D_j)}. \end{aligned}$$

If either $D_i \cap C = \emptyset$ or $D_j \cap C = \emptyset$, then

$$\int_{D_i} \int_{D_j} \nu(Q)^{-\theta} \frac{\mathrm{d}\nu}{\nu(D_i)} \frac{\mathrm{d}\nu}{\nu(D_j)} = 0.$$

Suppose now that both $D_i \cap C \neq \emptyset$ and $D_j \cap C \neq \emptyset$. We consider three cases.

Case 1. If $i = j$ then $D_i = D_j$ and

$$\begin{aligned} \int_{D_i} \int_{D_j} \nu(Q)^{-\theta} \frac{\mathrm{d}\nu}{\nu(D_i)} \frac{\mathrm{d}\nu}{\nu(D_j)} &= \int_{D_i} \int_{D_i} \nu(Q)^{-\theta} \frac{\mathrm{d}\nu}{\nu(D_i)} \frac{\mathrm{d}\nu}{\nu(D_i)} \\ &\leq \frac{1}{1-\theta} \nu(D_i)^{-\theta} \\ &= \frac{1}{1-\theta} \nu(Q_{[\beta n_k]}(T^i x, T^j x))^{-\theta}, \end{aligned}$$

by Lemma 5.2.

Case 2. If $i \neq j$ and $D_i \neq D_j$, then Q is constant and equal to $Q(T^i x, T^j x)$ on $D_i \times D_j$ and we get

$$\int_{D_i} \int_{D_j} \nu(Q)^{-\theta} \frac{\mathrm{d}\nu}{\nu(D_i)} \frac{\mathrm{d}\nu}{\nu(D_j)} = \nu(Q(T^i x, T^j x))^{-\theta}.$$

Case 3. If $i \neq j$ and $D_i = D_j$, then as in the case $i = j$, we get by Lemma 5.2 that

$$\begin{aligned} \int_{D_i} \int_{D_j} \nu(Q)^{-\theta} \frac{\mathrm{d}\nu}{\nu(D_i)} \frac{\mathrm{d}\nu}{\nu(D_j)} &\leq \frac{1}{1-\theta} \nu(D_i)^{-\theta} \\ &= \frac{1}{1-\theta} \nu(Q_{[\beta n_k]}(T^i x, T^j x))^{-\theta}. \end{aligned}$$

Taken together, these estimates show that

$$\int_{D_i} \int_{D_j} \nu(Q)^{-\theta} \frac{\mathrm{d}\nu}{\nu(D_i)} \frac{\mathrm{d}\nu}{\nu(D_j)} \leq \begin{cases} 0 & \text{if } \nu(D_i)\nu(D_j) = 0, \\ \frac{1}{1-\theta} \nu(Q_{[\beta n_k]}(T^i x, T^j x))^{-\theta} & \text{otherwise.} \end{cases}$$

Let

$$F_k(x, y) = \min\{\nu(Q_{[\beta n_k]}(x, y))^{-\theta}, 2^{\theta + \theta t_0 \beta n_k}\}.$$

By (23) we have that

$$\nu(Q_{[\beta n_k]}(x, y))^{-\theta} > 2^{\theta + \theta t_0 \beta n_k} \geq 2^{\theta + \theta t_0 [\beta n_k]}$$

only if $\nu(Q_{[\beta n_k]}(x, y)) = 0$. We therefore have

$$(1-\theta) I_\nu^\theta(\nu_k) \leq 4^{3-n_k} \sum_{i,j \in J} F_k(T^i x, T^j x).$$

This sum is split into two parts according to

$$\begin{aligned} (1-\theta) I_\nu^\theta(\nu_k) &= I_1(k) + I_2(k), \\ I_1(k) &= 4^{3-n_k} \cdot 2 \sum_{\substack{i,j \in J \\ i > j}} F_k(T^i x, T^j x), \\ I_2(k) &= 4^{3-n_k} \sum_{i \in J} F_k(T^i x, T^i x). \end{aligned}$$

We will estimate the expected value of $I_1(k)$. We have

$$\begin{aligned}\mathsf{E}\, I_1(k) &\le 4^{4-n_k} \sum_{\substack{i,j\in J\\ i>j}} \int F_k(T^i x, T^j x)\,\mathrm{d}\mu(x)\\ &= 4^{4-n_k} \sum_{\substack{i,j\in J\\ i>j}} \int F_k(T^{i-j} x, x)\,\mathrm{d}\mu(x).\end{aligned}$$

We will use the decay of correlations to estimate the integrals above. The following lemma which is a variation of Lemma 3 of Persson and Rams [23] will be used.

Lemma 7.1. *Suppose $F\colon [0,1]^2 \to \mathbb{R}$ is a piecewise continuous and non-negative function, and that V and M are two constants such that, for each fixed x, the function $f\colon y \mapsto F(x,y)$ satisfies* $\operatorname{var} f \le V$ *and* $\int f\,\mathrm{d}\mu \le M$. *Then*

$$\int F(T^n x, x)\,\mathrm{d}\mu(x) \le M + (V+M)p(n).$$

We will use Lemma 7.1 with $F = F_k$. Clearly, F_k is piecewise constant and hence piecewise continuous. Moreover, F_k is bounded by $2^{\theta+\theta t_0[\beta n_k]}$ by (23).

Let x be fixed and consider the function $f_x\colon y \mapsto F_k(x,y)$. The function f_x is increasing on $[0,x]$ and decreasing on $[x,1]$. Hence, f_x is of bounded variation, and

$$\operatorname{var} f_x \le 2^{1+\theta+\theta t_0[\beta n_k]}.$$

By (2), we have

$$\int f_x\,\mathrm{d}\mu \le \sum_{k=0}^{[\beta n]} \mu(D_k(x))^{1-\theta} \le M := \sum_{k=0}^{\infty} c_1^{1-\theta} 2^{-(1-\theta)t_1 k} < \infty.$$

It therefore follows by Lemma 7.1 that

$$\int F_k(T^{i-j}x, x)\,\mathrm{d}\mu(x) \le M + (V(k)+M)p(i-j).$$

Consequently, using that p is summable, we obtain

$$\begin{aligned}\mathsf{E}\, I_1(k) &\le 4^{4-n_k} \sum_{i=2^{n_k-1}+2}^{2^{n_k}} \sum_{j=2^{n_k-1}+1}^{i-1} \big(M + (V(k)+M)p(i-j)\big)\\ &= 4^{4-n_k} \sum_{i=2^{n_k-1}}^{2^{n_k}} \big(2^{n_k}M + (V(k)+M) \sum_{j=2^{n_k-1}+1}^{i-1} p(i-j)\big)\\ &\le 4^{4-n_k} \sum_{i=2^{n_k-1}}^{2^{n_k}} \big(2^{n_k}M + (V(k)+M) \sum_{j=0}^{\infty} p(j)\big)\\ &\le 16M + 16\cdot 2^{-n_k}(V(k)+M)\sum_{i=1}^{\infty}(i)\\ &\le 16M + c_0 2^{(\beta\theta t_0 - 1)n_k},\end{aligned}$$

where c_0 is a constant. Hence, $\mathsf{E}\, I_1(k)$ is uniformly bounded in k, provided $\theta \le \frac{1}{\beta t_0}$.

To analyse $I_2(k)$, we write

$$\begin{aligned}I_2(k) = 4^{3-n_k}\sum_{i\in J} F_k(T^i x, T^i x) &\le 4^{3-n_k}\sum_{i\in J} 2^{\theta+\theta t_0 \beta n_k}\\ &\le 4^{3-n_k} 2^{n_k} 2^{\theta+\theta\beta t_0 n_k} \le c_1 2^{(\theta\beta t_0 - 1)n_k}.\end{aligned}$$

Hence, $I_2(k)$ is bounded by c_1 if $\theta \le \frac{1}{\beta t_0}$.

Take $\theta < \frac{1}{\beta t_0} \le 1$. Then $I_2(k)$ is bounded by c_1 for all k and the expectation of $I_1(k)$ is bounded by $16M + c_0$ for all k. We can conclude that for almost all x, that is almost surely, there exists a subsequence k_l and a constant c_2 such that $I_1(k_l) < c_2$. We then have that

$$I_\nu^\theta(\nu_{k_l}) \le \frac{1}{1-\theta}(16M + c_0 + c_2)$$

for all l. Lemma 2.7 now implies that $E_\alpha(x) \in \mathscr{G}_\nu^\theta$.

From the statement that $E_\alpha(x) \in \mathscr{G}_\nu^\theta$ holds for a.e. x, we conclude in the same way as in the proof of Theorem 2.8 that $E_\alpha(x) \in \mathscr{G}_\mu^\theta$ holds for a.e. x, with $\theta = (s\alpha)^{-1}$. (We use that $\mu(C)$ can be made arbitrarily close to 1 and that β and t_0 can be taken as close to α and s as we please.)

Finally, as in the proof of Theorem 2.8, by taking t_2 close to $\overline{\dim}_{\mathrm{H}}\,\mu$, we find that

$$\dim_{\mathrm{H}} \bigcap_k E_\alpha(x_k) \geq \frac{1}{\alpha}\frac{\overline{\dim}_{\mathrm{H}}\,\mu}{s}$$

whenever the points $x_1, x_2, \ldots$ are elements of a set A of full μ measure.

References

[1] Aspenberg, Magnus; Persson, Tomas Shrinking targets in parametrised families, Math. Proc. Camb. Philos. Soc., Volume 166 (2019) no. 2, pp. 265-295 | MR | DOI | Zbl

[2] Benedicks, Michael; Carleson, Lennart The dynamics of the Hénon map, Ann. Math., Volume 133 (1991) no. 1, pp. 73-169 | DOI | Zbl

[3] Bugeaud, Yann; Liao, Lingmin Uniform diophantine approximation related to b-ary and β-expansions, Ergodic Theory Dyn. Syst., Volume 36 (2016) no. 1, pp. 1-22 | MR | DOI | Zbl

[4] Bugeaud, Yann; Wang, Bao-Wei Distribution of full cylinders and the diophantine properties of the orbits in β-expansions, J. Fractal Geom., Volume 1 (2014) no. 2, pp. 221-241 | MR | DOI | Zbl

[5] Durand, Arnaud On randomly placed arcs on the circle, Recent developments in fractals and related fields (Applied and Numerical Harmonic Analysis), Birkhäuser, 2010, 343–351 pages | MR | DOI | Zbl

[6] Ekström, Fredrik; Persson, Tomas Hausdorff dimension of random limsup sets, J. Lond. Math. Soc., Volume 98 (2018) no. 3, pp. 661-686 | MR | DOI | Zbl

[7] Falconer, Kenneth J. Classes of sets with large intersection, Mathematika, Volume 32 (1985), pp. 191-205 | MR | DOI | Zbl

[8] Falconer, Kenneth J. Sets with large intersection properties, J. Lond. Math. Soc., Volume 49 (1994) no. 2, pp. 267-280 | MR | DOI | Zbl

[9] Falconer, Kenneth J. Techniques in fractal geometry, John Wiley & Sons, 1997, xvii+256 pages | Zbl

[10] Fan, Ai-Hua; Feng, De-Jun; Wu, Jun Recurrence, dimension and entropy, J. Lond. Math. Soc., Volume 64 (2001) no. 1, pp. 229-244 | MR | DOI | Zbl

[11] Feng, De-Jun; Järvenpää, Esa; Järvenpää, Maarit; Suomala, Ville Dimensions of random covering sets in Riemann manifolds, Ann. Probab., Volume 46 (2018) no. 3, pp. 1201-1805 | MR | Zbl

[12] Frostman, Otto Potentiel d'équilibre et capacité des ensembles avec quelques applications a la théorie des fonctions, Meddelanden från Lunds Universitets Matematiska Seminarium, 3, 1935 | Zbl

[13] Fan, Ai-Hua; Schmeling, Jörg; Troubetzkoy, Serge A multifractal mass transference principle for Gibbs measures with applications to dynamical diophantine approximation, Proc. Lond. Math. Soc., Volume 107 (2013) no. 5, pp. 1173-1219 | MR | DOI | Zbl

[14] Fan, Ai-Hua; Wu, Jun On the covering by small random intervals, Ann. Inst. Henri Poincaré, Probab. Stat., Volume 40 (2004) no. 1, pp. 125-131 | Numdam | MR | DOI | Zbl

[15] Hill, Richard; Velani, Sanju L. The ergodic theory of shrinking targets, Invent. Math., Volume 119 (1995) no. 1, pp. 175-198 | MR | DOI | Zbl

[16] Järvenpää, Esa; Järvenpää, Maarit; Koivusalo, Henna; Li, Bing; Suomala, Ville Hausdorff dimension of affine random covering sets in torus, Ann. Inst. Henri Poincaré, Probab. Stat., Volume 50 (2014) no. 4, pp. 1371-1384 | MR | Numdam | DOI | Zbl

[17] Liao, Lingmin; Seuret, Stéphane Diophantine approximation by orbits of expanding Markov maps, Ergodic Theory Dyn. Syst., Volume 33 (2013) no. 2, pp. 585-608 | MR | DOI | Zbl

[18] Liverani, Carlangelo; Soussol, Benoit; Vaienti, Sandro Conformal measure and decay of correlation for covering weighted systems, Ergodic Theory Dyn. Syst., Volume 18 (1998) no. 6, pp. 1399-1420 | DOI | MR | Zbl

[19] Li, Bing; Wang, Bao-Wei; Wu, Jun; Xu, Jian The shrinking target problem in the dynamical system of continued fractions, Proc. Lond. Math. Soc., Volume 108 (2014) no. 1, pp. 159-186 | MR | Zbl

[20] Olsen, Lars A multifractal formalism, Adv. Math., Volume 116 (1995) no. 1, pp. 82-196 | MR | DOI | Zbl

[21] Persson, Tomas A note on random coverings of tori, Bull. Lond. Math. Soc., Volume 47 (2015) no. 1, pp. 7-12 | MR | DOI | Zbl

[22] Persson, Tomas; Reeve, Henry W. J. A Frostman-type lemma for sets with large intersections, and an application to diophantine approximation, Proc. Edinb. Math. Soc., Volume 58 (2015) no. 2, pp. 521-542 | MR | DOI | Zbl

[23] Persson, Tomas; Rams, Michał On shrinking targets for piecewise expanding interval maps, Ergodic Theory Dyn. Syst., Volume 37 (2017) no. 2, pp. 646-663 | MR | DOI | Zbl

[24] Rogers, Claude A. Hausdorff measures, Cambridge University Press, 1970 | Zbl

[25] Seuret, Stéphane Inhomogeneous coverings of topological Markov shifts, Math. Proc. Camb. Philos. Soc., Volume 165 (2018) no. 2, pp. 341-357 | MR | DOI | Zbl

[26] Young, Lai-Sang Decay of correlations for certain quadratic maps, Commun. Math. Phys., Volume 146 (1992) no. 1, pp. 123-138 | MR | DOI | Zbl