




FIGURE 5. The proof that $\mathbb{M}$ is dense goes as follows.




(a) We first cover X by topologically trivial open sets (here Y_1, Y_2, Y_3) on which the degeneracies of H reduce to those of a 2×2 system.



(b) In Y_1 , the degenerate part of H reduces to that of a 2×2 system. Via the procedure of §3.1, we can produce H_1 , arbitrarily close to H , with no bad points in Y_1 . By Lemma 3.3, $\mathcal{B}(H_1)$ is a small perturbation of $\mathcal{B}(H) \setminus Y_1$.



(c) We repeat the procedure and produce H_2 , arbitrarily close to H_1 , with no bad points in Y_2 . As bad points are stable, $\mathcal{B}(H_2)$ is close to $\mathcal{B}(H_1)$. In particular passing from H_1 to H_2 does not generate bad points back in $Y_1 \setminus (Y_2 \cup Y_3)$, and removes bad points in Y_2 .



(d) We get new systems H_1, H_2, H_3 , recursively constructed, arbitrarily close to H , with no bad points in $Y_1, Y_1 \cup Y_2 \setminus Y_3, Y_1 \cup Y_2 \cup Y_3$, respectively. Since $Y_1 \cup Y_2 \cup Y_3$ cover X , H_3 is in $\mathbb{M}$ and is arbitrarily close to H .