

1. INTRODUCTION

Let ξ be a non-zero real number and let n be a positive integer. Dirichlet's theorem (1842) is one of the most basic results of Diophantine approximation. It shows that for any real number $H > 1$, there exists a non-zero integer point $(x_0, \dots, x_n) \in \mathbb{Z}^{n+1}$ such that

$$(1.1) \quad \max\{|x_1|, \dots, |x_n|\} \leq H \quad \text{and} \quad |x_0 + x_1\xi + \dots + x_n\xi^n| \leq H^{-n}.$$

It is natural to ask if we can improve the exponent n of H^{-n} , and this question gives rise to two Diophantine exponents. The so-called *uniform exponent of approximation* $\widehat{\omega}_n(\xi)$ (resp. the *ordinary exponent* $\omega_n(\xi)$), is the supremum of the real numbers $\omega > 0$ such that the system

$$\|P\| \leq H \quad \text{and} \quad 0 < |P(\xi)| \leq H^{-\omega}$$

admits a non-zero solution $P \in \mathbb{Z}[X]$ of degree at most n for each sufficiently large H (resp. for arbitrarily large H). Here, $\|P\|$ denotes the (naive) *height* of P , defined as the largest absolute value of its coefficients. These quantities have been extensively studied over the past half-century, see for example [5] for a nice overview of the subject. By Dirichlet's theorem, if ξ is not an algebraic number of degree $\leq n$, then we have

$$\omega_n(\xi) \geq \widehat{\omega}_n(\xi) \geq n,$$

and it is well known that those inequalities are equalities for almost all real numbers ξ (with respect to the Lebesgue measure). Note that if ξ is an algebraic number of degree d , then $\widehat{\omega}_n(\xi)$ and $\omega_n(\xi)$ are both equal to $\min\{n, d-1\}$ (it is a consequence of Schmidt's subspace theorem, see [5, Th. 2.10]). We can therefore restrict our study to the set of transcendental real numbers. The initial question “can we improve the exponent n in Dirichlet's theorem?” may be rephrased as follows: “does there exist a transcendental real number ξ satisfying $\widehat{\omega}_n(\xi) > n$?”. For $n = 1$ the answer is negative and rather elementary to prove, so the first non-trivial case is $n = 2$. Before the early 2000s, it was conjectured that no such number existed. This belief was swept away by Roy's extremal numbers [20], [21], [1], whose exponent $\widehat{\omega}_2$ is equal to the maximal possible value $(3 + \sqrt{5})/2 = 2.618\dots$. Since then, several families of transcendental real numbers whose uniform exponent $\widehat{\omega}_2$ is greater than 2 have been discovered (see for example [22], [6], [15, 16]). However, for $n \geq 3$ the mystery remains, and it is still an open question whether or not there exists $\xi \in \mathbb{R} \setminus \overline{\mathbb{Q}}$ with $\widehat{\omega}_n(\xi) > n$.

In this paper, we are interested in finding an upper bound for the uniform exponent $\widehat{\omega}_n(\xi)$, as this could provide clues to solving the initial problem. Brownawell's version of Gel'fond's criterion [3] implies that $\widehat{\omega}_n(\xi) \leq 3n$. In 1969, Davenport and Schmidt [10, Th. 2b] showed that for any transcendental real number ξ and any integer $n \geq 2$, we have

$$(1.2) \quad \widehat{\omega}_n(\xi) \leq 2n - 1.$$

Up to now, few improvements have been made. Bugeaud and Schleischitz [8, Th. 2.1] first got the upper bound

$$(1.3) \quad \widehat{\omega}_n(\xi) \leq n - \frac{1}{2} + \sqrt{n^2 - 2n + 1/4} = 2n - \frac{3}{2} + \varepsilon_n,$$

where $\varepsilon_n > 0$ tends to 0 as n tends to infinity. Recently, Marnat and Moshchevitin [13] proved an important conjecture of Schmidt and Summerer on the ratio $\widehat{\omega}_n(\xi)/\omega_n(\xi)$ (see also [19, Ch. 2] for an alternative proof based on parametric geometry of numbers). In [23], Schleischitz pointed out that we can use the aforementioned inequality in the proof of (1.3) to get

$$\widehat{\omega}_n(\xi) \leq 2n - 2,$$

for each $n \geq 10$. This is currently the best known upper bound. Let us also mention that using parametric geometry of numbers, Schleischitz [24, Th. 1.1] was able to replace the estimate (1.3) by

$$\widehat{\omega}_n(\xi) \leq \frac{3(n-1) + \sqrt{n^2 - 2n + 5}}{2} = 2n - 2 + \varepsilon'_n$$

where $\varepsilon'_n > 0$ tends to 0 as n tends to infinity. For $n = 3, \dots, 9$, bounds that are better than (1.2), but nevertheless (strictly) greater than $2n - 2$, are known. For example, it was proved in [8] that for each transcendental real number ξ , we have

$$\widehat{\omega}_3(\xi) \leq 3 + \sqrt{2} = 4.41 \dots,$$

see also the very recent work of Schleischitz [25]. In this paper, without relying on Marnat-Moshchevitin's inequality and with a different approach, we show in Section 7 that the upper bound $\widehat{\omega}_n(\xi) \leq 2n - 2$ holds for any $n \geq 4$. We also improve the upper bound for $\widehat{\omega}_3$.

THEOREM 1.1. — *Let $n \geq 3$ be an integer and $\xi \in \mathbb{R}$ be a transcendental real number. If $n \geq 4$, then*

$$\widehat{\omega}_n(\xi) \leq 2n - 2.$$

For $n = 3$, we have the weaker estimate $\widehat{\omega}_3(\xi) \leq 2 + \sqrt{5} = 4.23 \dots$.

We do not think that these upper bounds are optimal. It is interesting to note that Schleischitz, with a different method and under a technical condition, also found the estimates of Theorem 1.1, see [25]. Our main result below is a significant improvement on the previous results as n tends to infinity and does not require Marnat and Moshchevitin's inequality [13].

THEOREM 1.2. — *Set $a = 1/3$. There exists a computable constant $N \geq 1$ such that, for each $n \geq N$ and any transcendental real number $\xi \in \mathbb{R}$, we have*

$$\widehat{\omega}_n(\xi) \leq 2n - an^{1/3}.$$

The constant $a = 1/3$ is not optimal. Numerical calculations based on the results from Section 11 suggest that we could take N rather “small” in Theorem 1.2 (maybe $N \leq 10^4$). However, to keep the arguments and calculations as clear and simple as possible, we did not try to provide an explicit value of N .

Theorem 1.2 can be compared to [17, Th. 1.1], where we study $\widehat{\lambda}_n(\xi)$, the uniform exponent of rational simultaneous approximation to the successive powers $\Xi = (1, \xi, \xi^2, \dots, \xi^n)$ (which is known to be, in a sense, dual to $\widehat{\omega}_n(\xi)$), see Section 2 for the precise definition and more details. We were not able to deduce one result from the other, even though there are similarities in the arguments. For example, given a polynomial $P \in \mathbb{Z}[X]$ of degree at most n , which is a good approximation, we can associate the $k + 1$ polynomials $P, XP, \dots, X^k P$ of degree at most $n + k$. They provide information on $\widehat{\omega}_{n+k}(\xi)$. On the other hand, if we consider $\mathbf{y} \in \mathbb{Z}^{n+1}$ which is a good approximation of Ξ (for simultaneous approximation), we can associate the $k + 1$ blocks of successive $n + 1 - k$ coordinates of $\mathbf{y}$, which are rather good approximations of $(1, \xi, \dots, \xi^{n-k})$. They in turn provide information on $\widehat{\lambda}_{n-k}(\xi)$. Note that the difficulties in the proofs of both theorems are not in the same places. In particular, in this paper we have to work with *irreducible* polynomials, a rather heavy constraint. Also, one of the most delicate parts of our approach is to bound from above the ordinary exponent $\omega_n(\xi)$, whereas this is rather “simple” to do for the ordinary exponent $\lambda_n(\xi)$ in [17].

Before presenting our strategy, let us quickly explain Davenport and Schmidt’s proof of the upper bound (1.2). Given a real number $\widehat{\omega} < \widehat{\omega}_n(\xi)$, they show, using elementary means and Gelfond’s lemma, that there are infinitely many pairs of coprime polynomials $P, Q \in \mathbb{Z}[X]$ of degree at most n , such that

$$\|Q\| \leq \|P\| \quad \text{and} \quad \max\{|Q(\xi)|, |P(\xi)|\} \ll \|P\|^{-\widehat{\omega}},$$

(where the implicit constant only depends on n). It implies that the resultant $\text{Res}(P, Q)$, which is a non-zero integer, satisfies

$$1 \leq |\text{Res}(P, Q)| \ll \|P\|^{n-1} \|Q\|^{n-1} \max\{\|P\| |Q(\xi)|, \|Q\| |P(\xi)|\} \ll \|P\|^{2n-1-\widehat{\omega}}.$$

The first upper bound for $|\text{Res}(P, Q)|$ is classical, see Lemma 4.1. Since $\|P\|$ can be arbitrarily large, they deduced that the exponent $2n - 1 - \widehat{\omega}$ is non-negative. Estimate (1.2) follows by letting $\widehat{\omega}$ tend to $\widehat{\omega}_n(\xi)$. Note that the term $2n$ in (1.2) is directly related to the size of the $2n \times 2n$ determinant defining $\text{Res}(P, Q)$ (if we suppose that P and Q have degree exactly n).

The key idea in the proof of our main Theorem 1.2 is to work with a large number of “good” linearly independent polynomial approximations $Q_0, \dots, Q_{j+1}$ rather than just two polynomials P and Q as above. By doing this, we can replace $\text{Res}(P, Q)$ by a non-zero $(2n - j) \times (2n - j)$ determinant depending on the coefficients of $Q_0, \dots, Q_{j+1}$. Under the ideal and unlikely assumption that

$$(1.4) \quad \|Q_k\| \leq \|Q_0\| \quad \text{and} \quad |Q_k(\xi)| \ll \|Q_0\|^{-\widehat{\omega}} \quad (\text{for } k = 0, \dots, j),$$

the aforementioned determinant would be bounded from above by $\|Q_0\|^{2n-j-1-\widehat{\omega}}$. So, together with an additional non-vanishing assumption, it would lead to $\widehat{\omega}_n(\xi) \leq 2n - j - 1$. Several new difficulties arise when trying to make the above arguments work. We introduce the tools for the construction of the generalized resultant in Section 6. To ensure that this determinant does not vanish, we need the extra assumption that

$Q_0, \dots, Q_{j+1}$ are irreducible polynomials. The idea is to first fix a sequence of best approximations, that we call *minimal polynomials*, and then to consider their highest-degree irreducible factors (which also happen to be rather good approximations). We deal with this question in Section 5. Two obstacles remain. Firstly, note that it may be possible that the best polynomial approximations span a subspace of dimension 3, even when ξ is transcendental and n is large, see [14, Th. 1.3]. Therefore, as soon as $j > 1$ (we will later choose $j \asymp n^{1/3}$), we have to justify that we can find $j + 2$ linearly independent polynomials as above. The second major problem is the control of the sequence $Q_0, \dots, Q_{j+1}$. Estimates (1.4) seem out of reach, instead we get upper bounds of the form

$$(1.5) \quad \|Q_k\| \leq \|Q_0\| \quad \text{and} \quad |Q_k(\xi)| \ll \|Q_0\|^{-\widehat{\omega}_n \theta} \quad (\text{for } k = 0, \dots, j),$$

where $\theta < 1$ depends only on n and j , and is “close” to 1 if j is not too large compared to n . The main ingredients for showing this are related to *twisted heights*, see Sections 8.2 and the appendix, and an important inequality on the height of subspaces due to Schmidt. The parameter θ in (1.5) is a function of the exponent of best approximation $\omega_n(\xi)$. We show in Section 10 that if the uniform exponent satisfies $\widehat{\omega}_n(\xi) \geq 2n - d$ (with $d \ll n^{1/3}$), then the ordinary exponent $\omega_n(\xi)$ is bounded from above by $2n + 2d^2$, and the ratio $\widehat{\omega}_n(\xi)/\omega_n(\xi)$ is therefore close to 1. This part, which is essentially independent from the others, is rather delicate, because we work with the polynomials Q_i . They are certainly irreducible, but not as good approximations as the minimal polynomials. More precisely, there could be large gaps between the height of two successive Q_i . If we could drop the irreducibility condition and directly work with the sequence of minimal polynomials, we could possibly replace the upper bound $2n - \mathcal{O}(n^{1/3})$ with $2n - \mathcal{O}(n^{1/2})$ in Theorem 1.2. Section 11 is devoted to the proof of Theorem 1.2.

Acknowledgements. — I would like to thank Damien Roy for his attentive reading of this work and his many comments, which helped to improve its overall presentation and clarity. I also thank the referee for their work and valuable comments.

2. NOTATION

Throughout this paper, ξ denotes a transcendental real number. The floor (resp. ceiling) function is denoted by $\lfloor \cdot \rfloor$ (resp. $\lceil \cdot \rceil$). If $f, g : I \rightarrow [0, +\infty)$ are two functions on a set I , we write $f = \mathcal{O}(g)$ or $f \ll g$ or $g \gg f$ to mean that there is a positive constant c such that $f(x) \leq cg(x)$ for each $x \in I$. We write $f \asymp g$ when both $f \ll g$ and $g \ll f$ hold.

Let K be a field. If $\mathcal{A}$ is a subset of a K -vector space V , we denote by $\langle \mathcal{A} \rangle_K \subseteq V$ the K -vector space spanned by $\mathcal{A}$, with the convention that $\langle \emptyset \rangle_K = \{0\}$.

Given a ring A (typically $A = \mathbb{R}$ or $\mathbb{Z}$) and an integer $n \geq 0$, we denote by $A[X]$ the ring of polynomials in X with coefficients in A , and by $A[X]_{\leq n} \subseteq A[X]$ the subgroup of polynomials of degree at most n . We say that $P \in \mathbb{Z}[X]$ is *primitive* if it non-zero and

the greatest common divisor of its coefficients is 1. Given $P = \sum_{k=0}^n a_k X^k \in \mathbb{R}[X]$, we set

$$\|P\| = \max_{0 \leq k \leq n} |a_k|.$$

Gelfond's lemma is the following statement (see e.g. [4, Lem. A.3] as well as [3]). For any non-zero polynomials $P_1, \dots, P_r \in \mathbb{R}[X]$ with product $P = P_1 \cdots P_r$ of degree at most n , we have

$$(2.1) \quad e^{-n} \|P_1\| \cdots \|P_r\| < \|P\| < e^n \|P_1\| \cdots \|P_r\|.$$

In particular, for each non-zero polynomial $P \in \mathbb{Z}[X]_{\leq n}$ and each factor $Q \in \mathbb{Z}[X]$ of P , we have $e^{-n} \|Q\| < \|P\|$. We will often use (2.1) as follows. If $Q \in \mathbb{Z}[X]_{\leq n}$ is irreducible and if $P \in \mathbb{Z}[X]_{\leq n}$ is a non-zero polynomial which satisfies $\|P\| \leq e^{-n} \|Q\|$, then Q cannot divide P . They are thus coprime polynomials.

We recall the definition of the resultant, which, as explained in the introduction, is useful for estimating the exponent $\hat{\omega}_n(\xi)$ (see also Section 4). Let $P, Q \in \mathbb{Z}[X]$ be non-constant polynomials of degree p and q respectively, and let $a_i, b_j \in \mathbb{Z}$ such that $P(X) = \sum_{k=0}^p a_k X^k$ and $Q(X) = \sum_{k=0}^q b_k X^k$. Their *resultant* $\text{Res}(P, Q)$ is defined as the $(q+p)$ -dimensional determinant

$$(2.2) \quad \text{Res}(P, Q) = \begin{vmatrix} a_p & 0 & \cdots & b_q & 0 & \cdots \\ a_{p-1} & a_p & & b_{q-1} & b_q & \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots \\ a_0 & & & b_0 & & \\ 0 & a_0 & & 0 & b_0 & \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots \\ & & a_0 & & & b_p \end{vmatrix}.$$

$\underbrace{\hspace{10em}}_q \qquad \underbrace{\hspace{10em}}_p$

Besides the exponents of linear approximation ω_n and $\hat{\omega}_n$, we will also need the following exponents of simultaneous rational approximation. For each positive integer n , the exponent $\hat{\lambda}_n(\xi)$ (resp. $\lambda_n(\xi)$) is the supremum of the real numbers $\lambda \geq 0$ such that the system

$$|y_0| \leq Y \quad \text{and} \quad L(\mathbf{y}) \leq Y^{-\lambda} \quad \text{where} \quad L(\mathbf{y}) := \max_{1 \leq k \leq n} |y_0 \xi^k - y_k|,$$

admits a non-zero integer solution $\mathbf{y} = (y_0, \dots, y_n) \in \mathbb{Z}^{n+1}$ for each sufficiently large $Y \geq 1$ (resp. for arbitrarily large Y). Dirichlet's theorem [26, §II.1, Th. 1A] implies that $\hat{\lambda}_n(\xi) \geq 1/n$. The best upper bounds known to date for $\hat{\lambda}_n(\xi)$ when $n \geq 4$ are established in joint work with Roy in [17]. In particular, there is an explicit positive constant a such that

$$\hat{\lambda}_n(\xi) \leq \frac{1}{n/2 + an^{1/2} + 1/3},$$

and sharper results are also obtained when n is small.

3. MINIMAL POLYNOMIALS

A *sequence of minimal polynomials* (associated to n and ξ) is a sequence $(P_i)_{i \geq 0}$ of non-zero polynomials in $\mathbb{Z}[X]_{\leq n}$ satisfying the following properties:

- (i) the sequence $(\|P_i\|)_{i \geq 0}$ is strictly increasing;
- (ii) the sequence $(|P_i(\xi)|)_{i \geq 0}$ is strictly decreasing;
- (iii) if $|P(\xi)| < |P_i(\xi)|$ for some index $i \geq 0$ and a non-zero $P \in \mathbb{Z}[X]_{\leq n}$, then $\|P\| \geq \|P_{i+1}\|$.

Note that if we require the dominant coefficient of P_i to be positive (and since ξ is transcendental), then the above sequence is unique up to its first terms. Let $(P_i)_{i \geq 0}$ be a sequence as above. We have the classical formulas:

$$(3.1) \quad \widehat{\omega}_n(\xi) = \liminf_{i \rightarrow \infty} \frac{-\log |P_i(\xi)|}{\log \|P_{i+1}\|} \quad \text{and} \quad \omega_n(\xi) = \limsup_{i \rightarrow \infty} \frac{-\log |P_i(\xi)|}{\log \|P_i\|}.$$

In particular, given a positive real number $\widehat{\omega}$ with $\widehat{\omega} < \widehat{\omega}_n(\xi)$, then we have, for each sufficiently large index i ,

$$(3.2) \quad |P_i(\xi)| \leq \|P_{i+1}\|^{-\widehat{\omega}} \quad \text{and} \quad \|P_{i+1}\|^\tau \leq \|P_i\|, \quad \text{where } \tau := \frac{\widehat{\omega}}{\omega_n(\xi)},$$

(with the convention $\tau = 0$ if $\omega_n(\xi) = \infty$). The second inequality in (3.2) asks for an upper bound on $\omega_n(\xi)$. Given a non-zero $P \in \mathbb{Z}[X]$, we set $\omega(P) = 0$ if $\|P\| = 1$. Otherwise, we denote by $\omega(P)$ the real number satisfying

$$|P(\xi)| = \|P\|^{-\omega(P)}.$$

With this notation, we have

$$(3.3) \quad \omega_n(\xi) = \limsup_{\substack{\|P\| \rightarrow \infty \\ P \in \mathbb{Z}[X]_{\leq n}}} \omega(P) = \limsup_{i \rightarrow \infty} \omega(P_i) \quad \text{and} \quad \liminf_{i \rightarrow \infty} \omega(P_i) \geq \widehat{\omega}_n(\xi).$$

The following results are well-known. We prove them for the sake of completion. The first one follows from the arguments of the proof of [9, Lem. 2] (see also [21, Lem. 4.1]).

LEMMA 3.1. — *Let $i \geq 0$ and write $V_i = \langle P_i, P_{i+1} \rangle_{\mathbb{R}} \subseteq \mathbb{R}[X]_{\leq n}$. Then $\{P_i, P_{i+1}\}$ forms a $\mathbb{Z}$ -basis of the lattice $V_i \cap \mathbb{Z}[X]_{\leq n}$.*

Proof. — By contradiction, suppose that $\{P_i, P_{i+1}\}$ is not a $\mathbb{Z}$ -basis of $V_i \cap \mathbb{Z}[X]_{\leq n}$. Then there exists a non-zero $Q \in \mathbb{Z}[X]_{\leq n}$ which may be written as $Q = rP_i + sP_{i+1}$, where $r, s \in \mathbb{Q}$ satisfy $|r|, |s| \leq 1/2$. In particular, we have

$$\begin{aligned} \|Q\| &\leq |r|\|P_i\| + |s|\|P_{i+1}\| < \|P_{i+1}\|, \\ |Q(\xi)| &\leq |r||P_i(\xi)| + |s||P_{i+1}(\xi)| < |P_i(\xi)|. \end{aligned}$$

This contradicts the minimality property of P_i . □

The next result is analogous to the second part of [21, Lem. 4.1]. The construction of S_i is due to Davenport and Schmidt [9].

LEMMA 3.2. — For each $i \geq 0$, define

$$S_i = P_i(\xi)P_{i+1} - P_{i+1}(\xi)P_i \in \mathbb{R}[X]_{\leq n}.$$

Then

$$\frac{1}{2}\|S_i\| \leq \|P_{i+1}\| |P_i(\xi)| \leq 2\|S_i\|.$$

Moreover, if for integers $0 \leq i < j$ the space spanned by $P_i, P_{i+1}, \dots, P_j$ has dimension 2, then $S_{j-1} = \pm S_i$. In particular

$$\|P_{i+1}\| |P_i(\xi)| \asymp \|P_j\| |P_{j-1}(\xi)|.$$

REMARK 3.3. — Note that the quantity $\|S_i\|$ satisfies $\|S_i\| \asymp \mathcal{D}_\xi(V_i)$, where $\mathcal{D}_\xi$ is defined in Section 8.2 and $V_i = \langle P_i, P_{i+1} \rangle_{\mathbb{R}}$. We will study the function $\mathcal{D}_\xi$ more deeply later.

Proof. — We easily get $\|S_i\| \leq 2\|P_{i+1}\| |P_i(\xi)|$. Define $R_+, R_- \in \mathbb{Z}[X]_{\leq n}$ by

$$R_{\pm} = P_{i+1} \pm P_i.$$

Suppose that there exists $\varepsilon \in \{+, -\}$ such that $|R_\varepsilon(\xi)| \leq |P_i(\xi)|/2$. Then by minimality of P_i , we must have $\|R_\varepsilon\| \geq \|P_{i+1}\|$. Since $S_i = P_i(\xi)R_\varepsilon - R_\varepsilon(\xi)P_i$, we find

$$\|S_i\| \geq |P_i(\xi)| \|R_\varepsilon\| - |R_\varepsilon(\xi)| \|P_i\| \geq \frac{1}{2} \|P_{i+1}\| |P_i(\xi)|.$$

Assume that $|R_+(\xi)|, |R_-(\xi)| \geq |P_i(\xi)|/2$. This is equivalent to

$$|P_{i+1}(\xi)| \leq \frac{1}{2} |P_i(\xi)|.$$

Again, this yields $\|S_i\| \geq |P_i(\xi)| \|P_{i+1}\| - |P_{i+1}(\xi)| \|P_i\| \geq \|P_{i+1}\| |P_i(\xi)|/2$.

Now, let us write $V_i = \langle P_i, \dots, P_j \rangle_{\mathbb{R}}$, with $j > i$, and suppose that V_i has dimension 2. We need to prove that $S_{j-1} = \pm S_i$. If $j = i + 1$ it is automatic, we may therefore assume that $j \geq i + 2$. By Lemma 3.1, there exist $a, b \in \mathbb{Z}$ such that $P_i = aP_{i+1} + bP_{i+2}$. Since $\{P_i, P_{i+1}\}$ is also a $\mathbb{Z}$ -basis of V_i , we have $b = \pm 1$, and we deduce that

$$S_i = (aP_{i+1}(\xi) + bP_{i+2}(\xi))P_{i+1} - P_{i+1}(\xi)(aP_{i+1} + bP_{i+2}) = -bS_{i+1} = \pm S_{i+1}.$$

By induction, we get $S_i = \pm S_{i+1} = \dots = \pm S_{j-1}$. $\square$

The proof of [9, Lem. 3] (which deals with the case $n = 2$) yields the classical following result.

LEMMA 3.4. — Suppose $n \geq 2$. Then, there are infinitely many indices $i \geq 1$ for which P_{i-1} , P_i and P_{i+1} are linearly independent.

Proof. — By contradiction, suppose that there exists $i \geq 0$ such that $V = \langle P_i, P_{i+1}, \dots \rangle_{\mathbb{R}}$ has dimension 2. By Lemma 3.2 there exists $c > 0$ such that for each $j > i$ we have

$$0 < \|P_{i+1}\| |P_i(\xi)| \leq c \|P_j\| |P_{j-1}(\xi)|.$$

This leads to a contradiction since $\|P_j\| \|P_{j-1}\| \leq \|P_j\|^{1-\widehat{\omega}_n(\xi)+o(1)}$ tends to 0 as j tends to infinity. $\square$

REMARK 3.5. — As mentioned in the introduction, it is however possible that all polynomials P_i with i large enough lie in a subspace of dimension 3, see [14, Th. 1.3].

4. RESULTANT AND FIRST ESTIMATES

The following useful result can easily be derived from the proof of [10, §5] (see also of [3, Lem. 1]). We recall the arguments since they illustrate (in a simpler situation) how we will deal with generalized determinants.

LEMMA 4.1. — *Let p, q be positive integers with $p, q \leq n$. There exists a constant $c > 0$ depending on ξ and n only, with the following property. For any polynomials $P, Q \in \mathbb{Z}[X]$ of degree p and q respectively, we have*

$$|\operatorname{Res}(P, Q)| \leq c \|P\|^{q-1} \|Q\|^{p-1} \max \{ \|P\| |Q(\xi)|, \|Q\| |P(\xi)| \}.$$

Proof. — Let $a_i, b_j \in \mathbb{Z}$ such that $P(X) = \sum_{k=0}^p a_k X^k$ and $Q(X) = \sum_{k=0}^q b_k X^k$. For $i = 1, \dots, p+q-1$, we add to the last row of the determinant (2.2) its i -th row multiplied by ξ^{p+q-i} . This last row now becomes

$$(\xi^{q-1} P(\xi), \dots, \xi P(\xi), P(\xi), \xi^{p-1} Q(\xi), \dots, \xi Q(\xi), Q(\xi)).$$

Using the upper bounds $|a_i| \leq \|P\|$ and $|b_j| \leq \|Q\|$ for the other entries of (2.2), we obtain

$$|\operatorname{Res}(P, Q)| \ll \|P\|^{q-1} |P(\xi)| \|Q\|^p + \|P\|^q \|Q\|^{p-1} |Q(\xi)|,$$

where the implicit constant only depends on p, q and ξ . $\square$

The next result, which is also based on inequalities involving resultants, will be used in Section 10. It ensures that if $R \in \mathbb{Z}[X]$ is a “good” approximation, in the sense that $R(\xi)$ is very small compared to $\|R\|$, and if we write R as a product of coprime polynomials $B_1 \cdots B_k$, then one of those factors is also a “good” approximation, while the product of the others is not.

LEMMA 4.2. — *Let m, k be positive integers. There exists a constant $c > 0$ depending on m and ξ only, with the following property. Let $B_1, \dots, B_k \in \mathbb{Z}[X]$ be non-constant, pairwise coprime polynomials, and suppose that $R := B_1 \cdots B_k$ has degree at most m . Then, there exists $j \in \{1, \dots, k\}$ such that*

$$|B_j(\xi)| \leq c \|R\|^{m-1} |R(\xi)| \quad \text{and} \quad \prod_{\substack{i=1 \\ i \neq j}}^k |B_i(\xi)| \geq c^{-1} \|R\|^{-(m-1)}.$$

Proof. — If $k = 1$ this is trivial. We now suppose that $k \geq 2$ and write $d_i = \deg(B_i)$ for $i = 1, \dots, k$. By hypothesis, we have $\deg(R) = d_1 + \dots + d_k \leq m$. Note that

$$(4.1) \quad |R(\xi)| = \prod_{i=1}^k |B_i(\xi)| \quad \text{and} \quad \|R\| \asymp \prod_{i=1}^k \|B_i\|,$$

the second inequality coming from Gelfond's lemma (with an implicit constant depending only on m). Choose $j \in \{1, \dots, k\}$ such that $|B_j(\xi)|$ is minimal and fix $i \in \{1, \dots, k\}$ with $i \neq j$. Since B_i and B_j are coprime, their resultant $\text{Res}(B_i, B_j)$ is a non-zero integer. Using Lemma 4.1, we find

$$\begin{aligned} 1 \leq |\text{Res}(B_i, B_j)| &\ll \|B_i\|^{d_j-1} \|B_j\|^{d_i-1} (\|B_j\| |B_i(\xi)| + \|B_i\| |B_j(\xi)|) \\ &\ll \|B_i\|^{d_j} \|B_j\|^{d_i} |B_i(\xi)|, \end{aligned}$$

with an implicit constant depending only on ξ and m , hence

$$-\log |B_i(\xi)| \leq d_j \log \|B_i\| + d_i \log \|B_j\| + \mathcal{O}(1).$$

On the other hand, by summing the above inequalities for $i \neq j$, and by using (4.1), we obtain

$$\begin{aligned} \sum_{\substack{i=1 \\ i \neq j}}^k -\log |B_i(\xi)| &\leq d_j \sum_{\substack{i=1 \\ i \neq j}}^k \log \|B_i\| + (m - d_j) \log \|B_j\| + \mathcal{O}(1) \\ &\leq (m - 1) \log \|R\| + \mathcal{O}(1). \end{aligned}$$

We easily deduce that

$$\prod_{\substack{i=1 \\ i \neq j}}^k |B_i(\xi)| \gg \|R\|^{-(m-1)} \quad \text{and} \quad |R(\xi)| = \prod_{i=1}^k |B_i(\xi)| \gg |B_j(\xi)| \|R\|^{-(m-1)}. \quad \square$$

5. A SEQUENCE OF IRREDUCIBLE POLYNOMIALS

As explained in the introduction, to get the upper bound $\widehat{\omega}_n(\xi) \leq 2n - 1$, the strategy of Davenport and Schmidt [10] consists in considering the resultant $\text{Res}(P, Q)$ of two “good” polynomial approximations $P, Q \in \mathbb{Z}[X]_{\leq n}$. To ensure that $\text{Res}(P, Q)$ does not vanish, they need a polynomial P which is irreducible (for it is then easy to find Q so that P and Q are coprime). The same difficulty appears in [8]. Similarly, we will not work directly with a sequence of minimal polynomials. Instead, we will consider the largest irreducible factors of the minimal polynomials. Now, let n, d be integers with

$$2 \leq d < 1 + \frac{n}{2}.$$

In this section, we assume that the transcendental real number ξ satisfies $\widehat{\omega}_n(\xi) > 2n - d$ and we fix a real number $\widehat{\omega}$ (arbitrarily close to $\widehat{\omega}_n(\xi)$) such that

$$(5.1) \quad \widehat{\omega}_n(\xi) > \widehat{\omega} > 2n - d.$$

We denote by $(P_i)_{i \geq 0}$ a sequence of minimal polynomials associated to n and ξ . Our goal is to prove the existence of a sequence $(Q_i)_{i \geq 0}$ as below.

PROPOSITION 5.1. — *Suppose that (5.1) holds. Then, there exist a sequence $(Q_i)_{i \geq 0}$ of pairwise distinct polynomials in $\mathbb{Z}[X]_{\leq n}$ and an index $j_0 \geq 0$ with the following properties. The sequence $(\|Q_i\|)_{i \geq 0}$ is bounded below by 2, unbounded and non-decreasing, and for any $i \geq 0$*

- (i) Q_i is irreducible (over $\mathbb{Z}$) and has degree at least $n - d + 2$;
- (ii) Q_i divides P_j for some index $j \geq j_0$ (not necessarily unique), and for each $j \geq j_0$ there exists $k \geq 0$ such that Q_k divides P_j ;
- (iii) $|Q_i(\xi)| = \|Q_i\|^{-\omega(Q_i)} \leq \|Q_i\|^{-\widehat{\omega}}$, and we further have

$$(5.2) \quad \omega_n(\xi) = \limsup_{k \rightarrow \infty} \omega(Q_k) \quad \text{and} \quad \liminf_{k \rightarrow \infty} \omega(Q_k) \geq \widehat{\omega}_n(\xi).$$

- (iv) if Q_i divides a minimal polynomial P_j with $j \geq j_0$, then

$$(5.3) \quad \|P_j\| \leq \|Q_i\|^{1+\theta_i}, \quad \text{where } \theta_i = \frac{\omega(Q_i) - 2n + d}{n - 2d + 3};$$

- (v) we have

$$(5.4) \quad \|Q_{i+1}\|^\tau \leq \|Q_i\| \quad \text{where } \tau = \frac{\widehat{\omega}(\widehat{\omega} - n - d + 3)}{\omega_n(\xi)(\omega_n(\xi) - n - d + 3)},$$

with the convention $\tau = 0$ if $\omega_n(\xi) = \infty$.

The above proposition is essentially a consequence of Lemma 5.3 below. Assertion (iii) ensures that the polynomials Q_i are quite good approximations, and they can be used to compute the exponent of best approximation $\omega_n(\xi)$. Estimate (5.4) is the analog of the second inequality of (3.2) but is way more difficult to prove. The main reason behind this difficulty is that there may be many polynomials $P \in \mathbb{Z}[X]_{\leq n}$ with $\|Q_i\| < \|P\| < \|Q_{i+1}\|$ and $|P(\xi)| < |Q_i(\xi)|$.

In order to prove Proposition 5.1, we need the two technical lemmas below. Essentially, they will be used to prove that the factors of P_i of small degree are bad approximations. This will lead to the existence of a factor of large degree which is necessarily a rather good approximation.

LEMMA 5.2. — Suppose that (5.1) holds. Then, there exists a constant $c \in (0, 1)$ depending only on ξ and n such that for any non-zero polynomial $R \in \mathbb{Z}[X]_{\leq n-d+1}$ we have

$$(5.5) \quad |R(\xi)| \geq c \|R\|^{-(n+\deg(R)-1)} \geq c \|R\|^{-(2n-d)}.$$

In particular (5.5) holds for any $R \in \mathbb{Z}[X]_{\leq d-2}$.

Proof. — If R is constant we have $|R(\xi)| = \|R\|$ and the result is trivial. Now, suppose that R is irreducible and not constant. We adapt the arguments of Davenport and Schmidt [10, §5–6]. Set $H = e^{-n} \|R\|$. By definition of $\widehat{\omega}_n(\xi)$ and $\widehat{\omega}$, if H is sufficiently large, there exists a non-zero $P \in \mathbb{Z}[X]_{\leq n}$ such that

$$\|P\| \leq H \quad \text{and} \quad |P(\xi)| \leq H^{-\widehat{\omega}}.$$

By (2.1), the (irreducible) polynomial R is not a factor of P , they are thus coprime polynomials. Their resultant is a non-zero integer, and using Lemma 4.1, we obtain

$$\begin{aligned} 1 &\ll \|P\|^{\deg(R)-1} \|R\|^n |P(\xi)| + \|P\|^{\deg(R)} \|R\|^{n-1} |R(\xi)| \\ &\ll H^{n+\deg(R)-1-\widehat{\omega}} + H^{n+\deg(R)-1} |R(\xi)|. \end{aligned}$$

Since $\widehat{\omega} > 2n - d$ and $\deg(R) \leq n - d + 1$, the first term tends to 0 as H tends to infinity. Hence $1 \ll H^{n+\deg(R)-1} |R(\xi)|$, which implies (5.5).

If R is not irreducible, we write $R = \prod_{i=1}^s R_i$ with integer $s \geq 1$ and $R_1, \dots, R_s \in \mathbb{Z}[X]$ irreducible of degree $\leq \deg(R)$ (possibly constant). Combining $\|R\| \asymp \prod_{i=1}^s \|R_i\|$ together with (5.5) applied to the irreducible factors R_i , we find

$$|R(\xi)| = \prod_{i=1}^s |R_i(\xi)| \gg \prod_{i=1}^s \|R_i\|^{-(n+\deg(R)-1)} \gg \|R\|^{-(n+\deg(R)-1)}.$$

Finally, the last assertion comes from the fact that $d - 1 \leq n + d - 1$ (since $d \leq 1 + n/2$). $\square$

LEMMA 5.3. — *Suppose that (5.1) holds. There exist $i_0 \geq 0$ and a constant $c > 0$ such that for each $i \geq i_0$ the polynomial P_i has a unique irreducible factor $\tilde{P}_i \in \mathbb{Z}[X]$ of degree $\geq n - d + 2$ and positive leading coefficient. It satisfies*

$$(5.6) \quad |P_i(\xi)| \|P_i\|^{n+d-3} \geq c |\tilde{P}_i(\xi)| \|\tilde{P}_i\|^{n+d-3},$$

moreover $(\|\tilde{P}_i\|)_{i \geq i_0}$ tends to infinity and as i tends to infinity. For each i large enough, we have $\|\tilde{P}_i\| > 1$, and writing $|\tilde{P}_i(\xi)| = \|\tilde{P}_i\|^{-\omega(\tilde{P}_i)}$, we furthermore have

$$(5.7) \quad \omega_n(\xi) = \limsup_{i \rightarrow \infty} \omega(\tilde{P}_i) \quad \text{and} \quad \liminf_{i \rightarrow \infty} \omega(\tilde{P}_i) \geq \widehat{\omega}_n(\xi).$$

Proof. — First, note that since $d < 1 + n/2$, if we decompose P_i as a product of irreducibles, there is at most one factor of degree $\geq n - d + 2$. Fix $i \geq 0$ large enough so that $\omega(P_i) \geq \widehat{\omega}$, and write

$$P := P_i = \prod_{k=1}^s R_k,$$

where $R_1, \dots, R_s \in \mathbb{Z}[X]$ are irreducible polynomials (and s is a positive integer). Suppose that $\deg(R_k) \leq n - d + 1$ for each $k = 1, \dots, s$. Then, by Lemma 5.2 together with $\|P\| \asymp \prod_k \|R_k\|$, we find

$$\|P\|^{-\widehat{\omega}} \geq |P(\xi)| = \prod_{k=1}^s |R_k(\xi)| \gg \prod_{k=1}^s \|R_k\|^{-(2n-d)} \asymp \|P\|^{-(2n-d)}.$$

This is impossible if i is sufficiently large since $\widehat{\omega} > 2n - d$. Therefore, if i is large enough, one of the factors R_k has degree at least $n - d + 2$. Without loss of generality, we may suppose that it is $R := R_1$. Write $S := \prod_{k=2}^s R_k$, so that $P = RS$. We have $\deg(S) \leq d - 2$, and (5.5) of Lemma 5.2 yields

$$|S(\xi)| \gg \|S\|^{-(n+d-3)}.$$

Together with $\|P\| \asymp \|R\| \|S\|$, this leads to

$$|P(\xi)| = |R(\xi)| |S(\xi)| \gg |R(\xi)| \|S\|^{-(n+d-3)} \asymp |R(\xi)| \|R\|^{n+d-3} \|P\|^{-(n+d-3)},$$

and (5.6) follows easily by setting $\tilde{P}_i := R$. The rest of the proof is based only on (5.6) and the inequality $\|\tilde{P}_i\| \ll \|P_i\|$. Note that $|P_i(\xi)| \|P_i\|^{n+d-3} \ll \|P_i\|^{n+d-3-\widehat{\omega}}$ tends to 0 as i tends to infinity (using $d < 1 + n/2$ together with $\widehat{\omega} > 2n - d$). We deduce that

$|\tilde{P}_i(\xi)|\|\tilde{P}_i\|^{n+d-3}$ also tends to 0 as i tends to infinity, which is possible only if $\|\tilde{P}_i\|$ tends to infinity. In particular, if i is large enough we must have $\|\tilde{P}_i\| > 1$. Writing $|\tilde{P}_i(\xi)| = \|\tilde{P}_i\|^{-\omega(\tilde{P}_i)}$, we also have $\omega(\tilde{P}_i) > n + d - 3$. Now, using $\|\tilde{P}_i\| \ll \|P_i\|$, and taking the logarithms of the two sides of (5.6), we get

$$\begin{aligned} (\omega(P_i) - (n + d - 3)) \log \|P_i\| &\leq (\omega(\tilde{P}_i) - (n + d - 3)) \log \|\tilde{P}_i\| + \mathcal{O}(1) \\ &\leq (\omega(\tilde{P}_i) - (n + d - 3)) (\log \|P_i\| + \mathcal{O}(1)) + \mathcal{O}(1). \end{aligned}$$

By dividing by $\log \|P_i\|$ and by simplifying, we deduce that $\omega(\tilde{P}_i) \geq \omega(P_i)(1 - o(1))$ and (5.7) follows easily from (3.3). $\square$

Proof of Proposition 5.1. — Let $i_0 \geq 0$ and $(\tilde{P}_i)_{i \geq i_0}$ given by Lemma 5.3. Let $(Q_i)_{i \geq 0}$ be the (infinite) sequence of factors $(\tilde{P}_j)_{j \geq i_0}$ reordered by increasing height, without repetition. By Lemma 5.3, we may assume i_0 large enough so that $\|Q_i\| > 1$ for each i , as well as $|Q_i(\xi)| \leq \|Q_i\|^{-\hat{\omega}}$. This sequence clearly satisfies the first assertions (i) to (iii), the third one coming from (5.7) together with (3.3).

Now, let $i \geq 0$ and let $j \geq i_0$ be an index such that Q_i divides P_j . Since we have $\|Q_i\| \ll \|P_j\|$ by Gelfond's lemma, the index j tends to infinity as i tends to infinity. Then, estimate (5.6) can be rewritten as

$$(5.8) \quad |P_j(\xi)|^{-1} \|P_j\|^{-n-d+3} \ll |Q_i(\xi)|^{-1} \|Q_i\|^{-n-d+3} = \|Q_i\|^{\omega(Q_i)-n-d+3}.$$

Using $|P_j(\xi)|^{-1} \gg \|P_j\|^{\hat{\omega}}$ and $\hat{\omega} > 2n - d$, we get, for each large enough i ,

$$\|P_j\|^{n-2d+3} \leq \|Q_i\|^{\omega(Q_i)-n-d+3},$$

which is equivalent to (5.3). So, assertion (iv) holds assuming i_0 large enough.

It remains to prove assertion (v). Note that this is trivial if $\omega_n(\xi) = \infty$. Let us assume that $\omega_n(\xi) < \infty$ and fix a small $\varepsilon > 0$ to be chosen later. For each pair (i, j) as above with $j \geq i_0$ large enough as a function of ε , we have $\omega(P_j) > \hat{\omega}_n(\xi) - \varepsilon/2$ and $\omega(Q_i) < \omega_n(\xi) + \varepsilon/2$, and thus (5.8) yields

$$\|P_j\|^{\hat{\omega}_n(\xi) - \varepsilon - n - d + 3} \leq \|Q_i\|^{\omega_n(\xi) + \varepsilon - n - d + 3},$$

for each $i \geq 0$ and each $j \geq i_0$ such that Q_i divides P_j . We define k as the largest index such that

$$\|P_k\| \leq \|Q_i\|^{\theta(\varepsilon)}, \quad \text{where } \theta(\varepsilon) = \frac{\omega_n(\xi) + \varepsilon - n - d + 3}{\hat{\omega}_n(\xi) - \varepsilon - n - d + 3}.$$

Since $\|P_j\| \leq \|Q_i\|^{\theta(\varepsilon)}$, by maximality of k we have $i_0 \leq j \leq k$. Let ℓ be such that Q_ℓ divides P_{k+1} . We find

$$\|P_k\| \leq \|Q_i\|^{\theta(\varepsilon)} < \|P_{k+1}\| \leq \|Q_\ell\|^{\theta(\varepsilon)},$$

and therefore $\ell \geq i + 1$. On the other hand, since by Gelfond's lemma we have $\|Q_\ell\| \ll \|P_{k+1}\|$, we deduce from (3.2) that

$$\|Q_{i+1}\| \leq \|Q_\ell\| \ll \|P_{k+1}\| \ll \|P_k\|^{\omega_n(\xi)/\hat{\omega}} \leq \|Q_i\|^{\omega_n(\xi)\theta(\varepsilon)/\hat{\omega}}.$$

We now choose $\varepsilon > 0$ small enough so that

$$\theta(\varepsilon) < \frac{\omega_n(\xi) - n - d + 3}{\widehat{\omega} - n - d + 3}.$$

This is possible since $\widehat{\omega} < \widehat{\omega}_n(\xi)$, and it yields (5.4) for each $i \geq 0$, assuming that i_0 is large enough. $\square$

6. ON THE DIMENSION OF SOME POLYNOMIAL SUBSPACES

We start by introducing some families of vector spaces spanned by polynomials, and we study their dimensions.

DEFINITION 6.1. — Let $k \geq n$ be an integer and let $\mathcal{A}$ be a subset of $\mathbb{R}[X]_{\leq n}$. We define

$$\begin{aligned} \mathcal{B}_k(\mathcal{A}) &= \{Q, XQ, \dots, X^{k-\deg(Q)}Q; Q \in \mathcal{A} \setminus \{0\}\} \subseteq \mathbb{R}[X]_{\leq k}, \\ V_k(\mathcal{A}) &= \langle \mathcal{B}_k(\mathcal{A}) \rangle_{\mathbb{R}}, \\ g_{\mathcal{A}}(k) &= \dim V_k(\mathcal{A}). \end{aligned}$$

The spaces $V_k(\mathcal{A})$ play the role of the spaces $\mathcal{U}^k(\mathcal{A})$ in [17, §3] (for simultaneous approximation). We obtain analog properties. Note that if $\mathcal{A}$ contains at least one non-zero polynomial, then

$$(6.1) \quad V_n(\mathcal{A}) \subsetneq V_{n+1}(\mathcal{A}) \subsetneq \dots$$

The goal of this section is to prove the following result. We could not find a reference for the proposition below.

PROPOSITION 6.2. — Let k be an integer with $0 \leq k \leq n$, and let $\mathcal{A}$ be a set of $k+1$ linearly polynomials of $\mathbb{R}[X]_{\leq n}$. Suppose that the gcd of the elements of $\mathcal{A}$ is 1 (in other words, the ideal spanned by $\mathcal{A}$ is $\mathbb{R}[X]$). Then

$$(6.2) \quad V_{2n-k}(\mathcal{A}) = \mathbb{R}[X]_{\leq 2n-k}.$$

The case $k = 1$ is a classical result (it is implied by the fact that the resultant of two coprime polynomials is non-zero). The proof of Proposition 6.2 is given at the end of the section. Recall that a function $f : \{n, n+1, \dots\} \rightarrow \mathbb{R}$ is *concave* if for any $i > n$, it satisfies

$$f(i) - f(i-1) \geq f(i+1) - f(i).$$

The next result is a dual version of [17, Prop. 3.1] (where we deal with simultaneous approximation to the successive powers of ξ).

LEMMA 6.3. — Let $\mathcal{A} \neq \{0\}$ be a non-empty subset of $\mathbb{R}[X]_{\leq n}$. The function $g_{\mathcal{A}}$ is concave and (strictly) increasing on $\{n, n+1, \dots\}$.

Proof. — The series of inclusions (6.1) shows that the function $g_{\mathcal{A}}$ is increasing on $\{n, n+1, \dots\}$. For simplicity, we write $V_i = V_i(\mathcal{A})$ and $\mathcal{B}_i = \mathcal{B}_i(\mathcal{A})$ for each $i \geq n$. Given an integer $i \geq n$ we have $XV_i \subseteq V_{i+1}$, and we set

$$h(i) := \dim(V_{i+1}/XV_i) = g_{\mathcal{A}}(i+1) - g_{\mathcal{A}}(i).$$

We have to prove that h is decreasing on $\{n, n+1, \dots\}$. Fix $i \geq n+1$ and consider the linear map $\pi : V_i \rightarrow V_{i+1}/XV_i$ defined by $\pi(P) = P + XV_i$. Since $\mathcal{B}_i \cup X\mathcal{B}_i = \mathcal{B}_{i+1}$, we have $V_i + XV_i = V_{i+1}$. So π is surjective, and consequently $\text{Im } \pi = V_{i+1}/XV_i$ is isomorphic to $V_i/\ker \pi$. On the other hand, $XV_{i-1} \subseteq V_i \cap XV_i \subseteq \ker \pi$, so XV_{i-1} is subspace of $\ker \pi$. Hence

$$h(i-1) = \dim(V_i/XV_{i-1}) \geq \dim(V_i/\ker \pi) = \dim(V_{i+1}/XV_i) = h(i). \quad \square$$

LEMMA 6.4. — Let $P, Q \in \mathbb{R}[X]_{\leq n}$ be two coprime polynomials. Then, we have

$$\dim V_{n+j}(P, Q) \geq 2(j+1),$$

for each $j \in \{0, \dots, n-1\}$. In particular $V_{2n-1}(P, Q) = \mathbb{R}[X]_{\leq 2n-1}$.

Proof. — Let p (resp. q) denote the degree of P (resp. of Q). There exist $\alpha, \beta \in \mathbb{R}$ such that the polynomial $\tilde{P} := P(X)(X - \alpha)^{n-p}$ and $\tilde{Q} := Q(X)(X - \beta)^{n-q}$ are coprime (and of degree exactly n). Fix $j \in \{0, \dots, n-1\}$. Since $\tilde{P}$ and $\tilde{Q}$ are coprime and $j < n$, the linear map

$$\begin{aligned} \mathbb{R}[X]_{\leq j} \times \mathbb{R}[X]_{\leq j} &\longrightarrow \mathbb{R}[X]_{\leq n+j} \\ (R, S) &\longmapsto R\tilde{P} + S\tilde{Q} \end{aligned}$$

is injective, so its image $V_{n+j}(\tilde{P}, \tilde{Q}) \subseteq V_{n+j}(P, Q)$ has dimension $2(j+1)$. $\square$

Proof of Proposition 6.2. — For simplicity, we write $g = g_{\mathcal{A}}$. Recall that $\mathcal{A}$ has cardinality $k+1$, so that $g(n) \geq \text{card}(\mathcal{A}) = k+1$. If $k = n$, then (6.2) is automatic (since in that case $\mathcal{A}$ contains a basis of $\mathbb{R}[X]_{\leq n}$). So, we may assume that $k < n$. We first prove that for each sufficiently large m , we have

$$(6.3) \quad V_m(\mathcal{A}) = \mathbb{R}[X]_{\leq m}.$$

Indeed, since the ideal spanned by $\mathcal{A}$ is $\mathbb{R}[X]$, there exists an integer $\ell \geq n$ such that $1 \in V_{\ell}(\mathcal{A})$. Let P be a non-zero element in $\mathcal{A}$ of degree d , and set $m = \ell + d$. Then $V_m(\mathcal{A})$ contains $\mathbb{R}[X]_{\leq d}$, as well as the polynomials $P, XP, \dots, X^{\ell}P$. We easily deduce (6.3).

By contradiction, suppose that (6.2) does not hold, i.e.,

$$(6.4) \quad g(2n-k) \leq 2n-k.$$

We distinguish two cases. Suppose first that $g(2n-k) - g(2n-k-1) \geq 2$. By concavity, then $g(j) - g(j-1) \geq 2$ for each j with $n < j \leq 2n-k$, and we deduce that

$$g(2n-k) \geq g(n) + 2(n-k) \geq k+1 + 2(n-k) = 2n-k+1,$$

since $g(n) \geq \text{card}(\mathcal{A}) = k+1$. This contradicts (6.4), so $g(2n-k) - g(2n-k-1) \leq 1$. Since the function g is increasing and concave, it is linear with slope 1 on

$$\{2n-k, 2n-k+1, \dots\}.$$

Choosing $m > 2n-k$ such that (6.3) holds, we obtain by (6.4)

$$m+1 = g(m) = g(2n-k) + m - (2n-k) \leq m,$$

a contradiction. Hence (6.2) holds. $\square$




FIGURE 1. Graph of the piecewise linear function interpolating the values $g(i) = \dim V_i(\mathcal{A})$ at integers $i \in \{n, \dots, 2n-1\}$.

7. PROOF OF THEOREM 1.1 (CASE $d = 2$)

In this section, we deal with the case $d = 2$ to prove Theorem 1.1, namely that $\widehat{\omega}_3(\xi) \leq 2 + \sqrt{5} = 4.23 \dots$ and $\widehat{\omega}_n(\xi) \leq 2n - 2$ for each $n \geq 4$. The estimate $\widehat{\omega}_n(\xi) \leq 2n - 2$ was already known for $n \geq 10$, however for $n = 4, \dots, 9$ it is a new result. For $n = 3$, our bound improves on the bound $\widehat{\omega}_3(\xi) \leq 3 + \sqrt{2} = 4.41 \dots$ due to Bugeaud and Schleischitz [8]. Moreover, our proof does not require Marnat-Moshchevitin's inequality [13].

Proof of Theorem 1.1. — Suppose that $\widehat{\omega}_n(\xi) > 2n - 2$, and fix a real number $\widehat{\omega}$ such that

$$\widehat{\omega}_n(\xi) > \widehat{\omega} > 2n - 2.$$

Let $(P_i)_{i \geq 0}$ be a sequence of minimal polynomials associated to n and ξ as in Section 3. According to Lemma 5.3 (with $d = 2$) there exists an index $i_0 \geq 0$ such that P_i has degree n and is irreducible for each $i \geq i_0$. Consequently, up to a finite number of terms, the sequence $(P_i)_{i \geq 0}$ coincides with the sequence $(Q_i)_{i \geq 0}$ of Proposition 5.1. Let I denotes the set of indices $i \geq i_0 + 1$ such that P_{i-1} , P_i and P_{i+1} are linearly independent. By Lemmas 3.2 and 3.4, the set I is infinite, and for any consecutive $i < j$ in I , we have

$$\|P_{i+1}\| |P_i(\xi)| \asymp \|P_j\| |P_{j-1}(\xi)|.$$

Furthermore, the irreducible polynomials P_i and P_{i+1} are also coprime since $\|P_i\| < \|P_{i+1}\|$ and $|P_i(\xi)| > |P_{i+1}(\xi)|$. Lemma 4.1 yields

$$1 \leq |\text{Res}(P_i, P_{i+1})| \ll \|P_i\|^{n-1} \|P_{i+1}\|^n |P_i(\xi)| \ll \|P_i\|^{n-1} \|P_j\|^n |P_{j-1}(\xi)| \\ \ll \|P_i\|^{n-1} \|P_j\|^{n-\widehat{\omega}}.$$

We deduce that

$$(7.1) \quad \|P_j\| \leq \|P_i\|^\theta \quad \text{where } \theta = \frac{n-1}{\widehat{\omega}-n}.$$

Let $h < i < j$ be consecutive indices in I . We have the following configuration

$$\langle P_h, P_{h+1} \rangle_{\mathbb{R}} = \langle P_{i-1}, P_i \rangle_{\mathbb{R}} \neq \langle P_i, P_{i+1} \rangle_{\mathbb{R}},$$

so P_h, P_{h+1}, P_{i+1} are linearly independent. Proposition 6.2 combined with Lemma 6.4 implies that

$$(\mathbb{R}[X]_{\leq n-2} P_h \oplus \mathbb{R}[X]_{\leq n-2} P_{h+1}) + \mathbb{R}[X]_{\leq n-2} P_{i+1} = \mathbb{R}[X]_{\leq 2n-2}.$$

Choose $k \in \{0, \dots, n-2\}$ such that $(P_h, \dots, X^{n-2} P_h, P_{h+1}, \dots, X^{n-2} P_{h+1}, X^k P_{i+1})$ is a basis of $\mathbb{R}[X]_{\leq 2n-2}$. We denote by M the matrix of this basis expressed in the canonical basis $(1, X, \dots, X^{2n-2})$. Estimating $\det(M)$ as in the proof of Lemma 4.1 (in other words, for $\ell = 2, \dots, 2n-2$, we add to the first row of M the ℓ -th row multiplied by $\xi^{\ell-1}$), we get the estimates

$$1 \leq |\det(M)| \ll |P_h(\xi)| \|P_h\|^{n-2} \|P_{h+1}\|^{n-1} \|P_{i+1}\|.$$

Now, since $\|P_{h+1}\|^{n-1} |P_h(\xi)| \asymp \|P_{h+1}\|^{n-2} \|P_i\| |P_{i-1}(\xi)| \ll \|P_i\|^{n-1-\widehat{\omega}}$, we deduce that

$$(7.2) \quad \|P_i\|^{\widehat{\omega}-n+1} \ll \|P_h\|^{n-2} \|P_j\|.$$

For consecutive $i < j$ in I , define $\tau_i \in (0, 1)$ by

$$\|P_i\| = \|P_j\|^{\tau_i}$$

and set $\tau = \limsup_{i \in I, i \rightarrow \infty} \tau_i \in [0, 1]$. Let $h < i < j$ be consecutive indices in I as previously. By (7.2), we obtain

$$\widehat{\omega} - n + 1 \leq (n-2)\tau_h + \frac{1}{\tau_i} + o(1) \leq (n-2)\tau + \frac{1}{\tau_i} + o(1).$$

We infer that

$$(7.3) \quad p(\tau) \geq 0, \quad \text{where } p(t) = (n-2)t^2 - (\widehat{\omega} - n + 1)t + 1.$$

Note that

$$p(0) = 1, \quad p\left(\frac{1}{n-2}\right) = \frac{2n-2-\widehat{\omega}}{n-2} < 0 \quad \text{and} \quad p(1) = 2n-2-\widehat{\omega} < 0.$$

We deduce that p has one root $\alpha \in (0, 1/(n-2))$ and one root larger than 1. Since $\tau \in [0, 1]$ and $p(\tau) \geq 0$, we obtain $\tau \leq \alpha$. Combined with the estimate $\|P_i\| = \|P_j\|^{\tau_i} \ll \|P_i\|^{\theta\tau_i}$ valid for any $i \in I$ (this is a consequence of (7.1)), this leads to

$$(7.4) \quad 1 \leq \theta\tau \leq \theta\alpha < \frac{n-1}{(n-2)^2}.$$

We easily check that this is impossible when $n \geq 4$ (the right-hand side is strictly less than 1), thus $\widehat{\omega}_n(\xi) \leq 2n - 2$ for each $n \geq 4$.

We now deal with the case $n = 3$. Suppose by contradiction that $\widehat{\omega}_3(\xi) > 2 + \sqrt{5}$ and choose $\widehat{\omega}$ such that

$$\widehat{\omega}_3(\xi) > \widehat{\omega} > 2 + \sqrt{5}.$$

The polynomial p from (7.3) becomes $p(t) = t^2 - (\widehat{\omega} - 2)t + 1$. Denote by α its smallest root, and by $\beta = (\sqrt{5} - 1)/2$ the smallest root of the polynomial $t^2 - \sqrt{5}t + 1$. We find

$$0 = \beta^2 - \sqrt{5}\beta + 1 > \beta^2 - (\widehat{\omega} - 2)\beta + 1 = p(\beta),$$

hence $\alpha < \beta$. Combined with $\theta = 2/(\widehat{\omega} - 3) < 1/\beta$, this implies that $\theta\alpha < 1$, which contradicts (7.4). It follows that $\widehat{\omega}_3(\xi) \leq 2 + \sqrt{5}$. $\square$

8. MULTILINEAR ALGEBRA AND HEIGHT OF POLYNOMIAL SUBSPACES

This section is divided into two parts. We introduce and study a quantity $\mathcal{D}_\xi(V)$ associated to a subspace $V \subseteq \mathbb{R}^m$ defined over $\mathbb{Q}$ in Section 8.2. Intuitively, $\mathcal{D}_\xi(V)$ is small if V is spanned by good polynomial approximations of $\mathbb{Z}[X]$ (i.e., small when evaluated at ξ). This will be a key-point for estimating the height of the polynomials Q_i of Section 5. In order to define $\mathcal{D}_\xi$, we need some tools of multilinear algebra that we recall in Section 8.1. In the appendix we give another interpretation of $\mathcal{D}_\xi$ in term of twisted heights.

8.1. MULTILINEAR ALGEBRA AND HODGE DUALITY. — For each integer m , we view $\mathbb{R}^{m+1}$ as an Euclidean space for the usual scalar product $(\cdot | \cdot)$, and we denote by $\|\cdot\|_2$ the associated Euclidean norm. For each $k = 1, \dots, m+1$, we identify $\Lambda^k \mathbb{R}^{m+1}$ with $\mathbb{R}^N$, where $N = \binom{m+1}{k}$, via an ordering of the Plücker coordinates, and we denote by $\|\mathbf{y}\|_2$ the norm of a point $\mathbf{y} \in \Lambda^k \mathbb{R}^{m+1} \cong \mathbb{R}^N$. This is independent of the ordering. Let V be a k -dimensional subspace of $\mathbb{R}^{m+1}$ defined over $\mathbb{Q}$, i.e., such that $\langle V \cap \mathbb{Q}^{m+1} \rangle_{\mathbb{R}} = V$. Its (standard) height $\mathcal{H}(V)$ is the covolume of the lattice $V \cap \mathbb{Z}^{m+1}$ inside V (with the convention that $\mathcal{H}(V) = 1$ if $V = \{0\}$). Explicitly, we have

$$\mathcal{H}(V) := \|\mathbf{x}_1 \wedge \dots \wedge \mathbf{x}_k\|_2,$$

for any $\mathbb{Z}$ -basis $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ of the lattice $V \cap \mathbb{Z}^{m+1}$. Schmidt established the very nice inequality

$$\mathcal{H}(U \cap V) \mathcal{H}(U + V) \leq \mathcal{H}(U) \mathcal{H}(V),$$

valid for any subspaces U, V of $\mathbb{R}^{m+1}$ defined over $\mathbb{Q}$ (see [27, Ch. I, Lem. 8A]). In this paper, we need to work with a “twisted” height and the corresponding version of Schmidt’s inequality (obtained by following Schmidt’s original arguments).

Let $(\mathbf{e}_1, \dots, \mathbf{e}_{m+1})$ denotes the canonical basis of $\mathbb{R}^{m+1}$, and let k be an integer with $0 \leq k \leq m+1$. The Hodge star operator

$$* : \Lambda^k \mathbb{R}^{m+1} \xrightarrow{\sim} \Lambda^{m+1-k} \mathbb{R}^{m+1}$$

is defined by

$$*(\mathbf{e}_{i_1} \wedge \dots \wedge \mathbf{e}_{i_k}) = \varepsilon_{i_1, \dots, i_k} \mathbf{e}_{j_1} \wedge \dots \wedge \mathbf{e}_{j_{m+1-k}}$$

and we deduce that the I -coordinate of $\mathbf{Y}$ is equal to 0, which proves assertion (ii). Suppose now that $i_1 = 1$ and set $J = (1, i_2 + 1, \dots, i_k + 1)$. Then

$$D_I(\mathbf{y}_1, \dots, \mathbf{y}_k) = T^{m+1-k} \mathfrak{D}_J(\mathbf{z}_1, \dots, \mathbf{z}_k),$$

hence the first part of (iii). The second part is obtained by combining the above with assertion (i).

We deduce from the last two assertions that all the non-zero coordinates of $\mathbf{Y}$ are coordinates of $\mathbf{Z}$, thus $\|\mathbf{Y}\|_2 \leq \|\mathbf{Z}\|_2$, which proves the second inequality in (A.5). For the first estimate, we need to estimate the K -coordinates of $\mathbf{Z}$ with $K \in \mathcal{J}(k-1, m+1)$ of the form $(1, i_2, \dots, i_{k-1})$. According to assertion (i), they are exactly the determinants $\mathfrak{D}_J(\mathbf{z}_1, \dots, \mathbf{z}_k)$ with $J = (1, 2, j_3, \dots, j_k)$ in $\mathcal{J}(k, m+2)$.

Fix a $J \in \mathcal{J}(k, m+2)$ as above. The second row of the matrix $M(\mathbf{z}_1, \dots, \mathbf{z}_k)$ is a linear combination of the remaining rows (with coefficients in absolute value between 1 and $|\xi|^m$). We deduce that $\mathfrak{D}_J(\mathbf{z}_1, \dots, \mathbf{z}_k)$ can be written as a linear combination of $\mathfrak{D}_{J'}(\mathbf{z}_1, \dots, \mathbf{z}_k)$, where J' belong to the subset of $\mathcal{J}(k, m+2)$ consisting in the k -tuples whose second element is ≥ 3 . By assertion (iii), they are all coordinates of $\mathbf{Y}$, hence $|\mathfrak{D}_J(\mathbf{z}_1, \dots, \mathbf{z}_k)| \ll \|\mathbf{Y}\|_2$. We conclude that $\|\mathbf{Z}\|_2 \ll \|\mathbf{Y}\|_2$.

Finally, fix $(i_2, \dots, i_k) \in \mathcal{J}(k-1, m)$ and set $K = (i_2 + 1, \dots, i_k + 1)$. By definition of $\mathbf{Z}$, the K -coordinate of $\mathbf{Z}$ is equal to

$$\sum_{i=1}^k p_i \mathfrak{D}_I(\mathbf{x}_1, \dots, \widehat{\mathbf{x}}_i, \dots, \mathbf{x}_k) = \sum_{i=1}^k p_i \mathfrak{D}_J(\mathbf{x}_1^+, \dots, \widehat{\mathbf{x}}_i^+, \dots, \mathbf{x}_k^+).$$

By assertion (iii), this is also the $(1, K)$ -coordinate of $\mathbf{Y}$. So, the set of non-zero coordinates of $\mathbf{Y}$ is exactly equal to the set of non-zero coordinates of the point

$$\sum_{i=1}^k p_i \mathbf{x}_1^+ \wedge \dots \wedge \widehat{\mathbf{x}}_i^+ \wedge \dots \wedge \mathbf{x}_k^+.$$

Equation (A.6) follows from the second identity of (A.7). $\square$

REFERENCES

- [1] B. ARBOUR & D. ROY – “A Gel’fond type criterion in degree two”, *Acta Arith.* **111** (2004), no. 1, p. 97–103.
- [2] N. BOURBAKI – *Algebra I*, Springer-Verlag, New York, 1989.
- [3] W. D. BROWNAWELL – “Sequences of Diophantine approximations”, *J. Number Theory* **6** (1974), no. 1, p. 11–21.
- [4] Y. BUGEAUD – *Approximation by algebraic numbers*, Cambridge Tracts in Math., vol. 160, Cambridge University Press, Cambridge, 2004.
- [5] ———, “Exponents of Diophantine approximation”, in *Dynamics and analytic number theory* (D. Badziahin, A. Gorodnik & N. Peyerimhoff, eds.), London Math. Soc. Lect. Note Ser., vol. 437, Cambridge University Press, Cambridge, 2016, p. 96–135.
- [6] Y. BUGEAUD & M. LAURENT – “Exponents of Diophantine approximation and Sturmian continued fractions”, *Ann. Inst. Fourier (Grenoble)* **55** (2005), no. 3, p. 773–804.
- [7] ———, “On transfer inequalities in Diophantine approximation, II”, *Math. Z.* **265** (2010), no. 2, p. 249–262.
- [8] Y. BUGEAUD & J. SCHLEISCHITZ – “On uniform approximation to real numbers”, *Acta Arith.* **175** (2016), no. 3, p. 255–268.

- [9] H. DAVENPORT & W. SCHMIDT – “Approximation to real numbers by quadratic irrationals”, *Acta Arith.* **13** (1967), no. 2, p. 169–176.
- [10] H. DAVENPORT & W. SCHMIDT – “Approximation to real numbers by algebraic integers”, *Acta Arith.* **15** (1969), no. 4, p. 393–416.
- [11] W. HODGE & D. PEDOE – *Methods of algebraic geometry*, Cambridge University Press, Cambridge, 1947.
- [12] M. LAURENT – “Simultaneous rational approximation to the successive powers of a real number”, *Indag. Math. (N.S.)* **14** (2003), no. 1, p. 45–53.
- [13] A. MARNAT & N. MOSHCHEVITIN – “An optimal bound for the ratio between ordinary and uniform exponents of Diophantine approximation”, *Mathematika* **66** (2020), no. 3, p. 818–854.
- [14] N. G. MOSHCHEVITIN – “Best Diophantine approximations: the phenomenon of degenerate dimension”, in *Surveys in geometry and number theory: reports on contemporary Russian mathematics*, London Math. Soc. Lecture Note Ser., vol. 338, Cambridge University Press, Cambridge, 2007, p. 158–182.
- [15] A. POËLS – “Exponents of Diophantine approximation in dimension 2 for numbers of Sturmian type”, *Math. Z.* **294** (2020), no. 3, p. 951–993.
- [16] ———, “Exponents of Diophantine approximation in dimension two for a general class of numbers”, *Mosc. J. Comb. Number Theory* **11** (2022), no. 1, p. 37–69.
- [17] A. POËLS & D. ROY – “Simultaneous rational approximation to successive powers of a real number”, *Trans. Amer. Math. Soc.* **375** (2022), no. 09, p. 6385–6415.
- [18] ———, “Parametric geometry of numbers over a number field and extension of scalars”, *Bull. Soc. math. France* **151** (2023), p. 257–303.
- [19] M. RIVARD-COOKE – “Parametric geometry of numbers”, PhD Thesis, University of Ottawa, 2019, <https://ruor.uottawa.ca/handle/10393/38871>.
- [20] D. ROY – “Approximation simultanée d’un nombre et de son carré”, *Comptes Rendus Mathématique* **336** (2003), no. 1, p. 1–6.
- [21] ———, “Approximation to real numbers by cubic algebraic integers I”, *Proc. London Math. Soc. (3)* **88** (2004), no. 1, p. 42–62.
- [22] ———, “On two exponents of approximation related to a real number and its square”, *Canad. J. Math.* **59** (2007), no. 1, p. 211–224.
- [23] J. SCHLEISCHITZ – “Some notes on the regular graph defined by Schmidt and Summerer and uniform approximation”, *JP J. Algebra Number Theory Appl.* **39** (2017), no. 2, p. 115–150.
- [24] ———, “Uniform Diophantine approximation and best approximation polynomials”, *Acta Arith.* **185** (2018), no. 3, p. 249–274.
- [25] ———, “Uniform dual approximation to Veronese curves in small dimension”, 2024, [arXiv: 2405.10086](https://arxiv.org/abs/2405.10086).
- [26] W. M. SCHMIDT – *Diophantine approximation*, Lect. Notes in Math., vol. 785, Springer-Verlag, Berlin, 1980.
- [27] ———, *Diophantine approximations and Diophantine equations*, Lect. Notes in Math., vol. 1467, Springer-Verlag, Berlin, 1991.

Manuscript received 16th May 2024
accepted 18th July 2024

ANTHONY POËLS, Université Claude Bernard Lyon 1, Institut Camille Jordan UMR 5208,
69622 Villeurbanne, France
E-mail : poels@math.univ-lyon1.fr
Url : <http://apoels-math-u.net/>