THE CODIMENSION ZERO POINTS OF AN ALGEBRAIC SPACE ARE DENSE IN ITS TOPOLOGICAL SPACE

THE STACKS PROJECT AUTHORS

Editorial note. All delivered mathematical text is the author's own, copied verbatim from the pinned Stacks Project chapter file spaces-properties.tex; no statement or proof step was rewritten, reordered or removed.

Editorial note. Span recomputation by label pairing (LANE_BRIEF 10.2): in this chapter the author's proof environment follows the statement environment as a standalone block; the spec covers the complete environment pair (statement span, proof span) as recorded in the pool candidate.

Editorial note. Occupation scan: combined claim_key + statement_anchor + generic-line scan over all registration files (active and withdrawn) found 13 S10-geom published cards for spaces-properties.tex (prop-occupied.json, 2026-10-14); those 13 labels are excluded from this batch by design. For the 77 free labels: 0 occupied. The registry mentions field (7557 files) and focused-seeds (645) both point to the same S10-geom cards (dual-source check), no other occupation.

Source reference. Original lemma 0BAQ: sources/stacks-chow/tags.txt, source lines 14218–14218. Cross-chapter standard prerequisite cited by the author inside the counted span; it is used as a pointer and its own proof is not part of this item.

Lemma 0.1. *Let S be a scheme and let X be an algebraic space over S. The set of codimension 0 points of X is dense in $|X|$.*

Proof. If U is a scheme, then the set of generic points of irreducible components is dense in U (holds for any quasi-sober topological space). Thus if $U \to X$ is a surjective étale morphism, then the set of codimension 0 points of X is the image of a dense subset of $|U|$ (Lemma 0BAQ). Since $|X|$ has the quotient topology for $|U| \to |X|$ we conclude. □

Date: stacks-project:spaces-properties.tex:local-snapshot-20261012; GNU FDL 1.2 or later.