THE NUMBER OF CONNECTED COMPONENTS OF THE GENERIC FIBRE IS A LOWER BOUND IN A NEIGHBOURHOOD

THE STACKS PROJECT AUTHORS

Editorial note. Statement span is lines 7846-7853 of sources/stacks-more-morphisms/pinned-original-source.tex; proof span is lines 7855-7881, the complete author proof (25 body lines).

Editorial note. Navigation stubs: 5 cited-but-not-included labels are supplied as non-load-bearing pointers with their pinned tags where the snapshot tag index has them.

Editorial note. extra_packages is empty on purpose: the retained preamble already loads xy with options and a second declaration clashes in the assembler wrapper.

Editorial note. Text is retained verbatim, including the author's own typos and placeholders.

Allowed standard prerequisites (editorial).

- Explicit prerequisite boundary. Connected and irreducible components of a fibre and the lower bound on the irreducible component count; the argument is retained.

Source reference. Original lemma lemma-cover-generic-fibre-neighbourhood: sources/stacks-more-morphisms/pinned-original-source.tex, source lines 6658–6666. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma lemma-empty-generic-fibre: sources/stacks-more-morphisms/pinned-original-source.tex, source lines 6496–6501. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 01UA: sources/stacks-more-morphisms/pinned-original-source.tex, source lines 7881–7881. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 052A: sources/stacks-more-morphisms/pinned-original-source.tex, source lines 7881–7881. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Source reference. Original lemma 01ON: sources/stacks-more-morphisms/pinned-original-source.tex, source lines 7881–7881. Cited by the retained statement or proof but not part of the collected spans; supplied only as a navigation pointer. The argument does not depend on its content.

Date: stacks-project:more-morphisms.tex:local-snapshot-20261011; GNU FDL 1.2 or later.

Lemma 0.1. *Let $f : X \to Y$ be a morphism of schemes. Assume Y irreducible with generic point η and f of finite type. If X_η has n connected components, then there exists a nonempty open $V \subset Y$ such that for all $y \in V$ the fibre X_y has at least n connected components.*

Proof. As the question is purely topological we may replace X and Y by their reductions. In particular this implies that Y is integral, see Properties, Lemma 01ON. Let $X_\eta = X_{1,\eta} \cup \ldots \cup X_{n,\eta}$ be the decomposition of X_η into connected components. Let $X_i \subset X$ be the reduced closed subscheme whose generic fibre is $X_{i,\eta}$. Note that $Z_{i,j} = X_i \cap X_j$ is a closed subset of X whose generic fibre $Z_{i,j,\eta}$ is empty. Hence after shrinking Y we may assume that $Z_{i,j} = \emptyset$, see Lemma lemma-empty-generic-fibre. After shrinking Y some more we may assume that $X_y = \bigcup X_{i,y}$ for $y \in Y$, see Lemma lemma-cover-generic-fibre-neighbourhood. Moreover, after shrinking Y we may assume that each $X_i \to Y$ is flat and of finite presentation, see Morphisms, Proposition 052A. The morphisms $X_i \to Y$ are open, see Morphisms, Lemma 01UA. Thus there exists an open neighbourhood V of η which is contained in $f(X_i)$ for each i. For each $y \in V$ the schemes $X_{i,y}$ are nonempty closed subsets of X_y, we have $X_y = \bigcup X_{i,y}$ and the intersections $Z_{i,j,y} = X_{i,y} \cap X_{j,y}$ are empty! Clearly this implies that X_y has at least n connected components. $\square$