Forbidden homological minors in sufficiently connected PL manifolds

Sergey Avvakumov, Marguerite Bin, Xavier Goaoc

arxiv:2601.02920; CC BY 4.0

Editorial scope. The argument combines a homological Hanani-Tutte theorem with intersection-form labels and a hypergraph Ramsey extraction; a subdivision parity construction turns the assumed large simplex-skeleton minor into a forbidden Euclidean minor. Complete full author native mathematics and appendix context retained under own arxivCCBY4. All seven original geometric panels preserve their original page references. No independent correctness certification or author software rerun.

Allowed standard prerequisites (editorial).

- PL topology, homological intersection theory, Ramsey theory and standard convexity parameters. Original author statements and references preserved. Source-specific chain modifications, Ramsey parity reduction and graded parameter arguments included; standard cited prerequisites not recursively reproved.

1 Introduction

A classical line of research in discrete geometry investigates generalizations of properties of convex sets beyond convexity, with a particular attention to topological conditions. At least three distinct lines of inquiry emerged:

(A) Generalizing families of convex sets into acyclic/good covers, meaning set systems in topological spaces such that every

subfamily has empty or homologically/homotopically trivial intersection; an early example is Helly's topological theorem [15], see also the survey of Tancer [37] for an overview.

(B) Reformulating properties of convex sets of $\mathbb{R}^d$ as properties of linear maps into $\mathbb{R}^d$ and investigating their generalizations to continuous maps into $\mathbb{R}^d$, typically via theorems of Borsuk-Ulam type; an early example is the topological Radon theorem of Bajmóczy and Bárány [2] and more examples can be found e.g. in the survey of Bárány and Soberón [4].

(C) Analyzing set systems whose nerve enjoy properties of nerves of convex sets, like d-collapsibility or d-Lerayness; an early example is the sharpening of the Fractional Helly theorem by Kalai [19] and the work of Alon et al. [1] establishes several landmark results.

Two decades ago, Kalai and Meshulam [20] proposed conjectures relating approaches (A) and (C), towards what they called a theory of homological VC dimension. First, in order to generalize the notion of good cover, let us measure the complexity of the intersection patterns of a set system $\mathcal{F}$ in a topological space by its hth homological shatter function

$$\phi_{\mathcal{F}}^{(h)}\colon \left\{ \begin{array}{ccl} \mathbb{N} & \to & \mathbb{N}\cup\{\infty\} \\ k & \mapsto & \sup\left\{ \tilde{\beta}_i\left(\bigcap_{F\in\mathcal{G}} F;\mathbb{Z}_2\right) \;\middle|\; \mathcal{G}\subset\mathcal{F}, |\mathcal{G}|\leq k, 0\leq i\leq h \right\}. \end{array} \right. \tag{1}$$

Here h is some fixed parameter, and $\tilde{\beta}_i(\cdot;\mathbb{Z}_2)$ is the ith reduced Betti number with coefficients in $\mathbb{Z}_2$. (The set systems $\mathcal{F}$ with $\phi_{\mathcal{F}}^{(\infty)}\equiv 0$ are the acyclic covers and include good covers and convex sets.) The conjectures are about nerves of set systems whose intersection patterns have polynomially-growing topological complexity. In particular, their combination [20, Conjectures 6 and 7] (see also [13, Conjecture 1.9]) implies that polynomially-growing homological shatter functions give rise to a "positive density implies big clique" phenomenon:

Conjecture 1 (Kalai and Meshulam)**.** For any $d\in\mathbb{N}$ and any function $\Psi:\mathbb{N}\to\mathbb{N}$ such that $\Psi(n)=O(n^d)$, there exists $\beta:(0,1)\to(0,1)$

such that the following holds. For any $\alpha > 0$ and any set system $\mathcal{F}$ in $\mathbb{R}^d$ with $\phi_{\mathcal{F}}^{(d)} \leq \Psi$, if a proportion α of the $(d+1)$-element subsets of $\mathcal{F}$ have nonempty intersection, then some $\beta(\alpha)|\mathcal{F}|$ members of $\mathcal{F}$ have a point in common.

Such conditions that a positive density of d-faces in the nerve implies the existence of a linear-size face is called a fractional Helly theorem (see Section 1.1.1). Conjecture 1 is a topological analogue of a theorem of Matoušek [26] which asserts that every set system with polynomial (combinatorial) shatter function enjoys a fractional Helly theorem. Families of semi-algebraic sets of bounded complexity in $\mathbb{R}^d$ are an interesting example at the intersection of these two realms: classical results on the Betti numbers of algebraic varieties (see e.g. [5] and the references therein) ensure that both their combinatorial shatter functions and their homological shatter functions are polynomial. Conjecture 1 was confirmed for functions Ψ that are bounded ([13, Corollary 1.3], building on [16, 17, 30]). The main contribution of the present paper is to extend that confirmation to some diverging homological shatter functions and to set systems on certain manifolds.

1.1 Context and motivation

Before we state our results precisely (in Section 1.2) let us provide some context and motivation, as well as introduce some necessary terminology.

1.1.1 Combinatorial background: convexity parameters of set systems

The classical theorems of Helly, Radon and Carathéodory have initiated a rich theory of the combinatorial properties of convexity, whose landmarks include the centerpoint theorem, Tverberg's theorem, the colorful Helly and Carathéodory theorems, the fractional Helly theorem, the selection lemma, the weak ε-net theorem, the (p,q)-theorem, etc. We refer the interested reader to the monograph of Bárány [3] and the textbook of Matoušek [23]. These classical convexity theorems have algorithmic consequences for instance in optimization and geometric data

analysis [8, § 6 – 7] or in property testing [7], and one motivation for their extension beyond the convex setting is that several of these benefits generalize as well [25, 11]. We can associate to any set system $\mathcal{F}$ with ground set X some parameters inspired by convexity properties, for instance:

- The Helly number $\mathrm{h}_{\mathcal{F}}$ of $\mathcal{F}$ is the smallest integer h with the following property: If in a finite subfamily $\mathcal{G} \subset \mathcal{F}$, every h members of $\mathcal{G}$ intersect, then $\mathcal{G}$ has nonempty intersection. If no such h exists, we set $\mathrm{h}_{\mathcal{F}} = \infty$.

- The Radon number $\mathrm{r}_{\mathcal{F}}$ of $\mathcal{F}$ is the smallest integer r such that every r-element subset $S \subset X$ can be partitioned into two nonempty parts $S = P_1 \sqcup P_2$ such that $\mathrm{conv}_{\mathcal{F}}(P_1) \cap \mathrm{conv}_{\mathcal{F}}(P_2) \neq \emptyset$. (The set $\mathrm{conv}_{\mathcal{F}}(P)$, the $\mathcal{F}$-convex hull of a subset $P \subset X$, is the intersection of all the members of $\mathcal{F}$ that contain P.) If no such r exists, we set $\mathrm{r}_{\mathcal{F}} = \infty$.

Hence, letting $\mathcal{C}_d$ denote the set of all halfspaces in $\mathbb{R}^d$, Radon's lemma asserts that $\mathrm{r}_{\mathcal{C}_d} = d + 2$ and Helly theorem that $\mathrm{h}_{\mathcal{C}_d} = d + 1$. A classical result by Levi [22] asserts that for every set system $\mathcal{F}$ we have $\mathrm{h}_{\mathcal{F}} \leq \mathrm{r}_{\mathcal{F}} - 1$. (This is often stated for convexity spaces but it holds for set systems, see Appendix C.) Similar relations between such parameters have been investigated over the years, like for instance the partition conjecture of Eckhoff [10] refuted by Bukh [6]. Conjecture 1 pertains to a parameter inspired by the fractional Helly theorem [21, 19], which asserts that in $(d+1)$-wise intersection hypergraphs of convex sets of $\mathbb{R}^d$, positive density implies a linear-size clique. Here is the associated parameter:

- The fractional Helly number $\mathrm{fh}_{\mathcal{F}}$ of $\mathcal{F}$ is the smallest integer s such that there exists a function $\beta_{\mathcal{F}} : (0,1) \to (0,1)$ with the following property: For every finite subfamily $\mathcal{F}' \subset \mathcal{F}$, whenever a fraction α of the s-tuples of $\mathcal{F}'$ have nonempty intersection, a subset $\mathcal{G}$ of $\mathcal{F}'$ of size $\beta_{\mathcal{F}}(\alpha)|\mathcal{F}'|$ has nonempty intersection.

Again, the fractional Helly theorem states that $\mathrm{fh}_{\mathcal{C}_d} = d + 1$. The significance of the fractional Helly number was highlighted by Alon et al. [1], who proved that intersection-closed set systems

with bounded fractional Helly number enjoy a weak ε-net theorem, a Tverberg-type theorem, a selection lemma, etc. It is tempting to reformulate Conjecture 1 as

> For any function $\Psi : \mathbb{N} \to \mathbb{N}$ such that $\Psi(n) = O(n^d)$, every set set system $\mathcal{F}$ in $\mathbb{R}^d$ such that $\phi_{\mathcal{F}}^{(d)} \leq \Psi$ has fractional Helly number at most $d + 1$.

We note, however, that this statement is weaker than Conjecture 1 in that it does not assert that the function $\beta()$ underpinning the fractional Helly number depends only on Ψ and d.

1.1.2 Homological background: homological minors

A classical way to extend the theory of planar graphs is to consider embeddings of graphs into surfaces and of simplicial complexes into $\mathbb{R}^d$ or other topological spaces. There is a rich theory of embedding of graphs on surfaces, both structural (a classic being the Heawood inequality [34]) and computational (e.g. the use of graph genus for parameterized complexity). Let us mention, in particular, the strong Hanani-Tutte theorem which asserts that a graph is planar if it can be drawn so that every pair of independent edges cross an even number of times. This statement generalizes to the projective plane [32, 9] but was found to fail in genus 4 [12]. Going to dimension higher than 2 changes the nature of the problems drastically, already because Fáry's theorem no longer holds. (For every $d \geq 3$ there are simplicial complexes that embed in $\mathbb{R}^d$ piecewise linearly but not linearly.) It is thus sometimes convenient to relax the notion of embedding and work with chain maps, and this was done in particular to analyze intersection patterns [14, 30, 13] using a notion of homological minors [38]. Formally, the support of a singular chain is the union of (the images of) the singular simplices with nonzero coefficient in that chain, and the support of a simplicial chain is the subcomplex induced by the simplices with nonzero coefficient in that chain. We write $\mathrm{supp}(\sigma)$ for the support of a (singular or simplicial) chain σ. A chain map $a : C_*(K) \to \mathrm{C}_*^{\mathrm{sing}}(X)$ (resp. $a : C_*(K) \to \mathcal{C}_*(T)$) is nontrivial if, for every vertex v of K, the support of $a(v)$ has odd size. Two faces in a simplicial complex K are

adjacent if they have at least one vertex in common. A homological almost-embedding of a simplicial complex K into a topological space X (resp. into another simplicial complex L) is a nontrivial chain map $a : C_*(K) \to \mathrm{C}_*^{\mathrm{sing}}(X)$ (resp. $a : C_*(K) \to C_*(L)$) such that any two non-adjacent faces $\sigma, \tau \in K$ have images with disjoint support, that is $\mathrm{supp}(a(\sigma)) \cap \mathrm{supp}(a(\tau)) = \emptyset$. In particular, for every embedding f the associated chain map $f_\#$ is a homological (almost) embedding. Let Δ_N denote the N-dimensional simplex and for K a simplicial complex let $K^{(t)}$ denote its t-dimensional skeleton. It turns out that there is a homological version of the van Kampen-Flores theorem (see for instance [14, Corollary 14]):

Theorem 2. For any $d \geq 1$, $\Delta_{d+2}^{(\lceil d/2 \rceil)}$ does not homologically almost embed into $\mathbb{R}^d$.

A simplicial complex K is a homological minor of a topological space or simplicial complex X if there is a homological almost-embedding of K into X. Theorem 2 thus asserts that $\Delta_{d+2}^{(\lceil d/2 \rceil)}$ is not a homological minor of $\mathbb{R}^d$, i.e., that $\mathbb{R}^d$ has $\Delta_{d+2}^{(\lceil d/2 \rceil)}$ as forbidden homological minor.

1.1.3 Parameters of set systems with a forbidden homological minor

Matoušek [24] bounded the Helly number of topological set systems in $\mathbb{R}^d$ in which every subfamily intersects in a bounded number of connected components, all contractible. His approach starts from a set systems in $\mathbb{R}^d$, uses Ramsey theory to build a map from $\Delta_{d+2}^{(\lceil d/2 \rceil)}$ into $\mathbb{R}^d$ that is "constrained" by the known intersection patterns of $\mathcal{F}$ so that the intersection forced by the van Kampen-Flores theorem reveals a new intersection in $\mathcal{F}$. This approach was generalized by Goaoc et al. [14] to set systems in $\mathbb{R}^d$ of bounded $(\lceil d/2 \rceil)$-level topological complexity, where the h-level topological complexity of $\mathcal{F}$ is the maximum over $\mathbb{N}$ of the homological shatter function $\phi_{\mathcal{F}}^{(h)}$, that is

$$\mathrm{hc}_{\mathcal{F}}^{(h)} := \max \left\{ \max_{0 \leq i < h} \tilde{\beta}_i \left(\bigcap_{A \in \mathcal{G}} A; \mathbb{Z}_2 \right) \;\middle|\; \mathcal{G} \subset \mathcal{F} \right\}. \tag{2}$$

That generalization relied on homological minors and replaced the construction of the "constrained map" by the (simpler) construction of a "constrained chain map". The only aspect of the method that is specific to $\mathbb{R}^d$ is the use of the forbidden homological minor given by Theorem 2, so the approach readily generalizes to set systems whose ground set is a topological space with a forbidden homological minor, see the discussions in [30, §5.2] and [13, §2.2]. This method was refined and extended to analyze other parameters of set systems whose ground set has a forbidden homological minor [30, 28, 13].

1.1.4 Proof of Conjecture 1 for bounded homological shatter functions

Conjecture 1 was proven for bounded homological shatter functions in three steps:

- First, Holmsen and Lee [17, Theorem 1.1] proved that the fractional Helly number of any set system can be bounded by a function of its Radon number. Specifically, letting $\Psi_{\mathrm{r}\to\mathrm{fh}}(x)$ denote the supremum of the fractional Helly number of a set system with Radon number x, they proved that for every $r \geq 3$ we have $\Psi_{\mathrm{r}\to\mathrm{fh}}(r) \leq r^{r^{\lceil \log_2 r \rceil} + r \lceil \log_2 r \rceil}$.

- Second, Patáková [30, Theorem 2.1] proved that the Radon number of any set system whose ground set has K as forbidden homological minor can be bounded by a function of its $(\dim K)$-level topological complexity.

- Third, Goaoc, Holmsen and Patáková [13, Theorem 1.2] proved that for every set systems $\mathcal{F}$ whose ground set has K as forbidden homological minor, if the fractional Helly number $\mathrm{fh}_{\mathcal{F}}$ is bounded then it is at most $\mu(K)+1$, where $\mu(K)$ denotes the maximum sum of dimensions of two disjoint simplices in K.

1.2 Statement of the results

Our main contribution is to confirm Conjecture 1 for some diverging homological shatter functions and when the ground set

is a manifold. This decomposes into five independent results. Throughout the paper, we work with compact piecewise-linear (PL) manifolds (possibly with boundary); see [35, §1] for an introduction. We first generalize Theorem 2:

Theorem 3. For every integers $d \geq 3$ and b there exists $N = N(d, b)$ such that $\Delta_N^{(\lceil d/2 \rceil)}$ does not homologically almost embed in any compact, $(\lceil d/2 \rceil - 1)$-connected, d-dimensional PL manifold (possibly with boundary) $\mathcal{M}$ with $\beta_{\lceil d/2 \rceil}(\mathcal{M}; \mathbb{Z}_2) \leq b$.

This partially answers [30, Problem 3], [13, Conjecture 1.7] and [28, Conjecture 2] and extends the previous confirmation of Conjecture 1 from set systems in $\mathbb{R}^d$ to set systems on manifolds. This also extends several results on set systems with bounded topological complexity (Helly's theorem, Radon's theorem, (p, q)-theorem, ...) from $\mathbb{R}^d$ to sufficiently connected manifolds (see [30, 13]). We can relax the connectivity assumption (see Appendix B) but not remove it. One ingredient in the proof of Theorem 3 is the following analogue of the Hanani-Tutte theorem for homological almost-embeddings, which is of independent interest.

Theorem 4. Let K be a simplicial complex of dimension $k > 1$ and let $\mathcal{M}$ be a compact $2k$-dimensional PL manifold (possibly with boundary). If there exists a triangulation T of $\mathcal{M}$ and a non-trivial chain map $f : C_*(K; \mathbb{Z}_2) \to C_*(T; \mathbb{Z}_2)$ in general position such that the images of any two non-adjacent k-faces $\sigma, \tau \in K$ intersect in an even number of points, then K is a homological minor of $\mathcal{M}$.

We formalize what we mean by general position in Section 2. The homological shatter function $\Phi_{\mathcal{F}}^{(h)}$ defined in Equation (1) can be reformulated as a graded version of the topological complexity $\mathrm{hc}_{\mathcal{F}}^{(h)}$ defined in Equation (2), where the value for parameter t considers only intersections of subfamilies of size at most t: $\phi_{\mathcal{F}}^{(h)}(t) = \sup_{\substack{\mathcal{F}' \subset \mathcal{F} \\ |\mathcal{F}'| \leq t}} \mathrm{hc}_{\mathcal{F}'}^{(h)}$. We systematize this viewpoint and define graded analogues of other parameters of set systems. For instance here are the graded Radon and graded Helly numbers:

$$\mathrm{r}_{\mathcal{F}}(t) \coloneqq \sup_{\substack{\mathcal{F}' \subset \mathcal{F} \\ |\mathcal{F}'| \leq t}} \mathrm{r}_{\mathcal{F}'} \quad \text{and} \quad \mathrm{h}_{\mathcal{F}}(t) \coloneqq \sup_{\substack{\mathcal{F}' \subset \mathcal{F} \\ |\mathcal{F}'| \leq t}} \mathrm{h}_{\mathcal{F}'}\,. \tag{3}$$

It is straightforward to see that if the graded Helly numbers of a set system do not grow fast enough, then they are ultimately stationary and the set system has bounded Helly number (Lemma 3.1). We prove a similar condition for (graded) Radon numbers:

Theorem 5. Let $\mathcal{F}$ be a set system. If $\lim_{t\to\infty} \mathrm{r}_{\mathcal{F}}(t) - \log_2 t = -\infty$, then $\mathrm{r}_{\mathcal{F}} < \infty$.

As a application, we extend Patáková's theorem from bounded to sufficiently slowly diverging homological shatter function (Corollary 9). This, in turns yields fractional Helly theorems for sufficiently slowly diverging homological shatter functions.

Corollary 6. For every simplicial complex K there exists a function $\Psi_K : \mathbb{N} \to \mathbb{N}$ with $\lim_{t\to\infty} \Psi_K(t) = +\infty$ such that the following holds. If $\mathcal{F}$ is a set system whose ground set has K as forbidden homological minor and such that $\phi_{\mathcal{F}}^{(\dim K)}(t) \leq \Psi_K(t)$ for t large enough, then $\mathcal{F}$ has fractional Helly number at most $(\mu(K) + 1)$.

We then investigate more graded numbers and establish more relations between graded and ungraded numbers. As an application, we extend the Holmsen-Lee bound on $\Psi_{\mathrm{r}\to\mathrm{fh}}(\cdot)$. Let $\Xi : \mathbb{N} \to \mathbb{N}$ be the function defined by $\Xi(r) := r^{r^{\lceil \log_2 r \rceil} + r\lceil \log_2 r \rceil}$.

Theorem 7. Let $\Psi : \mathbb{N} \to \mathbb{N}$ and $t_0 \in \mathbb{N}$ such that $\Psi(t) < t+1$ for every $t \geq t_0$. If there exists an integer $t_1 \geq {t_0}^2$ such that $\Xi(\Psi(t_1)) < \frac{t_1}{t_0}$, then every set system whose graded Radon number function is bounded from above by Ψ has bounded fractional Helly number.

With Theorem 7 we can strengthen Corollary 6, as a close inspection of the proof reveals that the function β associated to the fractional Helly number depends only on Ψ and t_0. (It also allows a slightly faster growth than Corollary 9.) We postpone this to the full version of the paper.

2 On homological minors

In this section we prove Theorems 3 and 4. For completeness, we start with a consequence of the simplicial approximation theorem that allows us to work purely in simplicial homology:

Lemma 2.1. A simplicial complex K is a homological minor of a compact PL manifold (possibly with boundary) $\mathcal{M}$ if and only if K is a homological minor of some triangulation of $\mathcal{M}$.

Due to space limitation we defer the proof of Lemma 2.1 to Appendix A. When counting intersection points between chains, we focus on intersections that are stable under small perturbation. We therefore consider generic intersections, meaning intuitively that the chains intersect transversally. We formalize this in terms of linking numbers.[1] Let X be a triangulation of $\mathbb{S}^{2k-1}$. Let S_1 and S_2 be two subcomplexes of X with $|S_1|$ and $|S_2|$ homeomorphic to $\mathbb{S}^{k-1}$. Suppose that the simplicial complex $X \setminus S_2$ induced by X on the vertices not in S_2 has a geometric realization homotopy equivalent to $\mathbb{S}^{2k-1} \setminus \mathbb{S}^{k-1}$ (this can always be ensured up to taking a subdivision of X). The linking number of S_1 and S_2 in X is 1 if the homology class $[S_1]$ generates $H_{k-1}(X \setminus S_2; \mathbb{Z}_2) \cong \mathbb{Z}_2$, and 0 if $[S_1]$ is trivial in $H_{k-1}(X \setminus S_2; \mathbb{Z}_2)$. (Exchanging S_1 and S_2 in this definition yields the same result.) Let T be a triangulation of a compact PL manifold (possibly with boundary) $\mathcal{M}$. Two k-chains $z_1, z_2 \in C_k(T; \mathbb{Z}_2)$ intersect generically in vertex v if the closed star B of v in T satisfies: $B \cap z_1 \cap z_2 = \{v\}$, $D_1 := z_1 \cap B$ and $D_2 := z_2 \cap B$ are k-dimensional balls, and the spheres ∂D_1 and ∂D_2 have linking number 1 in ∂B. Two k-chains $z_1, z_2 \in C_k(T; \mathbb{Z}_2)$ are in general position if $z_1 \cap z_2$ consists of finitely many vertices, and z_1 and z_2 intersect generically in each of these vertices. Two k-chains $z_1, z_2 \in C_k(T; \mathbb{Z}_2)$ intersect evenly if they are in general position and $\mathrm{im}(z_1) \cap \mathrm{im}(z_2)$ has even size. For K a k-dimensional complex, a simplicial chain map $f : C_\bullet(K; \mathbb{Z}_2) \to C_\bullet(T; \mathbb{Z}_2)$ is in general position if for every non-adjacent k-faces $\sigma, \tau \in K$, the chains $f(\sigma)$ and $f(\tau)$ are in general position and for each vertex $v \in f(\sigma) \cap f(\tau)$, σ and τ are the only faces of K whose images under f intersect the closed star of v in T. We say that a simplicial map $f : K \to T$ is in general position if the associated chain map $f_\#$ is. For any compact PL manifold (possibly with boundary) $\mathcal{M}$ of dimension $2k$, there exists a map $\cap_\mathcal{M} : H_k(\mathcal{M}; \mathbb{Z}_2) \times H_k(\mathcal{M}; \mathbb{Z}_2) \to \mathbb{Z}_2$, called the intersection form of $\mathcal{M}$, such that $\cap_\mathcal{M}([z_1], [z_2]) \in \mathbb{Z}_2$ counts

[1] Intuitively, this generalizes the idea that in the plane, two curves cross at a point x if the branches of the curves alternate around x.

the intersection points of z_1 and z_2 modulo 2. We use no property of intersection forms besides their existence, and refer the interested reader to Prasolov [33, Chapter 2, §2.7] for a precise definition and to Paták and Tancer [29] for an brief account. The next result is due to Paták and Tancer [29, Proposition 21] and formulated following the presentation of Skopenkov [36], which is better suited for our purpose. (More precisely, we reformulated [36, Theorem 1.1.5] using [36, Lemma 2.1.1].) Note that the proofs by Paták-Tancer [29] and by Skopenkov [36] use only PL maps.

Theorem 8. Let L be a simplicial complex of dimension $k > 1$ and let $\mathcal{M}$ be a compact, $(k-1)$-connected, $2k$-dimensional PL manifold (possibly with boundary). The following statements are equivalent:

(i) There exists a triangulation T of $\mathcal{M}$ and a simplicial map $f : L \to T$ in general position such that the images of any two non-adjacent faces intersect evenly.

(ii) There exists a triangulation R of $\mathbb{R}^{2k}$ and a simplicial map $g : L \to R$ in general position and a map α that sends each k-face of L to an element of $H_k(\mathcal{M}; \mathbb{Z}_2)$ such that any two non-adjacent k-faces $\sigma, \tau \in L$ have images that intersect in an even number of points if and only if $\cap_{\mathcal{M}}(\alpha(\sigma), \alpha(\tau)) = 0$.

2.1 A homological Hanani-Tutte theorem

Let us now prove Theorem 4. Let K be a simplicial complex of dimension $k > 1$ and let T be a triangulation of a compact PL manifold (possibly with boundary) $\mathcal{M}$ of dimension $2k$. Let $f : C_*(K; \mathbb{Z}_2) \to C_*(T; \mathbb{Z}_2)$ be a non-trivial chain map in general position such that the images of any two non-adjacent k-faces $\sigma, \tau \in K$ intersect evenly. Our goal is to prove that K is a homological minor of $\mathcal{M}$. This requires repeatedly subdividing the triangulation T. In what follows, every time we subdivide a triangulation T_i into T_{i+1}, all chain maps to T_i and subcomplexes of T_i are also subdivided to T_{i+1}. Also, throughout the proof we identify every pure ℓ-dimensional simplicial complex with the unique

ℓ-chain with coefficients in $\mathbb{Z}_2$ it supports; we abuse the terminology and say that we add (pure) simplicial complexes over $\mathbb{Z}_2$ to mean that we add the corresponding chains. If f is a homological almost-embedding we are done. Otherwise, there exist some non-adjacent k-faces $\sigma, \tau \in K$ such that $f(\sigma) \cap f(\tau)$ is non-empty. By assumption, this intersection is a set of vertices of even size, so let $x, y \in f(\sigma) \cap f(\tau)$ be two distinct such vertices. Recall that f is in general position, $f(\sigma)$ and $f(\tau)$ intersect generically in x and in y. We set out to construct a refinement T' of T and a new map $f' : C_*(K; \mathbb{Z}_2) \to C_*(T'; \mathbb{Z}_2)$ that is also in general position, differs from f only on σ and τ, and satisfies $f'(\sigma) \cap f'(\tau) = (f(\sigma) \cap f(\tau)) \setminus \{x, y\}$ as well as $f'(\alpha) \cap f'(\beta) = f(\alpha) \cap f(\beta)$ for any $\alpha \in \{\sigma, \tau\}$ and any k-face $\beta \in K \setminus \{\sigma, \tau\}$. Iterating this procedure produces the announced homological almost-embedding of K into a triangulation of $\mathcal{M}$. Let B_x and B_y denote the closed stars of x




Figure 1: The setup for the construction of T' and f'.

and y in T. For $z \in \{x, y\}$ and $\alpha \in \{\sigma, \tau\}$ let $D_{\alpha,z} := \mathrm{supp}(f(\alpha)) \cap B_z$. Since $f(\sigma)$ and $f(\tau)$ intersect generically in x and y, the complexes $D_{\sigma,x}, D_{\sigma,y}, D_{\tau,x}$ and $D_{\tau,y}$ are k-dimensional balls, the $(k-1)$-spheres $\partial D_{\sigma,x}$ and $\partial D_{\tau,x}$ have linking number 1 in ∂B_x, and similarly $\partial D_{\sigma,y}$ and $\partial D_{\tau,y}$ have linking number 1 in ∂B_y. We subdivide T into T_1 so that the closed star B'_x of x and B'_y of y in T_1 are disjoint from ∂B_x and ∂B_y, respectively. In particular, for $z \in \{x, y\}$ and $\alpha \in \{\sigma, \tau\}$, letting $D'_{\alpha,z} := \mathrm{supp}(f(\alpha)) \cap B'_z$, the pair $(\partial D'_{\alpha,z}, \partial B'_z)$ is homeomorphic to $(\mathbb{S}^{k-1}, \mathbb{S}^{2k-1})$. The genericity of x and y in $f(\sigma) \cap f(\tau)$ is preserved through the subdivision $T \to T_1$ so the four complexes $D'_{\bullet,\bullet}$ are k-dimensional balls and their bounding spheres have the same linking numbers in $\partial B'_x$ and $\partial B'_y$ as their counterparts on ∂B_x and ∂B_y. Up to subdividing T_1 into T_2, there exist vertices v'_x and v'_y




Figure 2: The construction of B'.

in $\partial B'_x$ and $\partial B'_y$, respectively, and a path P' in the 1-skeleton of T_2 from v'_x to v'_y such that, letting $N(P')$ denote the closed star of P' in T_2, the closed star of $N(P')$ is disjoint from the image of f. The union $B' := B'_x \cup N(P') \cup B'_y$ is a ball. Observe that in $\partial B'$, the $(k-1)$-spheres $\partial D'_{\sigma,x}$ and $\partial D'_{\tau,x}$ retain the linking number 1 that they have on $\partial B'_x$. Similarly, in $\partial B'$, the $(k-1)$-spheres $\partial D'_{\sigma,y}$ and $\partial D'_{\tau,y}$ retain the linking number 1 that they have on $\partial B'_y$. Thus, up to further




Figure 3: The path P_σ.

subdividing T_2 into T_3, there exist a path P_σ in the 1-skeleton of $\partial B'$ that connects $\partial D'_{\sigma,x}$ to $\partial D'_{\sigma,y}$ and with relative interiors disjoint from the image of f. (Here we use that $\partial D'_{\alpha,z}$ is of codimension $k \geq 2$ in $\partial B'$.) Up to subdividing the triangulation further, we can pipe $D_{\sigma,x}$ and $D_{\sigma,y}$ together by a tube F_σ found in a neighborhood of the path P_σ [35, §5.10]. The tube $(F_\sigma, F_\sigma \cap D_{\sigma,x}, F_\sigma \cap D_{\sigma,y})$ is homeomorphic to $\big(\mathbb{D}^k \times [0,1], \mathbb{D}^k \times \{0\}, \mathbb{D}^k \times \{1\}\big)$. Moreover, by the (PL) general position theorem for embeddings [35, §5.3], we can take F_σ such that $\partial B'$ and ∂F_σ intersect in a generic way. By taking F_σ in a sufficiently small neighborhood U of P_σ so that $(U, \partial B' \cap U)$ is homeomorphic to $(\mathbb{R}^{2k}, \mathbb{R}^{2k-1} \times \{0\})$,

and the triple $(F_\sigma \cap \partial B', F_\sigma \cap D_{\sigma,x} \cap \partial B', F_\sigma \cap D_{\sigma,y} \cap \partial B')$ is homeomorphic to $\left(\mathbb{D}^{k-1} \times [0,1], \mathbb{D}^{k-1} \times \{0\}, \mathbb{D}^{k-1} \times \{1\}\right)$. Now, let C_σ be




Figure 4: The piping F_σ between $D_{\sigma,x}$ and $D_{\sigma,y}$ in a neighborhood of P_σ.

the sum $D_{\sigma,x} + \partial F_\sigma + D_{\sigma,y}$ (over $\mathbb{Z}_2$). Note that C_σ is contained in the union of $B_x \cup B_y$ and the closed star of $N(P')$, and therefore intersects the image of f only inside $B_x \cup B_y$. We note that $F_\sigma \cap \partial B'$ also pipes the $(k-1)$-spheres $D_{\sigma,x} \cap \partial B'$ and $D_{\sigma,y} \cap \partial B'$. Indeed, $(F_\sigma \cap \partial B', F_\sigma \cap D_{\sigma,x} \cap \partial B', F_\sigma \cap D_{\sigma,y} \cap \partial B') = (F_\sigma \cap \partial B', F_\sigma \cap \partial(D_{\sigma,x} \cap B'),\ F_\sigma \cap \partial(D_{\sigma,y} \cap B'))$ is homeomorphic to $\left(\mathbb{D}^{k-1} \times [0,1], \mathbb{D}^{k-1} \times \{0\}, \mathbb{D}^{k-1} \times \{1\}\right)$. It follows that $C_\sigma \cap \partial B'$ is homeomorphic to the connected sum of two $(k-1)$-spheres, and is therefore homeomorphic to a $(k-1)$-sphere. We claim that, in $\partial B'$, the




Figure 5: The chain $C_\sigma = D_{\sigma,x} + \partial F_\sigma + D_{\sigma,y}$ over $\mathbb{Z}_2$.

linking number between $C_\sigma \cap \partial B'$ and $D_{\tau,x} \cap \partial B'$ equals the linking number between $D_{\sigma,x} \cap \partial B'$ and $D_{\tau,x} \cap \partial B'$, that is 1. This follows from the fact that the chain C_σ differs from $D_{\sigma,x}$ by $\partial F_\sigma + D_{\sigma,y}$, and that each of $\partial F_\sigma \cap \partial B'$ and $D_{\sigma,y} \cap \partial B'$ is a boundary in $\partial B' \setminus D_{\tau,x}$. The same claim holds if we exchange x for y. Altogether, we get that $D_{\tau,x} \cap \partial B'$ and $D_{\tau,y} \cap \partial B'$ are in the same homology class in




Figure 6: The chain $\eta \in C_k(\partial B' \setminus C_\sigma; \mathbb{Z}_2)$ such that $\partial\eta = D_{\tau,x} \cap \partial B' + D_{\tau,y} \cap \partial B'$ over $\mathbb{Z}_2$.

$\partial B' \setminus C_\sigma$. Hence, their sum is a boundary, and there exists a chain $\eta \in C_k(\partial B' \setminus C_\sigma; \mathbb{Z}_2)$ such that the support of $\partial\eta$ is the sum over $\mathbb{Z}_2$ of $D_{\tau,x} \cap \partial B'$ and $D_{\tau,y} \cap \partial B'$. We finally set

$$f'(\sigma) = \underbrace{f(\sigma) + D_{\sigma,x} + D_{\sigma,y}}_{f(\sigma) \text{ with its restriction to } B_x \cup B_y \text{ removed}} + C_\sigma \quad \text{and} \quad f'(\tau) = \underbrace{f(\tau) + D'_{\tau,x} + D'_{\tau,y}}_{f(\tau) \text{ with its restriction to } B' \text{ removed}} + \eta$$

We set $f'(\omega) = f(\omega)$ for every other face $\omega \in K$. Extending f'




Figure 7: The chains C_σ and η reroute $f(\sigma)$ and $f(\tau)$ so as to remove intersections in B_x and B_y.

linearly yields a chain map, since both $f'(\sigma) - f(\sigma)$ and $f'(\tau) - f(\tau)$ are cycles [2]. The chain map f' is as announced: $f'(\sigma) \cap f'(\tau) = f(\sigma) \cap f(\tau) \setminus \{x, y\}$ and every other intersection remains unchanged. This concludes the proof of Theorem 4.

[2] They are even boundaries, ensuring that f' is chain homotopic to f

2.2 Forbidden homological minors for manifolds

We now prove Theorem 3. First, note that the odd-dimensional case reduces to the even-dimensional one. Indeed, for every $k \geq 2$, if $\mathcal{M}$ is a compact, $(k-1)$-connected, $(2k-1)$-dimensional PL manifold (possibly with boundary), then $\mathcal{M} \times [0,1]$ is a compact, $(k-1)$-connected, $2k$-dimensional PL manifold (possibly with boundary). Moreover, $\beta_k(\mathcal{M} \times [0,1]; \mathbb{Z}_2) = \beta_k(\mathcal{M}; \mathbb{Z}_2)$ and any homological minor of $\mathcal{M}$ is a homological minor of $\mathcal{M} \times [0,1]$. If the statement holds for d even and $b \in \mathbb{N}$ with some $N(b,d)$, then it holds with $d-1$ and b by putting $N(d-1,b) := N(d,b)$. So let us now consider the even-dimensional case. Let $k \geq 2$ and $b \in \mathbb{N}$, and let $\mathcal{M}$ be a compact, $(k-1)$-connected, $2k$-dimensional PL manifold (possibly with boundary). Let us fix N and suppose that $K = \Delta_N^{(k)}$ is a homological minor of $\mathcal{M}$. By Lemma 2.1, there exist a triangulation T of $\mathcal{M}$ and a (simplicial) homological almost-embedding $C_*(K) \to C_*(T)$. Actually, letting $S := T^{(k)}$, we have that there exists a homological almost-embedding $a : C_*(K; \mathbb{Z}_2) \to C_*(S; \mathbb{Z}_2)$. We apply Theorem 8 with $L = S$. Condition (i) holds with $T = S$ and f the identity, so Condition (ii) also holds. Hence, there exists a triangulation R of $\mathbb{R}^{2k}$, a simplicial map $g : S \to R$ in general position, and a map α that sends each k-face of S to an element of $H_k(\mathcal{M}; \mathbb{Z}_2)$ such that any two non-adjacent k-faces $\sigma, \tau \in S$ intersect evenly if and only if $\cap_{\mathcal{M}}(\alpha(\sigma), \alpha(\tau)) = 0$. We let $\tilde{\alpha} : C_k(S) \to H_k(\mathcal{M}; \mathbb{Z}_2)$ denote the linear extension of α. Consider the chain map $b : C_*(K; \mathbb{Z}_2) \to C_*(R; \mathbb{Z}_2)$ defined by $b = g_\# \circ a$. Note that b is nontrivial since a is nontrivial and g is a simplicial map. Moreover, the fact that b is a chain map in general position follows from three observations:

- since g is a simplicial map in general position, $g_\#$ is a chain map in general position,
- since a is a homological almost-embedding, it is also a chain map in general position, and
- the composition of a homological almost-embedding and a chain map in general position is a chain map in general position.

Notice that for $N \geq 2k+3$, not every independent k-faces of K can have images under b that intersect evenly. Indeed Theorem 4

would then imply that K is a homological minor of $\mathbb{R}^{2k}$, which would contradict the homological van Kampen-Flores theorem (Theorem 2). We use Ramsey's theorem to show that this contradiction can be reached for some subcomplex of K. So consider two non-adjacent k-faces $\sigma, \tau \in K$ and put $a(\sigma) = \sigma_1 + \sigma_2 + \ldots + \sigma_s$ and $a(\tau) = \tau_1 + \tau_2 + \ldots + \tau_t$. Since a is a homological almost-embedding, $\mathrm{supp}(a(\sigma))$ and $\mathrm{supp}(a(\tau))$ are disjoint, and σ_i and τ_j are thus non-adjacent for every $(i,j) \in [s] \times [t]$. Hence, given $(i,j) \in [s] \times [t]$, $g(\sigma_i)$ and $g(\tau_j)$ intersect evenly if and only if $\cap_{\mathcal{M}}(\alpha(\sigma_i), \alpha(\tau_j)) = 0$. We can thus count the intersections of $b(\sigma)$ and $b(\tau)$ (the following equalities are modulo 2 and $\mathrm{Card}\,()$ denotes the cardinal):

$$\begin{aligned}\mathrm{Card}\,(b(\sigma) \cap b(\tau)) &= \sum_{i\in[s],j\in[t]} \mathrm{Card}\,(g(\sigma_i) \cap g(\tau_j)) \\ &= \sum_{i\in[s],j\in[t]} \cap_{\mathcal{M}}(\alpha(\sigma_i), \alpha(\tau_j)) \\ &= \sum_{i\in[s]} \cap_{\mathcal{M}}(\alpha(\sigma_i), \tilde{\alpha}(a(\tau))) = \cap_{\mathcal{M}}(\tilde{\alpha}(a(\sigma)), \tilde{\alpha}(a(\tau)))\end{aligned}$$

Let $r = \beta_k(\mathcal{M}; \mathbb{Z}_2)$. The map $\beta : C_k(K) \to H_k(\mathcal{M}; \mathbb{Z}_2)$ defined by $\beta = \tilde{\alpha} \circ a$ induces a coloring of the k-simplices of K by the (at most 2^r) elements of $H_k(M; \mathbb{Z}_2)$. Let N' denote the number of vertices of $\mathrm{sd}\,\Delta^{(k)}_{2k+2}$. By the hypergraph Ramsey theorem, for N large enough (as a function of r and k) there exists a subset W of N' vertices in K such that β is constant over all k-simplices of $K[W]$; let us denote by $\square$ this constant value. In particular, for every k-faces σ, τ of $K[W]$ such that σ and τ are not adjacent, we have, modulo 2, $\mathrm{Card}\,(b(\sigma) \cap b(\tau)) = \cap_{\mathcal{M}}(\beta(\sigma_1), \beta(\tau_1)) = \cap_{\mathcal{M}}(\square, \square)$. Let us fix a bijection from the vertices of $\mathrm{sd}\,\Delta^{(k)}_{2k+2}$ to W and extend it to a chain map $j : C_*\left(\mathrm{sd}\,\Delta^{(k)}_{2k+2}\right) \to C_*(K[W])$. Also, let $h : C_*\left(\Delta^{(k)}_{2k+2}\right) \to C_*\left(\mathrm{sd}\,\Delta^{(k)}_{2k+2}\right)$ denote the subdivision chain map, where each i-face of $\Delta^{(k)}_{2k+2}$ is mapped to the sum of the i-faces of $\mathrm{sd}\,\Delta^{(k)}_{2k+2}$ that it contains. In particular, for every k-face σ of $\Delta^{(k)}_{2k+2}$, the chain $h(\sigma)$ is supported on $k!$ k-faces of $\mathrm{sd}\,\Delta^{(k)}_{2k+2}$. Let us examine the properties of $b \circ j \circ h$. First, it is a nontrivial chain map (because b, j and h are). Moreover, $j \circ h$ is a homological almost-embedding (since j and h are), and its composition with the chain

map in general position b yields a chain map in general position. Furthermore, any two non-adjacent k-faces $\sigma, \tau \in \Delta^{(k)}_{2k+2}$ have images under $b \circ j \circ h$ that intersect evenly. To see this, let us put $j \circ h(\sigma) = \sigma_1 + \sigma_2 + \ldots + \sigma_s$ and $j \circ h(\tau) = \tau_1 + \tau_2 + \ldots + \tau_s$. Since $j \circ h$ is a homological almost-embedding, $\mathrm{supp}(j \circ h(\sigma))$ and $\mathrm{supp}(j \circ h(\tau))$ are disjoint. It follows that for every $i, j \in [s]$ the k-simplices σ_i and τ_j are non-adjacent. We therefore have, modulo 2,

$$\begin{aligned}\mathrm{Card}\,(b \circ j \circ h(\sigma) \cap b \circ j \circ h(\tau)) &= \sum_{i,j \in [s]} \mathrm{Card}\,(b(\sigma_i) \cap b(\tau_j)) \\ &= \sum_{i,j \in [s]} \cap_{\mathcal{M}}(\beta(\sigma_i), \beta(\tau_j)) = \sum_{i,j \in [s]} \cap_{\mathcal{M}}(\square, \square).\end{aligned}$$

To sum up, the cardinal of $b \circ j \circ h(\sigma) \cap b \circ j \circ h(\tau)$ has the same parity as $\cap_{\mathcal{M}}(\square, \square) s^2$. Since $s = k!$ is even, σ and τ have images under $b \circ j \circ h$ that intersect evenly. To conclude, $b \circ j \circ h$ is a nontrivial chain map in general position from $\Delta^{(k)}_{2k+2}$ to R such that independent faces have images that intersect evenly. By Theorem 4 this means that $\Delta^{(k)}_{2k+2}$ is a homological minor of $\mathbb{R}^{2k}$, a contradiction with Theorem 2. Thus, for N large enough, the initial hypothesis that $\Delta^{(k)}_N$ is a homological minor of $\mathcal{M}$ cannot be true.

3 Graded parameters of set systems

In this section we present our contributions on set systems of parameters.

3.1 Graded Radon and Helly numbers

Each relation between parameters of a set system yields a relation between their graded analogues. From Levi's inequality we get the following inequality between the graded Helly and Radon numbers (defined in Section 1.1.1):

$$\forall t \in \mathbb{N}, \qquad \mathrm{h}_{\mathcal{F}}(t) = \sup_{\substack{\mathcal{F}' \subset \mathcal{F} \\ |\mathcal{F}'| \leq t}} \mathrm{h}_{\mathcal{F}'} \leq \sup_{\substack{\mathcal{F}' \subset \mathcal{F} \\ |\mathcal{F}'| \leq t}} (\mathrm{r}_{\mathcal{F}'} - 1) = \mathrm{r}_{\mathcal{F}}(t) - 1. \qquad (4)$$

It follows from the definitions of graded parameters that each one is a non-decreasing function that converges to the ungraded

parameter (possibly ∞). We notice that if a graded Helly number is asymptotically sublinear then it is bounded:

Lemma 3.1. Let $\mathcal{F}$ be a set system and $t_0 \in \mathbb{N}$. If $\mathrm{h}_{\mathcal{F}}(t) < t$ for all $t > t_0$, then $\mathrm{h}_{\mathcal{F}} \leq t_0$.

Proof. By definition, we have $\mathrm{h}_{\mathcal{F}}(t) \leq t$ for every $t \in \mathbb{N}$. Moreover, $\mathrm{h}_{\mathcal{F}}(t) \neq \mathrm{h}_{\mathcal{F}}(t-1)$ if and only if $\mathrm{h}_{\mathcal{F}}(t) = t$. The assumption and a straightforward induction therefore implies that for every $t > t_0$ we have $\mathrm{h}_{\mathcal{F}}(t) = \mathrm{h}_{\mathcal{F}}(t_0) \leq t_0$. □

When graded Radon numbers grow, it is at most linearly (see Appendix D) and not too slowly:

Lemma 3.2. Let $\mathcal{F}$ be a set system and $t \geq 2$ an integer. If $\mathrm{r}_{\mathcal{F}}(t) > \mathrm{r}_{\mathcal{F}}(t-1)$, then $\mathrm{r}_{\mathcal{F}}(t-1) \geq 1 + \log_2\left(1 + \frac{t}{\mathrm{h}_{\mathcal{F}}(t)}\right)$.

Proof. Let X denote the ground set of $\mathcal{F}$ and let $n = \mathrm{r}_{\mathcal{F}}(t-1)$. Suppose that $\mathrm{r}_{\mathcal{F}}(t) > n$, so that there exist a subset $\mathcal{G} = \{G_1, G_2, \ldots, G_t\} \subset \mathcal{F}$ and a subset $S \subseteq X$ of size n such that

(i) there is no partition of S into two parts whose $\mathcal{G}$-convex hulls intersect,

(ii) for every $i \in [t]$, there exists a partition $\mathcal{P}_i$ of S into two parts whose $(\mathcal{G} \setminus \{G_i\})$-convex hulls intersect.

Recall that given $\mathcal{F}' \subseteq \mathcal{F}$, the $\mathcal{F}'$-convex hull $\mathrm{conv}_{\mathcal{F}'}(P)$ of a subset $P \subset X$ is the intersection of all the members of $\mathcal{F}'$ that contain P. In particular, for any $P \subseteq X$ such that $P \not\subseteq G_i$ we have $\mathrm{conv}_{\mathcal{G}}(P) = \mathrm{conv}_{\mathcal{G}\setminus\{G_i\}}(P)$. Conditions (i) and (ii) therefore imply that every $G_i \in \mathcal{G}$ contains one or the other part of $\mathcal{P}_i$. There are at most $2^{n-1} - 1$ partitions of S in two nonempty parts. Let us assume that $t > (2^{n-1} - 1)h$ for some integer h, so that by the pigeonhole principle there exist $h+1$ indices $i_1, i_2, \ldots, i_{h+1}$ such that the partitions $\mathcal{P}_{i_1}, \mathcal{P}_{i_2}$, ..., $\mathcal{P}_{i_{h+1}}$ coincide. Let $\{P_1, P_2\}$ be that partition of S. Let us put $\mathcal{G}' = \{A \in \mathcal{G} \colon P_1 \subseteq A \text{ or } P_2 \subseteq A\}$. We make two observations:

- $\cap_{A \in \mathcal{G}'} A$ coincides with $\mathrm{conv}_{\mathcal{G}}(P_1) \cap \mathrm{conv}_{\mathcal{G}}(P_2)$ and is therefore empty.

- every choice of h elements in $\mathcal{G}'$ has nonempty intersection. Indeed, $\mathcal{G}'$ contains G_{i_1}, G_{i_2}, ..., $G_{i_{h+1}}$, so that any choice of h elements from $\mathcal{G}'$ is bound to miss G_{i_j} for at least one $j \in [h+1]$ and their intersection must then contain $\mathrm{conv}_{\mathcal{G}\setminus\{G_{i_j}\}}(P_1) \cap \mathrm{conv}_{\mathcal{G}\setminus\{G_{i_j}\}}(P_2) \neq \emptyset$.

For $h \geq \mathrm{h}_{\mathcal{F}}(t)$ these conditions are incompatible. We therefore have $t \leq (2^{n-1}-1)\,\mathrm{h}_{\mathcal{F}}(t)$, and the statement follows. □

We can now prove that any set system with sufficiently slowly growing graded Radon numbers has finite Radon number.

Proof of Theorem 5. Let $\mathcal{F}$ be a set system with infinite Radon number. If the Helly number $\mathrm{h}_{\mathcal{F}}$ is also infinite, then by Lemma 3.1 there exists an increasing sequence $\{t_i\}_{i\in\mathbb{N}}$ such that $\mathrm{h}_{\mathcal{F}}(t_i) = t_i$, and therefore $\mathrm{r}_{\mathcal{F}}(t_i) \geq t_i + 1$ by Inequality (4); this prevents $\mathrm{r}_{\mathcal{F}}(t) - \log_2 t$ from going to $-\infty$ as $t \to \infty$. So suppose that the Helly number $\mathrm{h}_{\mathcal{F}}$ is finite. The assumption that $\mathrm{r}_{\mathcal{F}} = \infty$ ensures that there exists an increasing sequence $\{t_i\}_{i\in\mathbb{N}}$ such that $\mathrm{r}_{\mathcal{F}}(t_i) > \mathrm{r}_{\mathcal{F}}(t_i - 1)$. Lemma 3.2 implies that $\mathrm{r}_{\mathcal{F}}(t_i) > \mathrm{r}_{\mathcal{F}}(t_i - 1) \geq \log_2 t_i - \log_2 \mathrm{h}_{\mathcal{F}}$. Again, this prevents $\mathrm{r}_{\mathcal{F}}(t) - \log_2 t$ from going to $-\infty$ as $t \to \infty$. The statement follows by contraposition. □

3.2 Consequences for topological set systems

Let us finally consider topological set systems with slowly growing homological shatter function and ground set with a forbidden homological minor.

Corollary 9. For every simplicial complex K there exists a function $S_K : \mathbb{N} \to \mathbb{N}$ with $\lim_{t\to\infty} S_K(t) = +\infty$ such that the following holds. Any set system $\mathcal{F}$ whose ground set has K as forbidden homological minor and satisfies $\phi_{\mathcal{F}}^{(\dim K)}(t) \leq S_K(t)$ for t large enough has finite Radon number.

Proof. Recall that $\Psi^{(K)}_{\mathrm{hc}\to\mathrm{r}}(x)$ denotes the supremum of the Radon number of a set system with $(\dim K)$-level topological complexity at most x and whose ground set has K as forbidden homological minor. Patáková [30] proved that $\Psi^{(K)}_{\mathrm{hc}\to\mathrm{r}}(x)$ is finite for every K and x. For $t \in \mathbb{N}$, we define $S_K(t) = \max\{x \in \mathbb{N} \mid \Psi^{(K)}_{\mathrm{hc}\to\mathrm{r}}(x) \leq \frac{1}{2}\log_2 t\}$.

This ensures that $\Psi^{(K)}_{\mathrm{hc}\to\mathrm{r}}(S_K(t)) \leq \frac{1}{2}\log_2 t$ for every $t \in \mathbb{N}$. Observe that $\lim_{t\to\infty} S_K(t) = \infty$ since $\Psi^{(K)}_{\mathrm{hc}\to\mathrm{r}}(x)$ is finite for every $x \in \mathbb{N}$. Now consider a set system $\mathcal{F}$ with function $\phi^{(\dim K)}_{\mathcal{F}} \leq S_K$ and whose ground set has K as forbidden homological minor. Let $t \in \mathbb{N}$ and consider a subset $\mathcal{F}' \subseteq \mathcal{F}$ of size t. The ground set of $\mathcal{F}'$ also has K as forbidden homological minor. Moreover, $\mathcal{F}'$ has $(\dim K)$-level topological complexity at most $\phi^{(\dim K)}_{\mathcal{F}}(t) \leq S_K(t)$. It follows that $\mathrm{r}_{\mathcal{F}'} \leq \Psi^{(K)}_{\mathrm{hc}\to\mathrm{r}}(S_K(t)) \leq \frac{1}{2}\log_2 t$. This holds for every $\mathcal{F}' \subseteq \mathcal{F}$ of size t, so $\mathrm{r}_{\mathcal{F}}(t) \leq \frac{1}{2}\log_2 t$. This inequality holds for every $t \in \mathbb{N}$ so Theorem 5 implies that $\mathcal{F}$ has bounded Radon number. □

We can finally prove a fractional Helly theorem for diverging homological shatter functions.

Proof of Corollary 6. Recall that $\Psi_{\mathrm{r}\to\mathrm{fh}}(y)$ denotes the supremum of the fractional Helly number of a set system with Radon number y. Holmsen and Lee [17, Theorem 1.1] proved that $\Psi_{\mathrm{r}\to\mathrm{fh}}(y)$ is finite for every y. Let S_K denote the function from Corollary 9. Now consider a set system $\mathcal{F}$ whose ground set has K as forbidden homological minor and satisfies $\phi^{(\dim K)}_{\mathcal{F}} \leq S_K$. Corollary 9 ensures that $\mathrm{r}_{\mathcal{F}}$, the Radon number of $\mathcal{F}$, is finite. It follows that the fractional Helly number of $\mathcal{F}$ is at most $\Psi_{\mathrm{r}\to\mathrm{fh}}(\mathrm{r}_{\mathcal{F}})$, and is therefore finite. From there, [13, Theorem 1.2] ensures that this fractional Helly number is at most $\mu(K)+1$. □

3.3 Other graded parameters and relations

With the intersection hypergraph of $\mathcal{F}$ in mind, we say that a set $\mathcal{G}$ in a set system $\mathcal{F}$ is a clique if the intersection of all members of $\mathcal{G}$ is non-empty. The colorful Helly theorem suggests the following parameter:

- The colorful Helly number $\mathrm{ch}_{\mathcal{F}}$ of $\mathcal{F}$ is the smallest number of colors m such that for every coloring of a subfamily $\mathcal{F}' \subset \mathcal{F}$ with m colors, if every subfamily that contains exactly one element of each color is a clique, then at least one color class is a clique.

We say that a set $\mathcal{G}$ in a set system $\mathcal{F}$ is a c-wise clique if every c-element subset of $\mathcal{G}$ is a clique. Clearly a clique is a c-wise clique,

and when $c \geq \mathrm{h}_{\mathcal{F}}$ the converse is true. To analyze set systems with large, infinite or unknown Helly numbers, it is useful to consider variants of the colorful and fractional Helly numbers where cliques are replaced by c-wise cliques:

- The cth colorful Helly number $\mathrm{ch}_{\mathcal{F}}^{(c)}$ of $\mathcal{F}$ is the smallest number of colors $m \geq c$ such that for every coloring of a subfamily $\mathcal{F}' \subset \mathcal{F}$ with m colors, if every subfamily of $\mathcal{F}'$ that contains exactly one element of each color forms a c-wise clique, then at least one color class is a c-wise clique.

- The cth fractional Helly number $\mathrm{fh}_{\mathcal{F}}^{(c)}$ of $\mathcal{F}$ is the smallest integer s such that there exists a function $\beta_{\mathcal{F}} : (0,1) \to (0,1)$ with the following property: For every finite subfamily $\mathcal{F}' \subset \mathcal{F}$, whenever a fraction α of the s-tuples of $\mathcal{F}'$ forms a c-wise clique, a subset $\mathcal{G}$ of $\mathcal{F}'$ of size $\beta_{\mathcal{F}}(\alpha)|\mathcal{F}'|$ forms a c-wise clique.

Obviously for every set system $\mathcal{F}$, if $c \geq \mathrm{h}_{\mathcal{F}}$, then $\mathrm{fh}_{\mathcal{F}}^{(c)} = \mathrm{fh}_{\mathcal{F}}$ and $\mathrm{ch}_{\mathcal{F}}^{(c)} = \mathrm{ch}_{\mathcal{F}}$. Holmsen [16, Theorem 1.2] proved that $\mathrm{fh}_{\mathcal{F}}^{(c)} \leq \mathrm{ch}_{\mathcal{F}}^{(c)}$ for every set system $\mathcal{F}$ and every $c \in \mathbb{N}$. A close inspection of that proof provides another bridge between the graded and ungraded parameters:

Lemma 3.3. Let $\ell > c$ be integers. Every set system $\mathcal{F}$ such that $\mathrm{ch}_{\mathcal{F}}^{(c)}(c\ell) \leq \ell$ satisfies $\mathrm{fh}_{\mathcal{F}}^{(c)} \leq \mathrm{ch}_{\mathcal{F}}^{(c)}(c\ell)$.

Proof. Let $c < \ell$, let $\mathcal{F}$ be a set system and let $\ell' := \mathrm{ch}_{\mathcal{F}}^{(c)}(c\ell)$. By definition of $\mathrm{ch}_{\mathcal{F}}^{(c)}(\cdot)$, the c-uniform hypergraph recording which c-element subsets of $\mathcal{F}$ form cliques cannot contain a certain pattern on $c\ell'$ vertices, namely the complete ℓ'-tuples of missing edges [17, §3]. This is the only property needed to ensure that $\mathrm{fh}_{\mathcal{F}}^{(c)} \leq \ell'$ [16, Theorem 1.2]. □

Let $\Psi_{\mathrm{r}\to\mathrm{ch}}^{(c)}(x)$ denote the supremum of the cth colorful Helly number of a set system with Radon number x. Holmsen and Lee [17, Theorem 2.2] proved that $\Psi_{\mathrm{r}\to\mathrm{ch}}^{(c)}(x) \leq \max(\Xi(x), c)$ for every $c \geq x - 1$, where $\Xi : \mathbb{N} \to \mathbb{N}$ is the function defined by

$\Xi(r) := r^{r^{\lceil \log_2 r \rceil} + r\lceil \log_2 r \rceil}$. Applying this inequality to subsets of size t we get:

$$\mathrm{ch}_{\mathcal{F}}^{(c)}(t) \leq \max\left(\Xi\left(\mathrm{r}_{\mathcal{F}}(t)\right), c\right) \quad \text{for every } \mathcal{F}, c, t \text{ such that } c \geq \mathrm{h}_{\mathcal{F}}(t). \tag{5}$$

We can now prove that sufficiently slowly growing graded Radon numbers imply a bounded Fractional Helly number.

Proof of Theorem 7. Let $\Psi : \mathbb{N} \to \mathbb{N}$ and $t_0 \in \mathbb{N}$ such that $\Psi(t) < t+1$ for every $t \geq t_0$. Also suppose that there exists an integer $t_1 \geq {t_0}^2$ such that $\Xi\left(\Psi(t_1)\right) < \frac{t_1}{t_0}$. We now consider a set system $\mathcal{F}$ such that $\mathrm{r}_{\mathcal{F}}(t) \leq \Psi(t)$ for every $t \in \mathbb{N}$ and argue that $\mathrm{fh}_{\mathcal{F}}$, the fractional Helly number of $\mathcal{F}$, is bounded. By Levi's inequality (4) we have $\mathrm{h}_{\mathcal{F}}(t) \leq \mathrm{r}_{\mathcal{F}}(t) - 1 \leq \Psi(t) - 1$. It follows that $\mathrm{h}_{\mathcal{F}}(t) < t$ for every $t \geq t_0$, so by Lemma 3.1 we have $\mathrm{h}_{\mathcal{F}} \leq t_0$. Hence, the graded version (5) of the Holmsen-Lee inequality applies with $c = t_0$ and every t, that is $\mathrm{ch}_{\mathcal{F}}^{(t_0)}(t) \leq \max\left(\Xi\left(\Psi(t)\right), t_0\right)$. Let us apply Lemma 3.3 with $c = t_0$ and $\ell = \frac{t_1}{t_0}$. Observe that $\ell > c$ holds because $t_1 > {t_0}^2$, and that $\mathrm{ch}_{\mathcal{F}}^{(c)}(c\ell) \leq \ell$ follows from the assumption that $\Xi\left(\Psi(t_1)\right) \leq \frac{t_1}{t_0}$, as

$$\mathrm{ch}_{\mathcal{F}}^{(c)}(c\ell) = \mathrm{ch}_{\mathcal{F}}^{(t_0)}(t_1) \leq \max\left(\Xi\left(\Psi(t_1)\right), t_0\right) \leq \frac{t_1}{t_0} = \ell.$$

Hence, $\mathrm{fh}_{\mathcal{F}}^{(t_0)} \leq \mathrm{ch}_{\mathcal{F}}^{(t_0)}(t_1)$. Since $\mathrm{h}_{\mathcal{F}} \leq t_0$ we have $\mathrm{fh}_{\mathcal{F}}^{(t_0)} = \mathrm{fh}_{\mathcal{F}}$ and the statement follows. □

A Homological minors of manifolds and their triangulations

Here we give the proof of:

Lemma 8. A simplicial complex K is a homological minor of a compact PL manifold (possibly with boundary) $\mathcal{M}$ if and only if K is a homological minor of some triangulation of $\mathcal{M}$.

Proof. Let K and L be simplicial complexes. A map $\tau' : |K| \to |L|$ is an approximation of a map $\tau : |K| \to |L|$ with respect to L if for every $x \in |K|$, $\tau'(x)$ belongs to the inclusion-minimal closed simplex

of L containing the point $\tau(x)$. (See [39].) The relative simplicial approximation theorem [39] states that any continuous map into $|L|$ whose restriction to the boundary is simplicial admits an approximation (with respect to L) that is a simplicial map leaving the boundary unchanged. Moreover, it gives a relative homotopy (fixing the boundary) between the continuous map and the simplicial approximation, which remains an approximation (with respect to L) for every $t \in [0,1]$. Now, let $a : C_*(K) \to \mathrm{C}_*^{\mathrm{sing}}(|T|)$ be a homological almost-embedding and let T be a triangulation of $\mathcal{M}$ fine enough that no closed simplex of T intersects the supports $\mathrm{supp}(a(\sigma))$ and $\mathrm{supp}(a(\tau))$ of two non-adjacent faces σ, τ of K. We apply the relative simplicial approximation theorem to construct a simplicial chain map $b\colon C_*(K) \longrightarrow C_*(T)$ satisfying the following property: for every $\sigma \in K$, denoting $a(\sigma) = \tau_0 + \tau_1 + \cdots + \tau_s$, we have $b(\sigma) = \tau_0' + \tau_1' + \cdots + \tau_s'$, where each τ_i' is a simplicial approximation of the singular simplex τ_i. Our choice of T then guarantees that b is again a homological almost-embedding. Observe that the chain map a is supported on a finite number of singular simplices $\tau : \Delta^i \to \mathcal{M}$. We inductively construct simplicial approximations of all such map, starting from $i = 0$ to $i = \dim(K)$. In the process, we make sure that the approximation τ' of each singular simplex τ is compatible with the other approximations in the following sense: if $\partial\tau = \omega_0 + \omega_1 + \ldots + \omega_t$, we require the approximation τ' to satisfy $\partial \circ \tau' = \omega_0' + \omega_1' + \ldots + \omega_t'$, where the ω_j' are the previously chosen approximations. Specifically, suppose that we are processing a map $\tau : \Delta^t \to \mathcal{M}$. Let $\omega_0, \omega_1, \ldots, \omega_t$ denote the facets of τ. By induction, we assume that we already computed a suitable simplicial approximation ω_i' of ω_i for $0 \leq i \leq t$. We cannot apply the relative simplicial approximation theorem to τ directly, as it may not be simplicial on the boundary $\partial\Delta^t$. We therefore define an auxiliary map $\widetilde{\tau} : \Delta^t \to \mathcal{M}$ with boundary is $\omega_0' + \omega_1' + \ldots + \omega_t'$ by concatenation of the map τ and the homotopy between $\tau \circ \partial$ and $\tau' \circ \partial$ given by the previous use of simplicial approximation theorem to each map constituting $\partial\tau$. We then apply the relative simplicial approximation theorem to $\widetilde{\tau}$. The simplicial map we obtain satisfies our compatibility condition. So suppose that we have our collection of simplicial approximations of all singular simplices in the support of the chain map a, where the simplicial approximations of

τ being denoted τ'. For every $\sigma \in K$, letting $a(\sigma) = \tau_1 + \cdots + \tau_s$, we put $b(\sigma) = \tau'_1 + \cdots + \tau'_s$. It is a chain map by the compatibility of our simplicial approximations, it is nontrivial because a is, and the images of independent simplices have disjoint supports by our choice of T and the property of simplicial approximations. Altogether, b is a homological almost-embedding of K into T. □

B Theorem 3 for relaxed connectivity assumptions

In this appendix we relax the conditions in Theorem 3 that the d-manifold be $(\lceil d/2 \rceil - 1)$-connected. A classical method to increase connectivity of a manifold is to use surgeries. Let $\mathcal{M}$ a d-manifold and let $p \leq d$. A manifold $\mathcal{M}'$ is produced by a p-surgery from $\mathcal{M}$ if there exists an embedding $\phi : \mathbb{S}^p \times \mathbb{D}^q \to \mathcal{M}$, with $p + q = d$, such that $\mathcal{M}' = \left(M \backslash (\mathring{\text{im}}(\phi))\right) \cup_{f_{|\mathbb{S}^p \times \mathbb{S}^{q-1}}} \left(\mathbb{D}^{p+1} \times \mathbb{S}^{q-1}\right)$. It turns out that surgeries do not destroy homological minors of sufficiently small dimension.

Lemma B.1. Let K be a k-dimensional homological minor of a $2k$-manifold (possibly with boundary) $\mathcal{M}$. If $\mathcal{M}'$ is a manifold produced by a p-surgery from $\mathcal{M}$, with $p < k$, then K is a homological minor of $\mathcal{M}'$.

Proof. Let $a : C_*(K) \to \mathrm{C}_*^{\text{sing}}(\mathcal{M})$ be a homological almost-embedding from K to $\mathcal{M}$. We can assume that $A := \bigcup_{\omega \in K} \text{supp}(a(\omega))$ is a simplicial subcomplex inside some triangulation of $\mathcal{M}$. Let $\mathcal{M}'$ be a manifold produced by a p-surgery from $\mathcal{M}$. That is, there exists an embedding $\phi : \mathbb{S}^p \times \mathbb{D}^q \to \mathcal{M}$, with $q = 2k - p$, such that $\mathcal{M}' = \left(M \backslash (\mathring{\text{im}}(\phi))\right) \cup_{f_{|\mathbb{S}^p \times \mathbb{S}^{q-1}}} \left(\mathbb{D}^{p+1} \times \mathbb{S}^{q-1}\right)$. We define $\phi_0 : \mathbb{S}^p \to \mathcal{M}$ by $\phi_0(x) = \phi(x, 0)$ (which we can assume PL) and by the (PL) general position theorem for embeddings [35, §5.3], there is a homeomorphism $h : \mathcal{M} \to \mathcal{M}$ (a perturbation of the identity) such that $dim\,(h(A) \cap im(\phi_0)) \leq p + k - 2k < 0$. The chain map $b = h_{\#} \circ a$ is a homological almost-embedding from K to $\mathcal{M} \setminus im(\phi_0)$. Since $\mathcal{M} \setminus im(\phi_0)$ embeds into $\mathcal{M}'$ via the inclusion i, the same chain map the chain map $i_{\#} \circ b$ shows that K is a homological minor of $\mathcal{M}'$. □

A manifold is called stably parallelizable or a π-manifold if its tangent bundle is stably parallelizable, meaning that its sum with some trivial bundle is trivial (see [27]). Milnor proved [27, Corollary to Theorem 2] that any compact triangulated differentiable π-manifold of dimension $2k$ can be made $(k-1)$-connected by a finite sequence of p-surgeries with $p < k$. The same result holds [27, Corollary to Theorem 3] for any compact triangulated differentiable manifold that is $(k-1)$-parallelizable (i.e, if its tangent bundle restricted to its $(k-1)$-skeleton is trivial). Lemma B.1 ensure that these surgeries preserve k-dimensional homological minors, so we get the following generalization of Theorem 3:

Theorem 10. Let $\mathcal{M}$ be a compact PL manifold (possibly with boundary) of dimension d. If $\mathcal{M}$ is a π-manifold or is $(k-1)$-parallelizable, then there exists N such that $\Delta_N^{\left(\lceil \frac{d}{2} \rceil\right)}$ is not a homological minor of $\mathcal{M}$.

For example, $\mathbb{T}^d$ is parallelizable for every $d \geq 1$, so for $d \geq 3$ the manifold $\mathbb{T}^d$ is covered by Theorem 10 (but not by Theorem 3). Also, any closed orientable 3-manifold is parallelizable, and any product of stably parallelizable manifolds is stably parallelizable. Examples of manifolds that are covered by neither Theorem 3 nor Theorem 10 is $\mathbb{RP}^d$ for even values of d.

C Convexity spaces VS set systems

In this appendix, we provide for completeness a discussion of why the results of Levi [22] and Holmsen and Lee [17] stated for convexity spaces hold in the context of set systems. See also the discussion in the v3 of Patakova's preprint [31, Appendix A]. A common setting for studying generalizations of convexity is to consider a set X and a set $\mathcal{C}$ of subsets of X satisfying three properties:

(i) $\emptyset$ and X are in $\mathcal{C}$,

(ii) $\mathcal{C}$ is closed under intersections, and

(iii) $\mathcal{C}$ is closed under unions of nested families.

A pair $(X, \mathcal{C})$ satisfying all three properties is called a convexity space. These structures were extensively investigated by associating to a general convexity space various parameters such as its Helly number, its Radon number, etc. and establishing relations between these parameters that hold for arbitrary convexity spaces. This includes for instance Levi's inequality between Helly and Radon numbers [22]. Let $\mathcal{F}$ be a set system with ground set X. The $\mathcal{F}$-convex hull $\mathrm{conv}_{\mathcal{F}}(P)$ of a subset $P \subset X$ is the intersection of all the members of $\mathcal{F}$ that contain P. When $\mathcal{F}$ is not a convexity space, we do not have that $\mathrm{conv}_{\mathcal{F}}(P)$ is in $\mathcal{F}$ but this is of no concern to us.

- The Helly number $\mathrm{h}_{\mathcal{F}}$ of $\mathcal{F}$ is the smallest integer h with the following property: If in a finite subfamily $\mathcal{G} \subset \mathcal{F}$, every h members of $\mathcal{G}$ intersect, then $\mathcal{G}$ has nonempty intersection. If no such h exists, we set $\mathrm{h}_{\mathcal{F}} = \infty$.

- The Radon number $\mathrm{r}_{\mathcal{F}}$ of $\mathcal{F}$ is the smallest integer r such that every r-element subset $S \subset X$ can be partitioned into two nonempty parts $S = P_1 \sqcup P_2$ such that $\mathrm{conv}_{\mathcal{F}}(P_1) \cap \mathrm{conv}_{\mathcal{F}}(P_2) \neq \emptyset$. If no such r exists, we set $\mathrm{r}_{\mathcal{F}} = \infty$.

Note that both definitions make sense also without axioms (i)–(iii) of convexity spaces. Here is a proof of Levi's inequality:

Lemma C.1. For every set system $\mathcal{F}$ with finite Radon number, $\mathrm{h}_{\mathcal{F}} \leq \mathrm{r}_{\mathcal{F}} - 1$.

Proof. Let $\mathcal{F}$ be a set system and let X denote its ground set. Let $\mathrm{h}(\mathcal{F})$ denote the set of all subsets $W \subseteq \mathcal{F}$ with the following properties: W has empty intersection and every proper subset of W has nonempty intersection. Let $|\cdot|$ denotes the cardinality. Note that:

- Every $W \in \mathrm{h}(\mathcal{F})$ refutes the possibility that the Helly number of $\mathcal{F}$ be at most $|W| - 1$.

- Conversely, these are the inclusion-minimal obstructions: For any $k \in \mathbb{N}$ and finite subset $\mathcal{G} \subset \mathcal{F}$, if every k members of $\mathcal{G}$ intersect and $\mathcal{G}$ has empty intersection, then there exists $W \subseteq \mathcal{G}$ such that $W \in \mathrm{h}(\mathcal{F})$ and $|W| \geq k + 1$.

It follows that $\mathrm{h}_{\mathcal{F}}$ is finite if and only if $\mathrm{h}(\mathcal{F})$ is finite, and $\mathrm{h}_{\mathcal{F}} = \max_{W \in \mathrm{h}(\mathcal{F})} |W| - 1$. Let $W \in \mathrm{h}(\mathcal{F})$. For every $A \in W$ let $x_A \in \cap_{B \in W \setminus \{A\}} B$. Let $S = \{x_A \colon A \in W\}$. Consider an arbitrary partition $S = P_1 \sqcup P_2$. For $i = 1, 2$ define $W_i = \{A \in W \colon x_A \notin P_i\}$. Observe that $W = W_1 \cup W_2$ and that W_i contains $\{A \in W \colon P_i \subset A\}$. It follows that

$$\begin{aligned}\mathrm{conv}_{\mathcal{F}}(P_1) \cap \mathrm{conv}_{\mathcal{F}}(P_2) &= \left(\bigcap_{\substack{A \in \mathcal{F} \\ P_1 \subset A}} A\right) \cap \left(\bigcap_{\substack{A \in \mathcal{F} \\ P_2 \subset A}} A\right) \\ &\subset \underbrace{\left(\bigcap_{\substack{A \in W \\ P_1 \subset A}} A\right)}_{\subset \bigcap_{A \in W_1} A} \cap \underbrace{\left(\bigcap_{\substack{A \in W \\ P_2 \subset A}} A\right)}_{\subset \bigcap_{A \in W_2} A} \subset \left(\bigcap_{A \in W} A\right) = \emptyset\end{aligned}$$

Thus, there is no partition $S = P_1 \sqcup P_2$ such that $\mathrm{conv}_{\mathcal{F}}(P_1) \cap \mathrm{conv}_{\mathcal{F}}(P_2) \neq \emptyset$. It follows that $\mathrm{r}_{\mathcal{F}} \geq |W|$. Since this holds for every $W \in \mathrm{h}(\mathcal{F})$, we have $\mathrm{r}_{\mathcal{F}} \geq \max_{W \in \mathrm{h}(\mathcal{F})} |W| = \mathrm{h}_{\mathcal{F}} + 1$. □

The proof of our Corollary 6 uses Theorem 1.1 of Holmsen and Lee [17], which is stated for convexity spaces. The proof of [17, Theorem 1.1] uses two ingredients:

1. a colorful Helly theorem for convexity spaces with bounded Radon number [17, Theorem 2.2], where convexity spaces are only used to apply Levi's inequality and an inequality of Jamison [18, Ineq. (6)] on partition numbers, and

2. a lower bound on the size of the largest clique in a uniform hypergraph with a forbidden pattern [16, Theorem 1.2].

This only leaves us with Jamison's inequality to examine. Jamison's inequality is about analogues of Radon numbers for more than 2 parts, but dealing with multisets. Let $\mathcal{F}$ be a set system with ground set X. Given a multiset S of X we write $\bar{S}$ for the set of elements that appear in S, and define $\mathrm{conv}_{\mathcal{F}}(S) := \mathrm{conv}_{\mathcal{F}}(\bar{S})$. A partition of a multiset S into k parts is a union of k multisets $(S_1, S_2, \ldots, S_k)$ such that $\bar{S} = \bar{S}_1 \cup \bar{S}_2 \cup \ldots \cup \bar{S}_k$ and the multiplicity of an element in S equals the sum of its multiplicities in $S_1, S_2, \ldots, S_k$.

- For $k \geq 2$, the kth partition number $\mathrm{r}_{\mathcal{F}}^{(k)}$ of $\mathcal{F}$ is the smallest integer r such that every multiset S of X of size r can be partitioned into k nonempty parts $(P_1, P_2, \ldots, P_k)$ such that $\mathrm{conv}_{\mathcal{F}}(P_1) \cap \mathrm{conv}_{\mathcal{F}}(P_2) \cap \ldots \cap \mathrm{conv}_{\mathcal{F}}(P_k) \neq \emptyset$. If no such r exists, we set $\mathrm{r}_{\mathcal{F}}^{(k)} = \infty$.

In particular $\mathrm{r}_{\mathcal{F}}^{(2)} = \mathrm{r}_{\mathcal{F}}$. Jamison's inequality asserts that when $\mathcal{F}$ is a convexity space, $\mathrm{r}_{\mathcal{F}}^{(k)} \leq \mathrm{r}_{\mathcal{F}}^{(k)^{\log_2 k}}$. It follows from a recursive formula [18, Theorem 1] whose proof uses no property of convexity spaces. For completeness, we reproduce here the proof of a slightly simpler version of that inequality:

Lemma C.2. For any set system $\mathcal{F}$ and any $m, n \in \mathbb{N}$ we have $\mathrm{r}_{\mathcal{F}}^{(mn)} \leq \mathrm{r}_{\mathcal{F}}^{(m)}\, \mathrm{r}_{\mathcal{F}}^{(n)}$.

Proof. Let X be the ground set of $\mathcal{F}$ and let S be a multiset of size $\mathrm{r}_{\mathcal{F}}^{(m)}\, \mathrm{r}_{\mathcal{F}}^{(n)}$ of X. Fix any partition of S into $k = \mathrm{r}_{\mathcal{F}}^{(m)}$ multisets $(S_1, S_2, \ldots, S_k)$, each of size $\mathrm{r}_{\mathcal{F}}^{(n)}$. Each S_i admits a partition $(P_1^i, P_2^i, \ldots, P_n^i)$ such that $\mathrm{conv}_{\mathcal{F}}(P_1^i) \cap \mathrm{conv}_{\mathcal{F}}(P_2^i) \cap \ldots \cap \mathrm{conv}_{\mathcal{F}}(P_n^i) \neq \emptyset$. Let p_i be an element in that intersection. Now let P denote the multiset $\{p_1, p_2, \ldots, p_k\}$. Since $k = \mathrm{r}_{\mathcal{F}}^{(m)}$ there is a partition of P into m multisets $(P_1, P_2, \ldots, P_m)$ and an element $x \in X$ such that $x \in \mathrm{conv}_{\mathcal{F}}(P_1) \cap \mathrm{conv}_{\mathcal{F}}(P_2) \cap \ldots \cap \mathrm{conv}_{\mathcal{F}}(P_m)$. Now, for every $a \,\mathrm{im}[m]$ let $A(a) = \{i \colon p_i \in P_a\}$. For any $(a, b) \in [m] \times [n]$ we put $Q_{a,b} := \cup_{i \in A(a)} P_b^i$. The $Q_{a,b}$ therefore form a partition of S into mn multisets. Moreover, since for all i and b, $p_i \in \mathrm{conv}_{\mathcal{F}}(P_b^i)$, we have that each $\mathrm{conv}_{\mathcal{F}}(Q_{a,b})$ contains every p_i with $i \in A(a)$, and therefore also contains x. □

Finally, we note that the proofs by Alon et al. [1] that a Fractional Helly theorem implies the existence of weak ε-nets, (p, q)-theorems, etc. does require the set system to be closed under intersection.

D An upper bound on the growth of graded Radon numbers

A pigeonhole argument yields that for every set system $\mathcal{F}$ the graded Radon numbers satisfy $\mathrm{r}_{\mathcal{F}}(t) < 2^t$. Indeed, the (graded) Radon numbers do not change when we identify two elements in the ground set of $\mathcal{F}$ that belong to exactly the same members of $\mathcal{F}$. It turns out that a much sharper bound holds.

Proposition D.1. For any set system $\mathcal{F}$ and any $t \in \mathbb{N}$, we have $\mathrm{r}_{\mathcal{F}}(t) \leq t + 1$.

Proof. Suppose there exists $\mathcal{F}' \subseteq \mathcal{F}$ of size t of Radon number greater than $t + 1$, that is with some set $S = \{p_1, \ldots, p_{t+1}\}$ of $t + 1$ points in X with no $\mathcal{F}'$-Radon partition. Observe that if a partition $\mathcal{P} = (\mathcal{P}_0, \mathcal{P}_1)$ of $\{p_1, \ldots, p_{t+1}\}$ is not a Radon partition, then the $\mathcal{F}'$-hull of $\mathcal{P}_0$ does not contain any member of $\mathcal{P}_1$. In particular, there exists some $F_i \in \mathcal{F}'$ that contains $\mathcal{P}_0$ but not $\mathcal{P}_1$. Let us apply this observation to the partitions $\mathcal{P}^{(j)} = (\{p_1, \ldots, p_{t+1}\} \backslash \{p_j\}, \{p_j\})$ for $j \in \{1, \ldots, t\}$. For every $j \in [t]$ there exists $F_{i_j} \in \mathcal{F}'$ such that $F_{i_j} \cap S = S \setminus \{p_j\}$. Note that the $\{F_{i_j}\}_{j \in [t]}$ must be pairwise disjoint, that is $\mathcal{F}' = \{F_{i_j}\}_{j \in [t]}$. To conclude, observe that no member of $\mathcal{F}'$ contains $\{p_1, p_2, \ldots, p_t\}$. We thus have

$$\mathrm{conv}_{\mathcal{F}'}(\{p_1, \ldots, p_t\}) = X,$$

which contains p_{t+1}. Altogether, $(\{p_1, \ldots, p_t\}, \{p_{t+1}\})$ is a Radon partition, which contradicts the initial assumption. □

References

[1] Noga Alon, Gil Kalai, Jirí Matoušek, and Roy Meshulam. Transversal numbers for hypergraphs arising in geometry. Advances in Applied Mathematics, 29:79–101, 2002.

[2] Ervin G. Bajmóczy and Imre Bárány. On a common generalization of borsuk's and radon's theorem. Acta Mathematica Hungarica, 34(3-4):347–350, 1979.

[3] Imre Bárány. Combinatorial convexity, volume 77. American Mathematical Soc., 2021.

[4] Imre Bárány and Pablo Soberón. Tverberg's theorem is 50 years old: a survey. Bulletin of the American Mathematical Society, 55(4):459–492, 2018.

[5] Saugata Basu and Anthony Rizzie. Multi-degree bounds on the betti numbers of real varieties and semi-algebraic sets and applications. Discrete & Computational Geometry, 59:553–620, 2018.

[6] Boris Bukh. Radon partitions in convexity spaces, 2010.

[7] Sourav Chakraborty, Rameshwar Pratap, Sasanka Roy, and Shubhangi Saraf. Helly-type theorems in property testing. International Journal of Computational Geometry & Applications, 28(04):365–379, 2018.

[8] Jesús De Loera, Xavier Goaoc, Frédéric Meunier, and Nabil Mustafa. The discrete yet ubiquitous theorems of carathéodory, helly, sperner, tucker, and tverberg. Bulletin of the American Mathematical Society, 56(3):415–511, 2019.

[9] Éric Colin de Verdière, Vojtěch Kaluža, Pavel Paták, Zuzana Patáková, and Martin Tancer. A direct proof of the strong hanani-tutte theorem on the projective plane. Journal of Graph Algorithms and Applications, 21(5):939–981, 2017.

[10] Jürgen Eckhoff. The partition conjecture. Discrete Mathematics, 221(1):61–78, 2000.

[11] Rogers Epstein and Sandeep Silwal. Property Testing of LP-Type Problems. In Artur Czumaj, Anuj Dawar, and Emanuela Merelli, editors, 47th International Colloquium on Automata, Languages, and Programming (ICALP 2020), volume 168 of Leibniz International Proceedings in Informatics (LIPIcs), pages 98:1–98:18, 2020.

[12] Radoslav Fulek and Jan Kynčl. Counterexample to an extension of the hanani-tutte theorem on the surface of genus 4. Combinatorica, 39(6):1267–1279, 2019.

[13] Xavier Goaoc, Andreas F. Holmsen, and Zuzana Patáková. Intersection patterns in spaces with a forbidden homological minor, 2024.

[14] Xavier Goaoc, Pavel Paták, Zuzana Patáková, Martin Tancer, and Uli Wagner. Bounding Helly numbers via Betti numbers. In A Journey Through Discrete Mathematics: A Tribute to Jiří Matoušek, pages 407–447. Springer, 2017.

[15] Eduard Helly. Über Systeme von abgeschlossenen Mengen mit gemeinschaftlichen Punkten. Monatshefte für Mathematik und Physik, 37:281–302, 1930.

[16] Andreas F. Holmsen. Large cliques in hypergraphs with forbidden substructures. Combinatorica, 40(4):527–537, Aug 2020.

[17] Andreas F. Holmsen and Donggyu Lee. Radon numbers and the fractional Helly theorem. Israel Journal of Mathematics, 241(1):433–447, Mar 2021.

[18] Robert Jamison. Partition numbers for trees and ordered sets. Pacific Journal of Mathematics, 96(1):115–140, 1981.

[19] Gil Kalai. Intersection patterns of convex sets. Israel Journal of Mathematics, 48(2):161–174, 1984.

[20] Gil Kalai. Combinatorial expectations from commutative algebra. In I. Peeva anv V. Welker, editor, Combinatorial Commutative Algebra, volume 1(3), pages 1729–1734. Oberwolfach Reports, 2004.

[21] Meir Katchalski and Andy Liu. A problem of geometry in $\mathbb{R}^n$. Proceedings of the American Mathematical Society, 75(2):284–288, 1979.

[22] Friedrich W. Levi. On Helly's theorem and the axioms of convexity. Journal of the Indian Mathematical Society, 15:65–76, 1951.

[23] Jiři Matoušek. Lectures on discrete geometry, volume 212. Springer Science & Business Media, 2013.

[24] Jirí Matoušek. A Helly-type theorem for unions of convex sets. Discrete & Computational Geometry, 18:1–12, 1997.

[25] Jirí Matoušek, Micha Sharir, and Emo Welzl. A subexponential bound for linear programming. Algorithmica, 16(4-5):498–516, 1996.

[26] Jirí Matoušek. Bounded VC-dimension implies a fractional Helly theorem. Discrete & Computational Geometry, 31:251–255, 2004.

[27] John Milnor. A procedure for killing homotopy groups of differentiable manifolds. In Proceedings of symposia in pure mathematics, pages 39–55. American Mathematical Society, 1961.

[28] Pavel Paták. A sharper ramsey theorem for constrained drawings. Journal of Graph Theory, 109(4):401–411, 2025.

[29] Pavel Paták and Martin Tancer. Embeddings of k-complexes into 2 k-manifolds. Discrete & Computational Geometry, 71(3):960–991, 2024.

[30] Zuzana Patáková. Bounding Radon number via Betti numbers. International Mathematics Research Notices, 2024(11):9482–9500, 2024.

[31] Zuzana Patáková. Bounding radon numbers via betti numbers, 2024.

[32] Michael J Pelsmajer, Marcus Schaefer, and Despina Stasi. Strong hanani–tutte on the projective plane. SIAM Journal on Discrete Mathematics, 23(3):1317–1323, 2009.

[33] Victor V. Prasolov. Elements of homology theory, volume 81. American Mathematical Society, 2007.

[34] Gerhard Ringel. Map color theorem, volume 209. Springer Science & Business Media, 2012.

[35] Colin Rourke and Brian Sanderson. Introduction to Piecewise-Linear Topology. Springer Study Edition. Springer Berlin, Heidelberg, 1972.

[36] Arkadiy Skopenkov. Embeddings of k-complexes in $2k$-manifolds and minimum rank of partial symmetric matrices, 2024.

[37] Martin Tancer. Intersection patterns of convex sets via simplicial complexes: a survey. In Thirty essays on geometric graph theory, pages 521–540. Springer, 2012.

[38] Uli Wagner. Minors, embeddability, and extremal problems for hypergraphs. In Thirty Essays on Geometric Graph Theory, pages 569–607. Springer, 2012.

[39] Eric Christopher Zeeman. Relative simplicial approximation. In Mathematical Proceedings of the Cambridge Philosophical Society, volume 60, pages 39–43. Cambridge University Press, 1964.