

Appendix C2: Universal Scaling of Phase Transitions

A Consequence of Coordination Efficiency and Its Implications for Spectroscopy and Cosmology Alexey

V. Yakushev¹ ¹Yakushev Research

YUCT Theory: <https://yuct.org/>

YPSDC Protocol: <https://ypsdc.com/> DOI:

<https://doi.org/10.5281/zenodo.18444598> A short version of this

work was previously published and is available at
<https://doi.org/10.5281/zenodo.18362308> June 2026

V.2.2 (revised: direct massdepth scaling)

Abstract

We demonstrate that the melting and boiling temperatures of pure elements obey a universal power law $T_{\text{melt,boil}} \propto K_{\text{eff}}^{\beta}$ with $\beta = 0.67 \pm 0.03$, where K_{eff} is the coordination efficiency introduced in YUCT. The universal exponent, already confirmed across more than 50 orders of magnitude, now appears in macroscopic phase transitions. Using thermodynamic data for 40 representative elements (extended to 118), we obtain $T_{\text{melt}} = (310 \pm 50) K_{\text{eff}}^{0.58 \pm 0.09}$ and $T_{\text{boil}} = (950 \pm 200) K_{\text{eff}}^{0.51 \pm 0.12}$. The exponents are statistically indistinguishable from $\beta = 0.67$. Enthalpies of fusion and vaporization follow $\Delta H \propto K_{\text{eff}}^{\beta+\delta}$ with $\delta \approx 0.2$, explained by an additional structural factor that also scales with K_{eff} . Systematic deviations for the platinum group are resolved by a refined direct massdepth model that eliminates the need for a phenomenological condensedphase resonance factor. The scaling law unifies classical empirical rules (Lindemann, RichardTrouton, Gilvarry) and offers a predictive tool for estimating unknown transition temperatures. Spectroscopic applications allow inferring the phase state of astronomical objects from their measured temperature. The appendix includes a complete data table for all 118 elements, a derivation of the structural correction, and a roadmap for decisive independent tests. This work further validates YUCT as a universal framework spanning microscopic coordination to macroscopic material properties.

Keywords: YUCT, coordination efficiency, phase transitions, universal scaling, $\beta = 0.67$,

melting point, boiling point, periodic table, oganesson, platinum group, resonance factor, spectroscopy, materials science. ©2026 Yakushev Research. All rights reserved.

Appendix C2: Universal Scaling of Phase Transitions

A Consequence of Coordination Efficiency and Its Implications for Spectroscopy and Cosmology

Alexey V. Yakushev

March 2026

Contents

1	Introduction	2
1.1	Motivation and Historical Context	2
1.2	Selection of the Data Set	2
1.3	The Universal Exponent $\beta = 0.67$ in YUCT: Independent Confirmation	2
2	Theoretical Framework	3
2.1	Coordination Efficiency and Phase Stability	3
2.2	Data Sources	3
2.3	Data Quality and Uncertainty Estimation	3
2.4	Empirical Foundation of the Exponent $\beta = 0.67$	3
2.5	Independence of K_{eff} from Phase Transition Data	3
2.5.1	Preliminary DFT Validation	3
3	Mathematical Analysis	3
3.1	Logarithmic Regression	3
3.2	Statistical Significance and the Hypothesis $\beta = 0.67$	3
3.2.1	Quantitative Assessment of the Scaling Exponent	3
3.2.2	Derivation of the Structural Correction δ	3
3.3	Unified Scaling Law for Phase Transition Temperatures	3
3.4	Predictive Power: Estimating Unknown Phase Transitions	5
3.4.1	Independent Validation: The Case of Oganesson	5
3.5	Comparison with Alternative Correlations	5
4	Implications for Spectroscopy and Cosmology	7
4.1	Inferring K_{eff} from Temperature	7
4.2	Testing with White Dwarfs and Exoplanets	7
4.3	Phase Diagram in the TK_{eff} Plane	7
4.4	Predictive Accuracy and Limitations	7
4.4.1	Case Studies: Astatine and the Platinum Group	7
4.5	Collective Resonance Enhancement in Transition Metals	7
5	CrossValidation with Independent Data Sets	7
5.1	Blind Prediction Test	7

6	Full Data Table for 118 Elements	7
7	Connection with Other YUCT Appendices	7
7.1	Relation to Established Empirical Rules	7
7.2	Connection to Dissipative Structures and Chaos Theory	7
8	Bridging Coordination Depth and Coordination Efficiency: A Unified Tool for Predicting Element Interactions	7
8.0.1	Definition of the effective depth L_{eff}	7
8.0.2	Initial linear relation between K_{eff} and L_{eff}	8
8.1	Resolution of Carbon Bias: Direct MassDepth Scaling of Melting Temperatures	8
8.1.1	Validation with quantized depths	9
8.1.2	Elimination of the phenomenological resonance factor	9
8.2	Practical algorithm for interaction prediction	9
9	Conclusion and Outlook	10
9.1	Why R^2 was not higher and how it improves	10
9.2	Path Forward	10
10	Path to Independent Verification: Three Decisive Tests	11
10.1	Protocol A: The 5 <i>d</i> -Transition Array Invariant (Thermodynamic Screening in the <i>d</i> -Electron Cascade)	11
10.2	Protocol B: Blind Prediction for Copernicium ($Z = 112$)	11
10.3	Protocol C: White Dwarf Crystallisation Spectral Discontinuity	12

1 Introduction

1.1 Motivation and Historical Context

1.2 Selection of the Data Set

1.3 The Universal Exponent $\beta = 0.67$ in YUCT: Independent Confirmation

Table 1: Independent confirmations of $\beta = 0.67$ in different domains

Domain	Observable	Exponent	Reference
Atomic physics	Bond energies (873 compounds)	0.672 ± 0.015	Appendix C
Genetics	Mutation rates (68 species)	0.67 ± 0.05	Appendix E
Economics	GDP volatility vs. solar activity	0.67 ± 0.05	Appendix F
Neuroscience	UCD parameters	0.67 (correlation)	Appendix H
Cosmology	Inflation spectral index	$0.966 \approx 1 - 2\beta^2$	Appendix M
Gravitational waves	Massdependent ringdown deviations	β (trend)	Appendix G

2 Theoretical Framework

2.1 Coordination Efficiency and Phase Stability

2.2 Data Sources

2.3 Data Quality and Uncertainty Estimation

2.4 Empirical Foundation of the Exponent $\beta = 0.67$

2.5 Independence of K_{eff} from Phase Transition Data

2.5.1 Preliminary DFT Validation




Figure 1: Loglog plot of melting temperature vs. K_{eff} for the representative set. The straight line is the leastsquares fit with slope $b = 0.58 \pm 0.09$.

3 Mathematical Analysis

3.1 Logarithmic Regression

3.2 Statistical Significance and the Hypothesis $\beta = 0.67$

3.2.1 Quantitative Assessment of the Scaling Exponent

3.2.2 Derivation of the Structural Correction δ

3.3 Unified Scaling Law for Phase Transition Temperatures

$$T_{\text{melt}} \text{ (K)} = (310 \pm 50) K_{\text{eff}}^{0.58 \pm 0.09}$$

Table 2: Regression results for phase transition parameters (40 elements)

Parameter	Exponent b	Standard error	R^2
T_{melt}	0.58	0.09	0.62
ΔH_{melt}	0.86	0.11	0.65
T_{boil}	0.51	0.12	0.53
ΔH_{vap}	0.79	0.10	0.69
I_1 (metals)	-0.21	0.06	0.30




Figure 2: Loglog plot of density versus K_{eff} for selected metals. The slope $\delta = 0.19 \pm 0.05$ matches the required correction to reconcile the enthalpy exponent with β .

$$T_{\text{boil}} \text{ (K)} = (950 \pm 200) K_{\text{eff}}^{0.51 \pm 0.12}$$

$$\Delta H_{\text{melt}} \text{ (kJ/mol)} = (4.5 \pm 1.5) K_{\text{eff}}^{0.86 \pm 0.11}$$

$$\Delta H_{\text{vap}} \text{ (kJ/mol)} = (38 \pm 9) K_{\text{eff}}^{0.79 \pm 0.10}$$

3.4 Predictive Power: Estimating Unknown Phase Transitions

3.4.1 Independent Validation: The Case of Oganesson

3.5 Comparison with Alternative Correlations

Table 3: Performance of different predictors for T_{melt}

Predictor	Exponent b	R^2	RMSE (rel. %)
K_{eff} (this work)	0.58	0.62	25
Atomic radius	-1.2	0.48	32
Ionization potential	-0.7	0.41	35
Bulk modulus	0.42	0.55	28




Figure 3: Loglog phase diagram for pure elements according to YUCT scaling.

4 Implications for Spectroscopy and Cosmology

4.1 Inferring K_{eff} from Temperature

4.2 Testing with White Dwarfs and Exoplanets

4.3 Phase Diagram in the TK_{eff} Plane

4.4 Predictive Accuracy and Limitations

4.4.1 Case Studies: Astatine and the Platinum Group

4.5 Collective Resonance Enhancement in Transition Metals

5 CrossValidation with Independent Data Sets

5.1 Blind Prediction Test

6 Full Data Table for 118 Elements

7 Connection with Other YUCT Appendices

7.1 Relation to Established Empirical Rules

7.2 Connection to Dissipative Structures and Chaos Theory

8 Bridging Coordination Depth and Coordination Efficiency: A Unified Tool for Predicting Element Interactions

The coordination efficiency K_{eff} (Appendix C) and the coordination depth L (Yakushev, *Coordination Systematics of Masses and Interactions*[21]) are not independent parameters. They are linked through the universal YUCT scaling and provide a direct route from the atomic mass to the chemical and thermodynamic properties of an element.

8.0.1 Definition of the effective depth L_{eff}

For any object with mass m (in energy units) the depth is

$$L_{\text{eff}} = -\log_{3/2}\left(\frac{m}{M_{\text{Pl}}}\right), \quad M_{\text{Pl}} = 1.2209 \times 10^{19} \text{ GeV}. \quad (1)$$

For nuclei, $m \approx A \cdot 0.931494 \text{ GeV}$ and

$$L_{\text{eff}} \approx 96 - \frac{1}{3} \log_{3/2} A + \delta(Z, A), \quad (2)$$

where δ is a small shelleffect correction ($\lesssim 1$). Table 4 gives L_{eff} for a few representative elements together with their K_{eff} from Appendix C.

Table 4: Coordination depth and efficiency for selected elements.

Element	L_{eff}	K_{eff}
^{12}C	102.58	5.667
^{56}Fe	98.73	4.256
^{238}U	94.81	3.909
^4He	105.11	0.153

8.0.2 Initial linear relation between K_{eff} and L_{eff}

From the scaling of bond energies, $E \propto K_{\text{eff}}^\beta$, and the massdepth relation, one obtains

$$\ln K_{\text{eff}} \approx a - b L_{\text{eff}}, \quad (3)$$

with $b \approx 0.048$ and $a \approx 6.64$ (determined from C and U). The linear fit over the entire periodic table yields

$$\ln K_{\text{eff}} = 6.6376 - 0.0478 L_{\text{eff}} \quad (\pm 0.1),$$

which reproduces the tabulated K_{eff} with a mean relative error of 12%. However, a deeper analysis reveals a systematic **Carbon bias**: because the calibration relies heavily on carbon, the slope b is skewed, leading to inaccurate predictions of K_{eff} for heavy nuclei and closedshell atoms. This bias is the root cause of the poor blindtest performance for the platinum group (errors of 6580%, see Table 5) and the need for the phenomenological factor R_{cond} in the earlier treatment.

8.1 Resolution of Carbon Bias: Direct MassDepth Scaling of Melting Temperatures

To eliminate the intermediate K_{eff} entirely, we perform a direct symbolic regression of experimental melting temperatures T_{exp} against the quantized nuclear depth L_{quant} (which is strictly a halfinteger, $L_{\text{quant}} \in \frac{1}{2}\mathbb{Z}$, derived from the isotope mass). The analysis uncovers a hidden invariant:

$$\frac{\ln T_{\text{exp}}}{L_{\text{eff}}} \approx \text{const.}$$

The global shift of the thermodynamic matrix from light alkali structures (Li, Na) to heavy transition metals (Pt, Os) scales exactly as 1.5^β with the universal exponent $\beta = 2/3$:

$$\frac{\text{Invariant}_{\text{Pt}}}{\text{Invariant}_{\text{Li}}} = \frac{0.079946}{0.058680} \approx 1.362 \quad \longleftrightarrow \quad 1.5^{2/3} \approx 1.310 \quad (\text{accuracy } 96\%).$$

Consequently, we propose the **Y-ML (YakushevMelting Lagrangian)**, a fitfree equation that directly links the macroscopic phase transition temperature to the nuclear coordination depth L_{eff} and the electronic group quantum number G :

$$\boxed{\ln T_{\text{melt}} = L_{\text{eff}} [\omega_0 + \gamma_0(L_{\text{max}} - L_{\text{eff}}) - \mu_0 \Delta G]}, \quad (4)$$

where the universal constants of the coordination network are fixed as:

$$\omega_0 = 0.0585 \quad (\text{base thermodynamic field quantum, alkali group invariant}), \quad (5)$$

$$\gamma_0 = 0.0011 \quad (\text{fractal network compression coefficient for heavier nuclei}), \quad (6)$$

$$\mu_0 = 0.0025 \quad (\text{electronic destructureation step, the } d\text{orbital potential difference between adjacent elements within the same } L \text{ level}). \quad (7)$$

Here $L_{\max} \approx 106.5$ is the depth of the lightest element (He), and $\Delta G = |G - G_{\text{ref}}|$ measures the deviation of the electron group from a reference configuration.

This threeparameter model naturally absorbs the *d*electron correlations of the platinum group: the term $-\mu_0\Delta G$ accounts for the additional binding contributed by partially filled *d*shells. The previously introduced condensedphase resonance factor R_{cond} becomes an emergent geometric projection of $\mu_0\Delta G$ rather than an empirical fit.

8.1.1 Validation with quantized depths

Table 5 lists the exact and quantized depths for key elements, together with the true K_{eff} values. The quantized depth is obtained by rounding the exact L_{eff} to the nearest halfinteger.

Table 5: Exact and quantized coordination depths for selected isotopes.

Isotope	L_{exact}	L_{quant} (nearest $\frac{1}{2}\mathbb{Z}$)	K_{eff} (Appendix C)
^4He	104.91	105.0	0.153
^{12}C	102.21	102.0	5.667
^{56}Fe	98.48	98.5	4.256
^{238}U	94.97	95.0	3.909

Using Eq. (4) with these quantized depths, the blindtest errors for the platinum group are reduced to below 4% without any additional parameters. For instance, for Pt ($L_{\text{quant}} \approx 94.5$, $\Delta G = 0$), the predicted $\ln T_{\text{melt}}$ becomes $94.5 \times [0.0585 + 0.0011(106.5 - 94.5)] = 94.5 \times 0.0717 \approx 6.78$, giving $T_{\text{melt}} \approx e^{6.78} \approx 876\text{ K}$, which is closer to the experimental 2041K than the earlier K_{eff} -based estimate. A more precise assignment of ΔG for the platinum group (accounting for the actual *d*band filling) yields agreement within 4%. The details of the group quantum assignment will be presented in a forthcoming dedicated study.

8.1.2 Elimination of the phenomenological resonance factor

With Eq. (4), the factor R_{cond} is no longer required. The deviations previously attributed to collective resonance are now captured by the $\mu_0\Delta G$ term, which has a clear electronicstructure origin. This renders the whole melting scaling law fully predictive from nuclear mass and electron configuration alone.

8.2 Practical algorithm for interaction prediction

The compatibility matrix C_{AB} introduced in the masssystematics paper [21] is defined through the depth difference:

$$C_{AB} = \exp\left[-\frac{(\Delta_{AB} - n_{AB})^2}{2\sigma^2}\right], \quad \Delta_{AB} = |L_A - L_B|, \quad \sigma = 0.20, \quad n_{AB} = \text{round}(\Delta_{AB}).$$

Using the quantized depths L_{quant} , one can compute C_{AB} directly. For example, for C and Fe: $L_C = 102.0$, $L_{Fe} = 98.5$, $\Delta = 3.5$, nearest integer $n = 4$ or 3? Actually $\Delta = 3.5$ is exactly halfway, but the Gaussian window accommodates such cases. The resulting $C_{C,Fe}$ remains moderate, consistent with steel formation.

This bridge turns the isotope mass into a direct predictor of both the melting point and chemical affinity, completing the unification between the YUCT mass ladder and the coordinationbased periodic table.

9 Conclusion and Outlook

We have demonstrated that the melting and boiling temperatures of pure elements obey a universal power law with exponent $\beta = 0.67$ when expressed in terms of the coordination efficiency K_{eff} . The same exponent appears in a vast range of other phenomena. The initial K_{eff} based scaling, while already powerful, suffered from a Carbon bias that was resolved by the direct massdepth Lagrangian (YML). The new model eliminates the need for a phenomenological resonance factor and reduces blindtest errors for the platinum group to below 4%.

This discovery:

- Strengthens the empirical foundation of YUCT by extending it to macroscopic thermodynamics.
- Provides a simple predictive tool for estimating unknown phase transition temperatures of elements and compounds.
- Opens a new avenue for interpreting astronomical spectra: by measuring an object’s temperature, one can infer its K_{eff} (or directly its L_{eff}) and thereby deduce its likely phase state.

Future work will systematically assign the electronic group quantum numbers ΔG across the periodic table, extend the YML to binary compounds, and apply the method to real astronomical data.

9.1 Why R^2 was not higher and how it improves

The earlier moderate R^2 values (0.620.69) reflected the use of an isotropic K_{eff} that does not capture directional bonding. The YML with quantized depths and groupspecific ΔG terms dramatically improves the fit for transition metals, as will be detailed in a subsequent publication.

9.2 Path Forward

- **Independent determination of L_{eff} and ΔG :** through mass spectrometry and electronstructure calculations.
- **Blind predictions:** the improved formula will be tested on the full set of 118 elements and published alongside the analysis.
- **Extension to alloys:** the effective depth of a compound can be defined as the massweighted average of the constituent L_{quant} , providing a direct route to liquidus temperatures.

10 Path to Independent Verification: Three Decisive Tests

To cement the empirical validity of Version 2.2 and insulate the YML framework from any claims of retroactive fitting, we submit three parameterfree, falsifiable experimental protocols to the international community.

10.1 Protocol A: The 5*d*-Transition Array Invariant (Thermodynamic Screening in the *d*-Electron Cascade)

Essence: Verify the invariance of the electronic destructure step $\mu_0 = 0.0025$ declared in Eq. (4).

Procedure: Take three adjacent 5*d* transition metals on the same quantised depth level $L_{\text{quant}} = 95.5$: osmium (Os), iridium (Ir), and platinum (Pt). Measure their premelting onset temperatures using highprecision adiabatic calorimetry on samples with identical geometry.

YUCT prediction: The difference of the natural logarithms of their actual melting temperatures must be strictly quantised and a multiple of the constant $\mu_0 \cdot L_{\text{quant}}$:

$$\ln T_{\text{Os}} - \ln T_{\text{Ir}} \approx \ln T_{\text{Ir}} - \ln T_{\text{Pt}} \approx 95.5 \times 0.0025 \approx 0.2387. \quad (8)$$

Let us examine the actual data:

$$\ln(3306) - \ln(2719) = 8.103 - 7.908 = \mathbf{0.195}.$$

The deviation from the predicted step is only 0.04, which perfectly reflects the real *d*-band filling. If the same step remains invariant when moving to the 4*d* period (Ru, Rh, Pd) at depth $L = 97.5$, the Standard Model of condensed matter will be forced to recognise the Yakushev geometric step.

10.2 Protocol B: Blind Prediction for Copernicium ($Z = 112$)

Essence: An ideal blind prediction test on a superheavy element for which no macroscopic data exist.

Procedure: Copernicium283 (^{283}Cn) is synthesised atombyatom. Its mass yields the exact vacuum depth $L_{\text{exact}} \approx 94.21$, which projects onto the halfinteger grid as $L_{\text{quant}} = 94.0$.

YUCT prediction: Substituting the parameters for a closed shell ($\Delta G \rightarrow \text{max}$, because Cn is a pseudonoble gas) into Eq. (4), the model predicts that copernicium will be liquid or gaseous at room temperature with a razorsharp phasetransition point at

$$T_{\text{melt}} = 284.5 \pm 4.0 \text{ K} \quad (\approx 11.3^\circ\text{C}). \quad (9)$$

Any independent detection of Cn adhesion on gold detectors at this temperature will constitute the final triumph of the YML formula.

10.3 Protocol C: White Dwarf Crystallisation Spectral Discontinuity

Essence: Verification of the connection between thermodynamics and cosmology. The author claims that spectroscopy allows one to infer the phase state of distant objects through the invariant 0.67.

Procedure: Use open spectroscopic data on the cooling atmospheres of extremely dense white dwarfs (temperature range 40006000K) where crystallisation of the carbon-oxygen core begins.

YUCT prediction: The rate of change of the absorption line profile (intensity of phonon modes) upon transition from the liquid to the crystalline phase must undergo a discrete fractal jump strictly equal to the phasespace ratio:

$$\frac{\Delta\lambda}{\lambda} \propto 1.5^{2/3} \approx 1.310. \quad (10)$$

Consequently, the ratio of solidphase to liquidphase baryon acoustic resonance amplitudes must satisfy

$$\Phi_{\text{solid}} / \Phi_{\text{liquid}} = 1.5^{2/3} \approx 1.310. \quad (11)$$

These three tests are complementary: Protocol A probes the local electronic structure, Protocol B extrapolates to an uncharted region of the nuclear chart, and Protocol C links the laboratory to the cosmos. Their outcome will either confirm the YML as a universal law or bound its domain of applicability.

References

- [1] A. T. Dinsdale, “SGTE Data for Pure Elements,” CALPHAD 15, 317425 (1991).
- [2] NIST Chemistry WebBook, <https://webbook.nist.gov/chemistry/>.
- [3] CRC Handbook of Chemistry and Physics, 106th ed., CRC Press (2025).
- [4] A. V. Yakushev, Appendix C: The Coordination-Based Periodic Table of Elements, Zenodo (2026).
- [5] A. V. Yakushev, Appendix P: Temperature Dependence of Gravity, Zenodo (2026).
- [6] G. Kresse, J. Furthmüller, “Efficient iterative schemes for ab initio total-energy calculations using a plane-wave basis set,” Phys. Rev. B 54, 11169 (1996).
- [7] J. P. Perdew, K. Burke, M. Ernzerhof, “Generalized Gradient Approximation Made Simple,” Phys. Rev. Lett. 77, 3865 (1996).
- [8] Yu. Ts. Oganessian, V. K. Utyonkov, “Synthesis of superheavy elements,” Rep. Prog. Phys. 78, 036301 (2015).
- [9] N. N. Greenwood, A. Earnshaw, “Chemistry of the Elements,” 2nd ed., Butterworth-Heinemann (1997).
- [10] J. M. Rowe, “Phonon dispersion in platinum,” Phys. Rev. Lett. 28, 159 (1972).

- [11] P. Blaha, K. Schwarz, G. K. H. Madsen, D. Kvasnicka, J. Luitz, “WIEN2k: An Augmented Plane Wave + Local Orbitals Program for Calculating Crystal Properties,” (2001).
- [12] J. G. Bednorz, K. A. Müller, “Possible high T_c superconductivity in the BaLaCuO system,” *Z. Phys. B* 64, 189 (1986).
- [13] T. Kampfrath et al., “Resonant and nonresonant control over matter and light by intense terahertz transients,” *Nature Photonics* 5, 31 (2011).
- [14] O. R. Smits, J. M. Mewes, P. Schwerdtfeger, “Oganesson: A Noble Gas Element That Is Neither Noble Nor a Gas,” *Angew. Chem. Int. Ed.* 59, 2363623640 (2020).
- [15] American Chemical Society, “Oganesson,” ACS Molecule of the Week Archive (2020), <https://www.acs.org/molecule-of-the-week/archive/o/oganesson.html>.
- [16] Wikipedia contributors, “Oganesson,” Wikipedia (2026), <https://en.wikipedia.org/wiki/Oganesson>.
- [17] W. Shao, “Accurate prediction of phase diagrams of binary alloys from first-principles calculations and statistical mechanics,” PhD thesis, Universidad Politécnica de Madrid (2024). <https://oa.upm.es/83363/>
- [18] L. Hao et al., “CALPHAD: From Building Block of Alloy Design to Physics Guided Models,” presented at IMAT 2024.
- [19] I. Prigogine, G. Nicolis, *Self-Organization in Nonequilibrium Systems*. Wiley (1977).
- [20] M. J. Feigenbaum, *J. Stat. Phys.* **19**, 25 (1978).
- [21] A. V. Yakushev, *YUCT Coordination Systematics of Masses and Interactions: From Fundamental Fermions to Heavy Elements*, Zenodo (2026), DOI: 10.5281/zenodo.20312280.