

Origin of the energy band structure

in periodic lattices; “molecular orbital” and “electron wave diffraction” reasoning.

July 1, 2021

Origin of the band gaps

- ✚ Electrons in the crystal are arranged in energy band gaps.

Origin of the band gaps

- ✦ Electrons in the crystal are arranged in energy band gaps.
- ✦ The free electron wavefunctions are of the form:

$$\psi(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}}$$

they represent running waves and carry momentum $\mathbf{p} = \hbar\mathbf{k}$.

Origin of the band gaps

- ✱ Electrons in the crystal are arranged in energy band gaps.

- ✱ The free electron wavefunctions are of the form:

$$\psi(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}}$$

they represent running waves and carry momentum $\mathbf{p} = \hbar\mathbf{k}$.

- ✱ The band structure of a crystal can often be explained by the nearly free electron model for which the band electrons are treated as perturbed only weakly by the periodic potential of the ion cores. This model answers almost all the qualitative questions about the behavior of electrons in metals.

Origin of the band gaps

- ✦ Electrons in the crystal are arranged in energy band gaps.


- ✦ The free electron wavefunctions are of the form:

$$\psi(\mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{r}}$$

they represent running waves and carry momentum $\mathbf{p} = \hbar\mathbf{k}$.

- ✦ The band structure of a crystal can often be explained by the nearly free electron model for which the band electrons are treated as perturbed only weakly by the periodic potential of the ion cores. This model answers almost all the qualitative questions about the behavior of electrons in metals.

- ✦ So the energy bands are given from the periodicity



Origin of the band gaps: "electron wave diffraction" reasoning

- ✦ We know that Bragg reflection is a characteristic feature of wave propagation in crystals. The energy bands can also be explained by Bragg reflection of electron waves in crystals.

Origin of the band gaps: "electron wave diffraction" reasoning

- * We know that Bragg reflection is a characteristic feature of wave propagation in crystals. The energy bands can also be explained by Bragg reflection of electron waves in crystals.
- * The Bragg condition $(k + G)^2 = k^2$ for diffraction of a wave of wavevector k becomes in one dimension when $k = \pm \frac{G}{2} = \pm n \frac{\pi}{2}$

Origin of the band gaps: "electron wave diffraction" reasoning

- * We know that Bragg reflection is a characteristic feature of wave propagation in crystals. The energy bands can also be explained by Bragg reflection of electron waves in crystals.
- * The Bragg condition $(k + G)^2 = k^2$ for diffraction of a wave of wavevector k becomes in one dimension when $k = \pm \frac{G}{2} = \pm n \frac{\pi}{2}$
- * At the edge of the first Brillouin zone band gap occurs.

Origin of the band gaps: "electron wave diffraction" reasoning

- * We know that Bragg reflection is a characteristic feature of wave propagation in crystals. The energy bands can also be explained by Bragg reflection of electron waves in crystals.
- * The Bragg condition $(k + G)^2 = k^2$ for diffraction of a wave of wavevector k becomes in one dimension when $k = \pm \frac{G}{2} = \pm n \frac{\pi}{a}$
- * At the edge of the first Brillouin zone band gap occurs.
- * The wavefunctions at $k = \pm \frac{\pi}{a}$ are not the traveling waves $\exp(i \frac{\pi x}{a})$ or $\exp(-i \frac{\pi x}{a})$ of free electrons. These are standing waves.

Origin of the band gaps: "electron wave diffraction" reasoning

- ✱ Hence, the time-independent state is represented by standing waves. We can form two different standing waves from the two traveling waves:

$$\begin{aligned}\psi(+) &= \exp(i\pi x/a) + \exp(-i\pi x/a) = 2 \cos(\pi x/a) \\ \psi(-) &= \exp(i\pi x/a) - \exp(-i\pi x/a) = 2i \sin(\pi x/a)\end{aligned}\tag{1}$$

Origin of the band gaps: "electron wave diffraction" reasoning

- Hence, the time-independent state is represented by standing waves. We can form two different standing waves from the two traveling waves:

$$\begin{aligned}\psi(+) &= \exp(i\pi x/a) + \exp(-i\pi x/a) = 2 \cos(\pi x/a) \\ \psi(-) &= \exp(i\pi x/a) - \exp(-i\pi x/a) = 2i \sin(\pi x/a)\end{aligned}\quad (1)$$

- The two standing waves $\psi(+)$ and $\psi(-)$ pile up electrons at different regions, and therefore the two waves have different values of the potential energy in the field of the ions of the lattice. This is the origin of the energy gap




Figure 1: The probability density on y axis is $\rho = \psi^* \psi = |\psi|^2$

Origin of the band gaps: "electron wave diffraction" reasoning

- ✱ Regarding the wave $\psi(+)$, this function piles up electrons (negative charge) on the positive ions, where the potential energy is lowest. The potential energy of an electron in the field of a positive ion is negative, so that the force between them is attractive.

Origin of the band gaps: "electron wave diffraction" reasoning

- ✱ Regarding the wave $\psi(+)$, this function piles up electrons (negative charge) on the positive ions, where the potential energy is lowest. The potential energy of an electron in the field of a positive ion is negative, so that the force between them is attractive.
- ✱ On the other hand the standing wave $\psi(-)$, which concentrates electrons away from the ion cores.

Origin of the band gaps: "electron wave diffraction" reasoning

- ✱ Regarding the wave $\psi(+)$, this function piles up electrons (negative charge) on the positive ions, where the potential energy is lowest. The potential energy of an electron in the field of a positive ion is negative, so that the force between them is attractive.
- ✱ On the other hand the standing wave $\psi(-)$, which concentrates electrons away from the ion cores.
- ✱ When we calculate the average or expectation values of the potential energy over these three charge distributions, we find that the potential energy of $\rho(+)$ is lower than that of the traveling wave (because you put in the same point $+$ and $-$ charges that cancel out each others), whereas the potential energy of $\rho(-)$ is higher than the traveling wave.

Origin of the band gaps: "electron wave diffraction" reasoning

- ✱ Regarding the wave $\psi(+)$, this function piles up electrons (negative charge) on the positive ions, where the potential energy is lowest. The potential energy of an electron in the field of a positive ion is negative, so that the force between them is attractive.
- ✱ On the other hand the standing wave $\psi(-)$, which concentrates electrons away from the ion cores.
- ✱ When we calculate the average or expectation values of the potential energy over these three charge distributions, we find that the potential energy of $\rho(+)$ is lower than that of the traveling wave (because you put in the same point $+$ and $-$ charges that cancel out each others), whereas the potential energy of $\rho(-)$ is higher than the traveling wave.
- ✱ We have an energy gap of width E_g if the energies of $\rho(+)$ and $\rho(-)$ differ by E_g

Origin of the band gaps: "electron wave diffraction" reasoning




Figure 2: Energy band gaps

Origin of the band gaps: Molecular orbital reasoning

- ✦ Another approach to describe the presence of energy bands and gaps in the energy structure of a crystal is the Molecular orbital description.

Origin of the band gaps: Molecular orbital reasoning

- ✦ Another approach to describe the presence of energy bands and gaps in the energy structure of a crystal is the Molecular orbital description.
- ✦ Thanks to this we can explain why some materials are good conductors and others are good insulators.

Origin of the band gaps: Molecular orbital reasoning

- ✱ Another approach to describe the presence of energy bands and gaps in the energy structure of a crystal is the Molecular orbital description.
- ✱ Thanks to this we can explain why some materials are good conductors and others are good insulators.
- ✱ We first review the argument used to explain the energy structure of a covalent bond. Consider two identical hydrogen atoms so far apart that there is no interaction whatsoever between them. Further suppose that the electron in each atom is in the same ground state: a $1s$ electron with an energy of -13.6 eV (ignore spin).

Origin of the band gaps: Molecular orbital reasoning

- * Another approach to describe the presence of energy bands and gaps in the energy structure of a crystal is the Molecular orbital description.
- * Thanks to this we can explain why some materials are good conductors and others are good insulators.
- * We first review the argument used to explain the energy structure of a covalent bond. Consider two identical hydrogen atoms so far apart that there is no interaction whatsoever between them. Further suppose that the electron in each atom is in the same ground state: a $1s$ electron with an energy of -13.6 eV (ignore spin).
- * When the hydrogen atoms are brought closer together, the individual wave functions of the electrons overlap and, by the exclusion principle, can no longer be in the same quantum state, which splits the original equivalent energy levels into two different energy levels.

Origin of the band gaps: Molecular orbital reasoning

- * $\langle + - \rangle$ If four hydrogen atoms are brought together, four levels are formed from the four possible symmetries—a single sine wave “hump” in each well, alternating up and down, and so on.

Origin of the band gaps: Molecular orbital reasoning

- * $\langle + - \rangle$ If four hydrogen atoms are brought together, four levels are formed from the four possible symmetries—a single sine wave “hump” in each well, alternating up and down, and so on.




Figure 3: The dependence of energy-level splitting on the average distance between (a) two atoms, (b) four atoms, and (c) a large number of atoms. For a large number of electrons, a continuous band of energies is produced.

Origin of the band gaps: Molecular orbital reasoning

- ✱ In the limit of a very large number N of atoms, we expect a spread of nearly continuous bands of electronic energy levels in a solid. Each of these bands is known as an energy band.

Origin of the band gaps: Molecular orbital reasoning

- ✱ In the limit of a very large number N of atoms, we expect a spread of nearly continuous bands of electronic energy levels in a solid. Each of these bands is known as an energy band.
- ✱ Energy bands differ in the number of electrons they hold. In the 1s and 2s energy bands, each energy level holds up to two electrons (spin up and spin down), so this band has a maximum occupancy of $2N$ electrons. In the 2p energy band, each energy level holds up to six electrons, so this band has a maximum occupancy of $6N$ electrons

Origin of the band gaps: Molecular orbital reasoning

- ✱ In the limit of a very large number N of atoms, we expect a spread of nearly continuous bands of electronic energy levels in a solid. Each of these bands is known as an energy band.
- ✱ Energy bands differ in the number of electrons they hold. In the $1s$ and $2s$ energy bands, each energy level holds up to two electrons (spin up and spin down), so this band has a maximum occupancy of $2N$ electrons. In the $2p$ energy band, each energy level holds up to six electrons, so this band has a maximum occupancy of $6N$ electrons.





Figure 4: The allowed states of energy and wave number are still technically quantized, but for large numbers of atoms, these states are so close together that they are considered to be continuous or “in the continuum.”

Origin of the band gaps: Molecular orbital reasoning



Periodic Potential, Kronig-Penney model

- ✦ A very representative and interesting quantum mechanical model that shows how an electron can move inside a periodic crystal is the Kronig-Penney model.




Figure 5: Kronig-Penney model with a periodic potential consisting of delta functions with magnitude V_0 with periodicity a .

Origin of the band gaps: Periodic Potential and Kronig-Penney model

- ✱ The Kronig-Penney model is an idealized quantum-mechanical system that describes electron motion in a period array of delta functions with magnitude V_0 :

$$U(x) = U_0 \sum_{n=-\infty}^{+\infty} \delta(x - na)$$

where the delta is the dirac delta and a is the lattice constant. The time-independent Schrödinger equation consists:

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + U(x) \right) \psi(x) = E\psi(x) \quad (2)$$

Origin of the band gaps: Periodic Potential and Kronig-Penney model

- ✱ The Kronig-Penney model is an idealized quantum-mechanical system that describes electron motion in a period array of delta functions with magnitude V_0 :

$$U(x) = U_0 \sum_{n=-\infty}^{+\infty} \delta(x - na)$$

where the delta is the dirac delta and a is the lattice constant. The time-independent Schrödinger equation consists:

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + U(x) \right) \psi(x) = E\psi(x) \quad (2)$$

- ✱ According to Bloch's theorem, the solution of the Schrödinger equation when the potential is periodic, can be written as:

$$\psi(x) = e^{iqx} u(x) \quad \text{with } q = n \frac{2\pi}{Na}$$

where $u(x)$ is a periodic function. In fact $u(x) = u(x + a)$.

Origin of the band gaps: Periodic Potential and Kronig-Penney model

- ✳ Then imposing the boundary condition for the wave function solution we a transcendental equation:

$$\cos(qa) = \frac{2mV_0a}{\hbar^2} \frac{\sin(ka)}{ka} + \cos(ka) \quad (3)$$

Origin of the band gaps: Periodic Potential and Kronig-Penney model

- ✦ Then imposing the boundary condition for the wave function solution we a transcendental equation:

$$\cos(qa) = \frac{2mV_0a}{\hbar^2} \frac{\sin(ka)}{ka} + \cos(ka) \quad (3)$$

- ✦ On the left we have a term that depend only on the lattice, on the other side of the equation we have terms that depends only on the energy, through the factor k . This equation is satisfied only for specific value of k and q .


Origin of the band gaps: Periodic Potential and Kronig-Penney model

- ✱ Then imposing the boundary condition for the wave function solution we a transcendental equation:

$$\cos(qa) = \frac{2mV_0a}{\hbar^2} \frac{\sin(ka)}{ka} + \cos(ka) \quad (3)$$

- ✱ On the left we have a term that depend only on the lattice, on the other side of the equation we have terms that depends only on the energy, through the factor k . This equation is satisfied only for specific value of k and q .
- ✱ The allowed values of the energy ϵ are given by those satisfying $qa = \left(\frac{2m\epsilon}{\hbar^2}\right)^{\frac{1}{2}} a$, since the number N of lattice point is big the spectrum of the possible solution is very dense for which that lie under the condition:
$$-1 \leq \frac{2mV_0a}{\hbar^2} \frac{\sin(ka)}{ka} + \cos(ka) \leq 1 .$$

Origin of the band gaps: Periodic Potential and Kronig-Penney model



Origin of the band gaps: Periodic Potential and Kronig-Penney model

