

## \*pn junction

balance between diffusion and drift currents for electrons and holes;  
application of the forward or reverse external bias;

---

July 1, 2021




Figure 1: Doping of a Si-semiconductor


One region (P) is doped with  $B$  atoms (3 valence electrons)

The other region (N) is doped with  $P$  atoms (5 valence electrons)

The electrons in excess in the N region go to the P region leaving an excess of positive charge ( $P^*$ ) apporporting negative charge in the  $P$  region.


# Diffusion process

- ✦ Particles will diffuse to equalize distribution.




# Diffusion process

- ✱ Particles will diffuse to equalize distribution.
- ✱ Particles diffuse from regions with higher concentration to lower and vice versa.



# Diffusion process

- \* Particles will diffuse to equalize distribution.
- \* Particles diffuse from regions with higher concentration to lower and vice versa.
- \* This is simply due to statistical properties.



An electric field starts forming and slows down the process of diffusion until the electrons have not enough energy to win the potential




Figure 2: PN junction

- ✦ The diffusion current consists in electrons moving from the  $N$  region to the  $P$  region (concentration gradient)

$$J_n = J_{n, \text{ drift}} + J_{n, \text{ diff}} = qn\mu_n\varepsilon + qD_n \frac{dn}{dx}$$
$$J_p = J_{p, \text{ drift}} + J_{p, \text{ diff}} = qp\mu_p\varepsilon - qD_p \frac{dp}{dx}$$

At equilibrium  $J = J_n + J_p = 0$


- \* The diffusion current consists in electrons moving from the  $N$  region to the  $P$  region (concentration gradient)
- \* The drift current is due to moving electrons as a consequence of the electric field and it is in the opposite direction of the diffusion current

$$J_n = J_{n, \text{ drift}} + J_{n, \text{ diff}} = qn\mu_n\varepsilon + qD_n \frac{dn}{dx}$$
$$J_p = J_{p, \text{ drift}} + J_{p, \text{ diff}} = qp\mu_p\varepsilon - qD_p \frac{dp}{dx}$$


At equilibrium  $J = J_n + J_p = 0$



$$J_{n,\text{diff}} = qD_n \frac{dn}{dx}$$



$$J_{p,\text{diff}} = -qD_p \frac{dp}{dx}$$



- ✱ When applying an electric field electrons will drift. The velocity will be proportional to the electric field with electron mobility  $\mu_h$  and  $\mu_e$ .

- ✱ When applying an electric field electrons will drift. The velocity will be proportional to the electric field with electron mobility  $\mu_h$  and  $\mu_e$ .
- ✱ They are linked to the velocity and the electric field in the following way:

$$\begin{aligned}\vec{v}_h &= \mu_p \vec{E} \\ \vec{v}_e &= -\mu_n \vec{E}\end{aligned}\tag{1}$$

- ✱ When applying an electric field electrons will drift. The velocity will be proportional to the electric field with electron mobility  $\mu_h$  and  $\mu_e$ .
- ✱ They are linked to the velocity and the electric field in the following way:


$$\begin{aligned}\vec{v}_h &= \mu_p \vec{E} \\ \vec{v}_e &= -\mu_n \vec{E}\end{aligned}\tag{1}$$

- ✱ The total charge flow is determined by a combination of drift and diffusion

$$\begin{aligned}J &= J_n + J_p \\ J_n &= J_{n,\text{drift}} + J_{n,\text{diff}} = qn\mu_n\varepsilon + qD_n \frac{dn}{dx} \\ J_p &= J_{p,\text{drift}} + J_{p,\text{diff}} = qp\mu_p\varepsilon + qD_p \frac{dp}{dx}\end{aligned}$$


# Contact bending

Time = 0, the contact is “just” established




## Band bending

At equilibrium the Fermi energy of the two parts are equal and the other levels bend in order to connect each other



**Figure 7.3** | Energy-band diagram of a pn junction in thermal equilibrium.

**Figure 3:** Energy levels in a PN junction





## Forward bias

Copyright © Physics and Radio-Electronics, All rights reserved



**Figure 4:** Forward biasing of a PN junction

# Biasing


## Forward bias



## Reverse bias








Energy band diagram of a p-n junction under reverse and forward bias

**Figure 5:** Biasing of a PN junction



**Figure 6:** I-V curve of a diode. The exponential trend is due to the Boltzmann distribution. The breakdown occurs when we provide enough energy to break bonds forming electron-hole pairs that travel according to the applied field and counter the depletion region field.