

## STA457TUT5

For an AR(2) process, we have that:

$$x_t = \phi_1 x_{t-1} + \phi_2 x_{t-2} + w_t$$

In our case,  $\phi_1 = 1.5$ ,  $\phi_2 = -0.75$ .

The difference equation for the ACF of the process is:

$$\rho(h) - \phi_1 \rho(h-1) - \phi_2 \rho(h-2) = 0$$

.

So we have a difference equation:

$$\rho(h) - 1.5\rho(h-1) + 0.75\rho(h-2) = 0$$

Solve for general expression of  $\rho(h)$ . Let  $z_1$  and  $z_2$  be the roots of the associated polynomial:

$$\phi(z) = 1 - \phi_1 z - \phi_2 z^2$$

$$\phi(z) = 1 - 1.5z + 0.75z^2$$

where  $z_1 = 1 + i\sqrt{3}/3$ ,  $z_2 = 1 - i\sqrt{3}/3$ .

The general solutions are:

$$\rho(h) = c_1 z_1^{-h} + c_2 z_2^{-h}$$

For initial conditions,

$$\rho(0) = c_1 + c_2 = 1$$

$$\rho(-1) = c_1 z_1 + c_2 z_2 = \phi_1 / (1 - \phi_2) = 6/7$$

We solve that :  $c_1 = 1/2 + (\sqrt{3}/14)i$ ,  $c_2 = 1/2 - (\sqrt{3}/14)i$

The general solution is given by:

$$\rho(h) = (1/2 + (\sqrt{3}/14)i)(1 + (\sqrt{3}/3)i)^{-h} + (1/2 - (\sqrt{3}/14)i)(1 - (\sqrt{3}/3)i)^{-h}$$

We use a loop to calculate the given recurrence relation.

```
loop <- function(n){  
  a <- 1  
  b <- 6/7  
  for (i in 2:n){  
    a_new <- b
```

```


    b <- 1.5*b - 0.75*a
    a <- a_new
  }
  return(b)
}

```

```

ACF<-c(1,6/7,loop(2),loop(3),loop(4),loop(5),loop(6),loop(7),loop(8),loop(9),loop(10))
X<-c(0:10)
plot(X,ACF,type = "h",col = "blue")
abline(h=0)

```



ARMAacf function to check the value we calculated.

```

ACF_r = ARMAacf(ar= c(1.5,-0.75),ma=0,10)
plot(ACF_r,type = "h",xlab = "lag",col = "red")
abline(h=0)

```

We use

