

The variance

Brian Caffo, Jeff Leek, Roger Peng

May 19, 2016

The variance

- ▶ The variance of a random variable is a measure of *spread*
- ▶ If X is a random variable with mean μ , the variance of X is defined as

$$\text{Var}(X) = E[(X - \mu)^2] = E[X^2] - E[X]^2$$

- ▶ The expected (squared) distance from the mean
- ▶ Densities with a higher variance are more spread out than densities with a lower variance
- ▶ The square root of the variance is called the **standard deviation**
- ▶ The standard deviation has the same units as X

Example


- ▶ What's the variance from the result of a toss of a die?
- ▶ $E[X] = 3.5$
- ▶ $E[X^2] = 1^2 \times \frac{1}{6} + 2^2 \times \frac{1}{6} + 3^2 \times \frac{1}{6} + 4^2 \times \frac{1}{6} + 5^2 \times \frac{1}{6} + 6^2 \times \frac{1}{6} = 15.17$
- ▶ $Var(X) = E[X^2] - E[X]^2 \approx 2.92$

Example

- ▶ What's the variance from the result of the toss of a coin with probability of heads (1) of p ?
- ▶ $E[X] = 0 \times (1 - p) + 1 \times p = p$
- ▶ $E[X^2] = E[X] = p$

$$\text{Var}(X) = E[X^2] - E[X]^2 = p - p^2 = p(1 - p)$$

Distributions with increasing variance



The sample variance

- ▶ The sample variance is


$$S^2 = \frac{\sum_{i=1} (X_i - \bar{X})^2}{n - 1}$$

(almost, but not quite, the average squared deviation from the sample mean)


- ▶ It is also a random variable
- ▶ It has an associated population distribution
- ▶ Its expected value is the population variance
- ▶ Its distribution gets more concentrated around the population variance with more data
- ▶ Its square root is the sample standard deviation

Simulation experiment

Simulating from a population with variance 1



Variances of x die rolls



Recall the mean

- ▶ Recall that the average of random sample from a population is itself a random variable
- ▶ We know that this distribution is centered around the population mean, $E[\bar{X}] = \mu$
- ▶ We also know what its variance is $Var(\bar{X}) = \sigma^2/n$
- ▶ This is very useful, since we don't have repeat sample means to get its variance; now we know how it relates to the population variance
- ▶ We call the standard deviation of a statistic a standard error

To summarize

- ▶ The sample variance, S^2 , estimates the population variance, σ^2
- ▶ The distribution of the sample variance is centered around σ^2
- ▶ The the variance of sample mean is σ^2/n
- ▶ Its logical estimate is s^2/n
- ▶ The logical estimate of the standard error is s/\sqrt{n}
- ▶ s , the standard deviation, talks about how variable the population is
- ▶ s/\sqrt{n} , the standard error, talks about how variable averages of random samples of size n from the population are

Simulation example

Standard normals have variance 1; means of n standard normals have standard deviation $1/\sqrt{n}$

```
nosim <- 1000  
n <- 10  
sd(apply(matrix(rnorm(nosim * n), nosim), 1, mean))
```

```
## [1] 0.3165173
```

```
1 / sqrt(n)
```

```
## [1] 0.3162278
```

Simulation example

Standard uniforms have variance $1/12$; means of random samples of n uniforms have sd $1/\sqrt{12 \times n}$

```
nosim <- 1000  
n <- 10  
sd(apply(matrix(runif(nosim * n), nosim), 1, mean))
```

```
## [1] 0.0936879
```

```
1 / sqrt(12 * n)
```

```
## [1] 0.09128709
```

Simulation example

Poisson(4) have variance 4; means of random samples of n

Poisson(4) have sd $2/\sqrt{n}$

```
nosim <- 1000  
n <- 10  
sd(apply(matrix(rpois(nosim * n, 4), nosim), 1, mean))
```

```
## [1] 0.6546695
```

```
2 / sqrt(n)
```

```
## [1] 0.6324555
```

Simulation example

Fair coin flips have variance 0.25; means of random samples of n coin flips have sd $1/(2\sqrt{n})$

```
nosim <- 1000  
n <- 10  
sd(apply(matrix(sample(0 : 1, nosim * n, replace = TRUE),  
                 nosim), 1, mean))
```

```
## [1] 0.1564943
```

```
1 / (2 * sqrt(n))
```

```
## [1] 0.1581139
```

Data example

```
library(UsingR); data(father.son);
```

```
## Loading required package: MASS
```

```
## Loading required package: HistData
```

```
## Loading required package: Hmisc
```

```
## Loading required package: lattice
```

```
## Loading required package: survival
```

```
## Loading required package: Formula
```


```
##
```

```
## Attaching package: 'Hmisc'
```

```
## The following objects are masked from 'package:base':
```

```
##
```


Plot of the son's heights



Let's interpret these numbers

```
round(c(var(x), var(x) / n, sd(x), sd(x) / sqrt(n)),2)
```

```
## [1] 7.92 0.01 2.81 0.09
```



Summarizing what we know about variances

- ▶ The sample variance estimates the population variance
- ▶ The distribution of the sample variance is centered at what its estimating
- ▶ It gets more concentrated around the population variance with larger sample sizes
- ▶ The variance of the sample mean is the population variance divided by n
- ▶ The square root is the standard error
- ▶ It turns out that we can say a lot about the distribution of averages from random samples, even though we only get one to look at in a given data set